

1-1-2007

# N-(4-Methylphenyl) Benzenepropanamide - the First Isolated Amide From the Genus *Paederia*

Debasish Bandyopadhyay

*The University of Texas Rio Grande Valley*, [debasish.bandyopadhyay@utrgv.edu](mailto:debasish.bandyopadhyay@utrgv.edu)

Anupam Nayak

Bidyut Basak

Avijit Banerji

Julie Banerji

*See next page for additional authors*

Follow this and additional works at: [https://scholarworks.utrgv.edu/chem\\_fac](https://scholarworks.utrgv.edu/chem_fac)

 Part of the [Chemistry Commons](#)

---

## Recommended Citation

Bandyopadhyay, Debasish; Nayak, Anupam; Basak, Bidyut; Banerji, Avijit; Banerji, Julie; Chatterjee, Asima; Prang, Thierry; and Neuman, Alain, "N-(4-Methylphenyl) Benzenepropanamide - the First Isolated Amide From the Genus *Paederia*" (2007). *Chemistry Faculty Publications and Presentations*. 59.

[https://scholarworks.utrgv.edu/chem\\_fac/59](https://scholarworks.utrgv.edu/chem_fac/59)

This Article is brought to you for free and open access by the College of Sciences at ScholarWorks @ UTRGV. It has been accepted for inclusion in Chemistry Faculty Publications and Presentations by an authorized administrator of ScholarWorks @ UTRGV. For more information, please contact [justin.white@utrgv.edu](mailto:justin.white@utrgv.edu), [william.flores01@utrgv.edu](mailto:william.flores01@utrgv.edu).

---

**Authors**

Debasish Bandyopadhyay, Anupam Nayak, Bidyut Basak, Avijit Banerji, Julie Banerji, Asima Chatterjee, Thierry Prang, and Alain Neuman

## *N*-(4-Methylphenyl) benzenepropanamide - the First Isolated Amide from the Genus *Paederia*

Debasish Bandyopadhyay<sup>a</sup>, Anupam Nayak<sup>a</sup>, Bidyut Basak<sup>a</sup>, Avijit Banerji<sup>a</sup>, Julie Banerji<sup>a\*</sup>, (Late) Asima Chatterjee<sup>a</sup>, Thierry Prangé<sup>b</sup> and Alain Neuman<sup>b</sup>

<sup>a</sup>Centre of Advanced Studies on Natural Products including Organic Synthesis, Department of Chemistry, Calcutta University, 92, A. P. C. Road, Kolkata – 700 009, India

<sup>b</sup>UMR 8015 CNRS 4, AV de l'Observatoire, 75006 Paris, France

juliebanerji@yahoo.co.in

Received: March 10<sup>th</sup>, 2007; Accepted: April 5<sup>th</sup>, 2007

Investigation of the stem of *Paederia foetida* (Rubiaceae) resulted in the isolation and characterization of *N*-(4-methylphenyl)-benzopropanamide, which was hitherto unknown as a natural product. This is the first report of an amide for the genus *Paederia*.

**Key words:** *Paederia foetida*, *N*-(4-methylphenyl)-benzopropanamide, Rubiaceae, X-ray crystallography.

*Paederia foetida* (Rubiaceae), locally called “Gandal” in India, grows in tropical parts of India, as well as in Central and Eastern Himalayas [1]. The stem of the plant has a reputation in folklore medicine and exhibits anti-inflammatory, anti-microbial, anti-oxidative and hepato-protective properties [2]. Several iridoid glycosides and lactones (paederinin, paederoside and paederia lactone) have been isolated from the stem of this plant [3]. The present article describes the isolation and characterization of an amide (**1**) from *P. foetida* stem; this compound was hitherto unknown as a natural product.

The ethereal extract of the stem was fractionated between *n*-hexane and ethyl acetate. The ethyl acetate fraction of the extract was subjected to column chromatography over silica gel (60-120 mesh). A white solid was isolated from the *n*-hexane-ethyl acetate (9:1) fraction, which on crystallization from methanolic *n*-hexane (2%) afforded a needle shaped microcrystalline compound (**1**).

The IR (KBr disc) spectrum of **1** revealed the presence of >NH and >C=O at 3299 and 1655 cm<sup>-1</sup>, respectively. The elemental composition of **1** could be ascertained as C<sub>16</sub>H<sub>17</sub>NO from the TOF mass spectrum. This exhibited a base peak at (m/z +Na) 262.1203, corresponding to C<sub>16</sub>H<sub>17</sub>NO, and a characteristic peak [M-C<sub>9</sub>H<sub>9</sub>O]<sup>+</sup> at m/z 107, from which the elemental composition C<sub>7</sub>H<sub>8</sub>N was determined by exact mass measurements (found: 107.9367). A fragment at m/z 91.8811 indicated the presence of a substituted toluene unit and another fragment at m/z 104.9208 indicated the presence of a Ph-CH<sub>2</sub>-CH<sub>2</sub> moiety. In the <sup>1</sup>H NMR spectrum, two triplets at δ 2.65 and δ 3.06 have been assigned to two adjacent methylenes, the latter being adjacent to a carbonyl group, as it resonated in a comparatively downfield region. A three-proton singlet at δ 1.71 was due to the *para*-methyl substituent at C-4 (Table 1). The <sup>13</sup>C NMR spectrum clearly showed the presence of sixteen carbon atoms in the molecule. Comparison of the <sup>13</sup>C NMR spectrum (fully decoupled) with DEPT 135<sup>0</sup> and DEPT 90<sup>0</sup> clearly indicated the presence of one methyl, two methylenes, nine aromatic methines and four quaternary carbons in the molecule, among which the signal at δ 170.24 was assigned to the carbonyl carbon. The NMR shifts of all the protons and





Figure 1: Structure of (**1**).

carbons have been depicted in Table 1. All these spectral data identify **1** as *N*-(4-methylphenyl)-benzopropanamide (Figure 1).

**Table 1:** NMR spectroscopic shifts of the amide (**1**) in CDCl<sub>3</sub>.

Position	<sup>1</sup> H (δ)	<sup>13</sup> C (δ)
2	—	170.24
3	3.1(t, 2H, <i>J</i> = 7.6 Hz)	39.42
4	2.65 (t, 2H, <i>J</i> = 7.6 Hz)	31.59
5	—	140.71
6, 10	7.1-7.4 (m, 10H)	128.60
7, 9	7.1-7.4 (m, 10H)	128.38
8	7.1-7.4 (m, 10H)	126.34
1'	—	133.95
2', 6'	7.1-7.4, (m, 10H)	120.03
3', 5'	7.1-7.4 (m, 10H)	129.41
4'	7.1-7.4 (m, 10H)	135.15
7'	1.71 (3H, s)	20.81

Finally the structure was confirmed by X-ray crystallographic analysis. The ORTEP projection is shown in Figure 2.



**Figure 2:** ORTEP projection of **1**.

In conclusion, this investigation has resulted in the first report of the isolation of *N*-(4-methylphenyl)-benzenepropanamide as a natural product.

## Experimental

The melting point was determined in an electrically controlled melting point apparatus (Sunvic, UK). The IR spectrum was recorded using a Perkin Elmer RX1 FT IR spectrophotometer. The <sup>1</sup>H and <sup>13</sup>C NMR spectra were obtained in CDCl<sub>3</sub> in a Bruker AVANCE 300 DIGITAL MHz NMR spectrometer.

## References

- [1] Chatterjee A, Pakrashi SC. (1997) *The Treatise on Indian Medicinal Plants*, Vol. 5, National Institute of Science Communication, New Delhi p. 94.
- [2] (a) De S, Ravishankar B, Bhavsar GC. (1994) Investigation of the anti-inflammatory effects of *Paederia foetida*. *Journal of Ethnopharmacology*, **43**, 31-38; (b) Srivastava MC, Tewari JP, Kant V. (1973) Anti-inflammatory activity of an indigenous plant-*Paederia foetida*. *Indian Journal of Medical Sciences*, **27**, 231-234.
- [3] Zou X, Liang J, Ding LS, Peng SL. (2006) Studies on chemical constituents of *Paederia scandense*. *Zhongguo Zhong Yao Za Zhi*, **31**, 1436-1441.

Chemical shifts were reported as δ (ppm) and the coupling constants (*J*) were given in Hz. The mass spectrum was recorded on a Qtof Micro YA 263 mass spectrometer. The X-ray crystallographic data were recorded on a Nonius CCD diffractometer operating the MoKα radiation (λ = 0.7107 Å). The crystals are monoclinic, space group P2<sub>1</sub>/c, with parameters a = 14.706(3); b = 4.870(4); c = 18.965(4) Å, β = 98.14(4)°, and refined to a R = 8.9 % (for all 1468 data). Crystallographic Data for compound **1** i.e. *N*-(4-methylphenyl)-benzopropanamide have been deposited with the Cambridge Crystallographic Data Centre as CCDC 643303. Copies of the data can be obtained, free of charge, on application to CCDC, 12 Union Road, Cambridge CB2 1EZ, UK (fax: +44 1223 336033 or e-mail: deposit@ccdc.cam.ac.uk).

**Plant material:** The stems of *P. foetida* were collected from the Ministry of Food and Supplies, Government of India, and compared with a voucher specimen maintained in the herbarium of Calcutta University.

**Extraction and Isolation:** The stem (0.53 Kg) was extracted at room temperature with ether. The ethereal fraction was concentrated under low pressure and a dark brown mass (22 g) was obtained. This was column chromatographed over silica gel (60-120 mesh) and eluted successively with *n*-hexane and *n*-hexane-ethyl acetate mixtures. The fraction obtained with *n*-hexane-ethyl acetate (9:1) afforded a white solid, which on repeated recrystallization using methanolic *n*-hexane afforded pure (57 mg) compound **1** as white needle-shaped crystals (m.p. 128°C).

**Acknowledgments** - The authors thank Mr Pitambar Ghosh and Mr Jagadish Ghosh for NMR and IR spectral measurements, respectively and to the University of Calcutta for financial assistance (Khaira endowment research fellowship to A. N.).