

Genetic instability of *Acetobacter pasteurianus* revealed by whole genome analyses

Yoshinao Azuma

Abstract

Acetobacter species have been used for brewing traditional vinegar and are known to have genetic instability. To clarify the mutability, *Acetobacter pasteurianus* NBRC 3283, which forms a multi-phenotype cell complex, was subjected to genome DNA sequencing. The genome analysis revealed that there are more than 280 transposons and 5 genes with hyper-mutable tandem repeats as common features in the genome consisting of a 2.9 Mb chromosome and 6 plasmids. There were 3 single nucleotide mutations and 5 transposon insertions in 32 isolates from the cell complex. The *A. pasteurianus* hyper-mutability was applied for breeding a temperature resistant strain grown at an unviable high-temperature (42°C). The genomic DNA sequence of a heritable mutant showing temperature resistance was analyzed by mutation mapping, illustrating that a 92 kb deletion and three single nucleotide mutations occurred in the genome during the adaptation. Alpha-proteobacteria including *A. pasteurianus* consists of many intracellular symbionts and parasites, and their genomes show increased evolution rates and intensive genome reduction. However, *A. pasteurianus* is assumed to be a free-living bacterium, it may have the potentiality to evolve to fit in natural niches of seasonal fruits and flowers with other organisms, such as yeasts and lactic acid bacteria.

Footnotes

The eight substrains of *A. pasteurianus* (IFO 3283-01, 03, 07, 12, 22, 26, 32 and IFO 3283-01-42C) reported herein have been re-deposited in the National Biological Resource Center (NBRC) and its genome DNA sequences were registered in DDBJ/EMBL/GenBank database. The accession numbers of NBRC and DDBJ are shown in **Table 1** and **Table 2**, and in **Supplemental Table S1** in detail. An accession number for the largest plasmid of *Gluconacetobacter xylinus* NBRC 3288, pGXY010, is BABN01000001. Supplemental data including a gene list of *A. pasteurianus* IFO 3283-01 and the color version of the figures in this paper may be presented in my web site: <http://web.me.com/yoshinaoazuma>.

Key words: evolution, genomics, acetic acid bacteria, *Gluconacetobacter xylinus*, adaptation

Abbreviations: IFO, Institute for Fermentation, Osaka; NBRC, National Biological Resource Center of Japan; SNP, single-nucleotide polymorphism; DAPI, 4'-6-diamidino-2-phenylindole; AAB, acetic acid bacteria; bp, base pair; kb, kilo base pair; aa, amino acid; nt, nucleotide; TLS, translesion synthesis.


Received 22 Nov. 2010

Department of Science and Technology on Food Safety, Kinki University, Wakayama 649-6493, Japan

Introduction

Acetic acid bacteria (AAB) is a widely divergent group within the alpha-proteobacteria and are isolated from a variety of natural fields such as fruits, flowers and fermented foods, and are rarely found in soils and insect guts (1-3). Recently, one species was identified as a human pathogen (4). Its metabolic uniqueness was also traditionally utilized to generate fermented food, especially vinegar, and also for industrial production of sorbose and dihydroxyacetone (5). AAB plays a crucial role in the quantity, taste, nutrition and hygienic quality of the productions, thus the reliability of strains in terms of functional properties and growth robustness is indispensable. AAB has a noticeable physiological instability and it is reported in at least two different terms of temporal acclimation and heritable adaptation (6-9).

Introductory Figure.



Phylogenetic tree of life and acetic acid bacteria based on small ribosomal RNA sequences

Acetobacter strains, which are among the most popular AAB for the production of vinegar in many countries, may gradually acquire resistance against higher concentrations of acetic acid when properly adapted to the conditions. However, the rapid loss of the acquired phenotype is observed when the cells are maintained in environments without acetate as a selection pressure (7,8). This acquisition of acetic acid resistance must be an example of the temporal acclimation or physiologic adaptation, and one reason why new tanks for fermentation are continuously inoculated with AAB from old tanks, but not from a preserved seed AAB in traditional vinegar production.

Heritable deficiencies in various physiological properties of *Acetobacter* strains, such as ethanol oxidation, acetic acid resistance and bacterial cellulose synthesis, are observed at high frequencies. Very little is known about the genetic background for the instability but phenotypic modifications by transposon insertion were reported in ethanol oxidation and acetic acid resistance (6,9), and cellulose formation (10). Relatively copious transposons were identified in the genome of AAB strains, such as *Gluconobacter oxydans* 621H (11) and *Granulibacter bethesdensis* CGDNIH1

(12).

Genomic mutations are fundamental phenomena in the evolution of any organism. Bacteria may be more directly influenced by any mutations because of its minimized genomes or the density of information in its genomes. New information about the genomic structures and mutational events of bacteria have clarified various systems of the genomic mutations, such as a horizontal gene transfer of mobile gene units (e.g. transposon, plasmid and phage), hyper-mutable tandem repeat, genome-wide rearrangement and genome reduction (13), adding to the well known mutation systems, such as nucleotide substitution, insertion and deletion, and gene duplication (14).

In a harmful environment, mutations could provide beneficial alleles, but most mutations are likely to be neutral or deleterious, at least slightly, to organisms. Bacterial genomes accumulate mutations at certain rates even in well-fitted environments, and a natural environment causing stresses and stimuli may elevate the mutation rates (15,16). Therefore, DNA repair and proofreading systems must have evolved to minimize the rates of mutations and natural selection acts on the reliable maintenance and transmission of genome information (17).

Genome modulation for survival under certain stressful environments may cause other types of mutation, i.e. genome wide rearrangement and genome reduction. Under nutrition depleted conditions, genome wide rearrangements were experimentally observed with a certain frequency in *Salmonella enterica* (18) and *Lactococcus lactis* (19), suggesting the existence of a biological system acquired in evolution due to repeatedly facing harmful environments. Extreme genome reduction is mostly observed in endosymbionts and obligate intracellular pathogens (20-23), but also in a free-living photosynthetic marine bacterium, *Prochlorococcus marinus* (24). The genetic variation of symbiont genomes in comparison to free-living bacteria illustrate a large scale of gene loss that includes genes for regulation of transcription, translation and replication as well as genes for intermediate metabolism and amino acid biosynthesis.

In association with adaptation to a new niche, unnecessary genes without significant functions may accumulate mutations and may be deleted from the genomes. One explanation for genome reduction was derived from a comparative genome analysis between *Buchnera aphidicola* and *E. coli* that showed large gene deletions by chromosome rearrangements and multiple events of disintegration dispersed over the whole genome (25,26). However, the mechanism associated with genome reduction or shrinkage is still controversial, with regard to whether a wide deletion may occur as a single event or as an accumulation of smaller deletions. A previous chlamydial genome analysis predicted the replication terminal region within the plasticity zone was involved in genome reduction according to enrichment of species specific, hypothetical and truncated genes (23).

The complete genomic DNA sequences of *Acetobacter pasteurianus*, which has been used for traditional vinegar productions, was herein determined to characterize the hyper-mutability of AAB, using a multi-phenotype cell complex formed by 21 years maintenance without crucial stresses. The strain that adapted to environments above the growth-limiting temperature was also subjected to genome analyses.

Materials and Methods

Strain and Culturing

IFO records (maintained in NBRC to date) show that *Acetobacter* spp. NBRC 3283, formerly assigned as *Acetobacter aceti* IFO 3283, was originally isolated from a pellicle (a kind of biofilm) on the surface of vinegar fermentation. The 16S rRNA sequences of the *Acetobacter* spp. NBRC 3283 in this work are all identical to one of *Acetobacter pasteurianus* but not to *A. aceti* or any other *Acetobacter* strains. Therefore, the nomenclature of this strain was taxonomically reassigned as *A. pasteurianus* in this paper. Different lots of *A. pasteurianus* NBRC 3283 were distributed and maintained as a typical *Acetobacter* strains for research. An NBRC 3283 stock sample for genome analyses was the closest to an original IFO 3283 but contained several substrains. To avoid any nomenclature confusion, the substrains isolated from the IFO 3283 (NBRC 3283) were designated, such as *A. pasteurianus* IFO 3283-01.


The *A. pasteurianus* IFO 3283 (NBRC 3283) and *Gluconacetobacter xylinus* NBRC 3288 were obtained from National Biological Resource Center of Japan (NBRC) and grown routinely at 30°C on YPG medium 1.0% yeast extract, 1.0% polypeptone and 2.0% glycerol. A rich medium, YPGD, contains 1.0% yeast extract, 1.0% polypeptone, 2.0% glycerol and 0.5% glucose, and a minimum medium, NCG, contains 0.5% yeast nitrogen base (BD Difco, Franklin Lakes, NJ), 0.5% casamino acid (BD Difco) and 2.0% glycerol. Potato solid medium was prepared with potato extract from 200 g of boiled potato per liter, 3.0% yeast extract (BD Difco), 0.5% meat extract (Wako Pure Chemical Industries, Osaka, Japan), 1.0% thioglycolate medium (Wako), 0.5% glucose, 1.5% glycerol, 1.5% CaCO₃,

1.5% agar (pH 7.0). Other media and supplementations, such as ethanol and acetic acid, are described in each figure legend. Short-term continuous cultivation was performed by a dilution method, i.e. when the optical density (OD600) of a culture reached approximately 3.0, the culture was diluted to OD600 = 0.1 with 100 ml of fresh medium in a 300 ml flask and continuously incubated at 30°C with aeration by rotation at 120 rpm.

Between 1954 and 1974, *A. pasteurianus* IFO 3283 (NBRC 3283) was maintained by a serial passage every three months at the Institute for Fermentation, Osaka (IFO). In each passage, a loaf of cells picked from a former slant tube stored at 5°C was spread onto the potato solid medium in a new tube and incubated at 28°C for 4 days to allow the cells to grow enough for the next passage, followed by storage at 5°C for 3 months. Reconstitution of the *A. pasteurianus* maintenance herein was performed with the potato medium within slant tubes of 18 mm in diameter in a manner similar to that carried out between 1954 and 1974 in IFO.

Escherichia coli JM109 was grown routinely at 37°C on LB medium containing 1.0% NaCl, 1.0% bacto tryptone and 0.5% yeast extract.

Methodological Figure



DNA Sequencing, assembling gap closing and variation detection

Three DNA shotgun libraries of *A. pasteurianus* IFO 3283 (NBRC 3283) complex were constructed with inserts of 1.5 and 5.0 kb in pUC118 vector (Takara, Otsu, Japan) and 35 kb in the pCC1FOS (Epicentre, Madison, WI) fosmid vector. The plasmid clone DNAs were directly amplified from *E. coli* colonies and fosmid clone DNAs were extracted from *E. coli* transformants using the Montage BAC96 MiniPrep Kit (Millipore, Billerica, MA), respectively, and DNAs were end-sequenced using dye-terminator chemistry on an ABI PRISM3730XL sequencer (ABI, Foster, CA), as described previously (27). The 42,048 raw sequence reads corresponding to approximately 9.3-fold coverage were assembled using the PHRED/PHRAP/CONSED software package (<http://www.phrap.org>) (28-30). A number of repeat regions detected in the assembling were masked and the sequences in the most of the masked regions were determined by primer walking to assemble the sequences into supercontigs. Seven supercontigs corresponding to a chromosome and 6 plasmids were constructed with remains of 30 gaps and 84 low quality regions under 40 in Phrap score. DNA fragments were then produced to fill all the gaps, except for one locus, by PCR based on the supercontig. The one locus with a few similar transposons was independently sequenced using a fosmid covering the region and Template Generation System II Kit (Finnzymes, Espoo, Finland). All regions with sequences of lower quality than 25 in Phred value were resequenced by primer walking. Assembling of genome DNA was validated by three methods, i.e., Pulse field gel electrophoresis (Bio-Rad, Hercules, CA) after genomic DNA digestion by *NotI*, *SfiI* and *FseI*, fosmid clone digestion by several restriction enzymes and an Optical Mapping system (OpGen, Madison, WI). When any inconsistency in assembling and coexistence of different nucleotides were detected, the sequence variations were confirmed by resequencing all 32 isolated clones.

The draft genomic DNA sequence of *Gluconacetobacter xylinus* NBRC 3288 was obtained by similar methods as described above, and the DNA sequence of the largest plasmid pGXY010 was analyzed in this work.

For the mutation mapping of *A. pasteurianus* IFO 3283-01-42C, which is bred from IFO 3283-01 by adaptation to 42°C, DNA sequencing was performed with the Genome Analyzer (Illumina, San Diego, CA). After the text editing of approximately one billion of raw sequence data, mutation mapping was performed using the MAQ software (31). All candidates of mutation sites were directly resequenced using PCR products as templates by dye-terminator chemistry on an ABI PRISM3730XL sequencer (Invitrogen ABI, Carlsbad, CA).

Genome DNA sequence analysis and annotation.

Gene identification was performed with a method described previously (23). rRNA and tRNA genes were predicted using the BLASTN (32) and tRNAscan-SE (33) programs, respectively. Open reading frames (ORFs) longer than 90 bp were first predicted using the combination of three programs, GenomeGambler (34), GeneHacker plus (35) and Glimmer 3.0 (36), and at last protein-coding genes manually selected out of the ORFs. Annotation for each gene was carried out by using the BLASTP (37) and FASTA3 (38) programs against the non-redundant protein database. Transmembrane protein genes were predicted using SOSUI (39). The genes and the annotations of IFO 3283-01 are listed in **Supplemental Table S2**.

Hyper-mutable tandem repeats (HTRs or short tandem repeats) were extracted using the JSTRING software, (40), on genomic DNA sequences from *A. pasteurianus* IFO 3283 (NBRC 3283) substrains, *Gluconobacter oxydans* 621H genome (11), *Gluconacetobacter diazotrophicus* PAL5 (GENBANK: NC-010123, NC-010124, NC-010125), *Acidiphilium cryptum* JF-5 (GENBANK: from NC-009467 to NC-009474, and NC-009484), and *Granulibacter bethesdensis* CGDNIH1 (12) and a *G. xylinus* plasmid, pGXY010 (BABN01000001 in this work).

Accession numbers of *A. pasteurianus* IFO 3283 (NBRC 3283) substrains.

Thirty-two isolates from the *A. pasteurianus* IFO 3283 (NBRC 3283) cell complex stored in 1974 were classified into seven major groups. Seven substrains, one from each group, i.e. IFO 3283-01, IFO 3283-03, IFO 3283-07, IFO 3283-12, IFO 3283-22, IFO 3283-26 and IFO 3283-32, were re-deposited and are available from National Biological Resource Center of Japan (NBRC; <http://www.nbrc.nite.go.jp/e/index.html>; **Table 1**). The complete genomic DNA sequences of the 7 strains including 6 plasmids are registered at GenBank/EMBL/DDBJ. The accession numbers are listed in **Table 1** and **Table 2** and in **Supplemental Table S1** in detail. The accession number for the largest plasmid, pGXY010, of *G. xylinus* NBRC 3288 is BABN01000001. The adapted strain, IFO 3283-01-42C, was re-deposited to NBRC and are available as NBRC 105185. The accession numbers of the genomic DNA sequence for IFO 3283-01-42C are listed in **Supplemental Table S1**.

High-throughput measurement of the total amount of neutral sugar.

An amount of total neutral sugar as the polysaccharide components were measured based on a Phenol-H₂SO₄ method (41). By this method, ribose and deoxyribose are detectable as well as glucose, mannose, galactose, and fucose, thus the amounts of nucleic acid were compensated. For a liquid culture condition, *A. pasteurianus* was grown to approximately OD₅₉₅ = 1 in the NCG medium and the culture was diluted with 0.85% NaCl solution to OD₅₉₅ = 0.1. For plate culture conditions, *A. pasteurianus* was grown for 3 days on the YPG plate medium and the cells were harvested and diluted with 0.85% NaCl solution to OD₅₉₅ = 0.1. The amounts of soluble and insoluble (retained with cells) neutral sugars were measured using the supernatants and cell pellets after centrifugation of the cell suspensions at 10,000 x g for 5 min at 4°C. Briefly, 20 µl of samples in a well of a 96-well titer plate was mixed with 20 µl of 5% phenol and then 100 µl of H₂SO₄ was added to the mixture. The absorbance of the mixture was measured at 485 nm using a micro-titer plate reader (Beckman, Fullerton, CA). A glucose solution was used for a concentration standard. The amount of nucleic acid in the sample was determined by measurement of the absorbance at 260 nm after mixing 126 µl of samples with 14 µl of 5% SDS solution in a well of a UV transparent multi-titer plate (Corning, New York, NY). Double strand DNA extracted from *A. pasteurianus* using a Genomic DNA extraction kit (Qiagen, Germantown, MD) was used as a concentration standard. The amount of protein in the sample was measured using a Protein assay kit (Bio-Rad, Hercules, CA) by absorbance at 595 nm after mixing 40 µl of the diluted cell suspension, 40 µl of 1% SDS solution and 20 µl of Protein assay solution in a well of a multi-titer plate. Bovine serum albumin was used as a concentration standard. Because DNA in the sample affects the measurements of total sugar and protein concentration, the amounts of DNA were estimated and subtracted from the total sugar and protein measurements, and then the sugar production per protein was calculated.

High-temperature adaptation and mutation locus analysis.

To attempt to breed a thermo-resistant *A. pasteurianus* strain, *A. pasteurianus* IFO 3283-01 cells, which exhibit less aggregation and produce sugar in liquid culture, were incubated at 40°C and the culture at approximately OD₆₀₀ = 3 was inoculated into 200 ml of fresh YPGD liquid medium pre-warmed at 40°C to be OD₆₀₀ = 0.1. After 4 passages, the culturing temperature was shifted from 40°C to 42°C. After a temperature tolerant strain was produced, serial

cultures at 42°C were still continued 50 times to confirm the viability at that temperature.

Genomic DNA was prepared from clones (40C(3)1 and 40C(5)1) out of the third and fifth cultures at 40°C, respectively, and from clones (42C(1)9, 42C(11)1, 42C(21)1 and 42C(30)1) the first successful culture at 42°C cultivation and 11, 21, 30 succeeding cultures at 42°C, respectively. The genome DNAs were analyzed by 1% agarose gel electrophoresis to check small size plasmids and hyper-mutable tandem repeat sequences, and by pulse field gel electrophoresis to study macro scale genome structures.

Five µg of genomic DNAs from the IFO 3283-01-42C, which is a clone from the first successful culture at 42°C, were sequenced using a Genome Analyzer (Illumina inc. San Diego CA) following the instruction of Post Genome Ins. (Tokyo, Japan). Two databases were constructed using the 10.0 million sequence reads obtained from the sequencing process and 2.3 million sequence reads with high qualities (at least 20 bases with Solexa score 25 or more in a read). These sequence read sets were separately mapped onto genome sequences of non-parental clones, such as *A. pasteurianus* IFO 3283-02, to evaluate whether the known mutation loci were identified, thus indicating that those data sets contained sufficient quality and quantity to extract mutation candidates. After extraction of mutation candidates using the genomic DNA sequence of the parental clone, IFO 3283-01, mutation sites were confirmed by PCR and DNA sequencing.

Results and Discussion

Multi-phenotype cell complex.




Fig 1. Mutation accumulation within maintenance of *A. pasteurianus* IFO 3283 for 21 years.

a) A schematic illustration of the maintenance and generations from 1954 to 1974. The bottom panel indicates cell multiplication for the first 5 days up to 6.7 generations (a solid line) and decreased cell viability during the slants storage at 5°C for 85 days (a broken line) in a passage of every 3 months. Gs in the panel mean generation numbers. b) Phenotypic variations in the IFO 3283 (NBRC 3283) cell complex stored in 1974. The top photo shows at least two types of colonies from multi-phenotype cell complex. Two independent clones, (R; rough colony and S; smooth) are shown in the two sets of bottom photos, the transparency (upper) and texture of the surface (middle) of colonies. The bottom panel shows that R type clones produce a pellicle (biofilm) on the surface of liquid culture, but not S clones.

Acetobacter pasteurianus IFO 3283 (NBRC 3283) was maintained by a series passages every three months since the deposit of the pure culture at Institute for Fermentation, Osaka (IFO) in 1954 until the establishment of the freeze-dry preservation method in 1974 (Fig. 1a). Since no colony isolation was performed in the maintenance to avoid the loss of the useful features of the strain, a multi-phenotype cell complex was formed characterized by different textures of rough (R) and smooth (S) colony surfaces, which may be based on a polysaccharide consisting of neutral sugars and a mucoid slime layer, respectively (42) (Fig. 1b). The proportions of the rough and smooth colonies were 57% and 43%, respectively.

To estimate the generations in the 21 years of maintenance, the three-month slant passage was reconstituted in this work using the potato medium. Within the first 4-day culturing in slant tubes, the cell number increased 101.2 folds, which is 6.7 generations (standard deviation (SD); 0.71). However, cell viability decreased to 0.88×10^{-6} (SD; 4.1)

after 85 days storage in 5°C and each survival cell need to increase 0.88×10^6 folds ($\times 2^{20.1}$) to recover the original amount of cells. Together, cells needed to multiply $\times 2^{20.1}$ and $\times 2^{6.7}$, corresponding to $\times 2^{26.7}$ or 26.7 generations, in each passage (Fig. 1a). Consequently, it was estimated that 26.8 generations per passage every 3 months and thus 2.2×10^3 generations occurred within the 21 years, accompanying the formation of the multi-phenotype cell complex.

A. pasteurianus genome DNA sequence and its variations.

Shotgun DNA sequencing of the *A. pasteurianus* IFO 3283 (NBRC 3283) genome was carried out using mixed genomic DNA extracted from the cell complex stored at IFO in 1974. After the assembly of the sequence reads with 9.3-fold coverage, an approximate 2.9 Mbp supercontig and 6 short contigs were constructed corresponding to the chromosome and 6 plasmids, respectively. Thirty-eight assembling gaps and sequence discrepancies were identified as candidate loci for genome variations, i.e. 30 gaps by transposon insertions, 5 possible sites of single nucleotide polymorphisms (SNPs) and 3 regions with hyper-mutable tandem repeats (HTRs).

The DNA sequences of the mutation candidates were completely determined using PCR products amplified from genomic DNAs of the 32 isolates (serially named from IFO 3283-01 to IFO 3283-32) from the IFO 3283 (NBRC 3283) cell complex, except one gap in the largest plasmid was sequenced by serial terminal deletions after cloning of the corresponding region from the IFO 3283-01. The results showed that 4 transposon insertions (numbered as Gap03, Gap14, Gap18, Gap22), 3 SNPs (SNP01, SNP02 and SNP03) and 3 HTRs (AP-TR01, AP-TR03 and AP-TR04) were identified as chromosome variations and a transposon insertion (Gap107) and a HTR (AP-TR07) in the largest plasmid were found as a plasmid variation (Table 1). In the other 25 loci of the gaps, sequencing using long genome fragments shows that there are a few similar transposons but no genome variations in each locus. The complexity of transposons might prevent the program to assemble the loci with shotgun sequence data. Neither nucleotide substitutions nor insertion/deletion were found in the other two SNP loci out of the five candidates.

The mutation rate of non-mutator *E. coli* (chromosome size; 4.64 Mb) in the exponential phase was previously estimated as $2 - 8 \times 10^{-4}$ per generation (43-45). When the rate is adapted to the *A. pasteurianus* (chromosome size; 2.9 Mb), its mutation rate may be $1.25 - 5.0 \times 10^{-4}$ per generation. Therefore the mutation amount are predicted by multiplication of the mutation rate and the generation numbers, resulting in 0.3 - 1.1 mutations within the 2.2×10^3 generations in the 21 years. Since the actual chromosome mutation number is 8 (3 SNPs and 5 transposon insertions) and *A. pasteurianus* seems to accumulate mutations faster than the non-mutator *E. coli*. The “experienced” hyper-mutability of *A. pasteurianus* might be explained by the high rate of mutation (6). It is also reported that the mutation rate of *E. coli* in the starvation or “stationary phase” increases up to 10 to 100 times higher than that in the exponential phase (46). However, *A. pasteurianus* cells were allowed to grow on a rich medium during the maintenance, 85 days storage in 5°C remarkably decreased the cell viability to 0.88×10^{-6} . This suggests that *A. pasteurianus* cells might be sensitive to the nutrition depletion and that increases the mutation rate.

General features of *A. pasteurianus* genome

Table 2. General features of *A. pasteurianus* IFO 3283-01 chromosome and plasmids.

*1, IFO 3283-01-42C cultured at 30°C

| IFO 3283-01 | Name | DDBJ accession No. | Size | Copy *1 | GC(%) | rRNA | tRNA | ORFs |
|-------------|------------|--------------------|-----------|---------|-------|-----------|------|------|
| Chromosome | APA01 | AP011121 | 2,907,495 | 1 | 50.7 | 5 operons | 57 | 2628 |
| Plasmid | pAPA01-011 | AP011122 | 191,799 | 0.8 | 59.3 | 0 | 0 | 178 |
| | pAPA01-020 | AP011123 | 182,940 | 0.9 | 53.5 | 0 | 0 | 174 |
| | pAPA01-030 | AP011124 | 49,961 | 0.8 | 53.9 | 0 | 0 | 63 |
| | pAPA01-040 | AP011125 | 3,204 | 7.4 | 53.7 | 0 | 0 | 3 |
| | pAPA01-050 | AP011126 | 3,035 | 19.1 | 55.8 | 0 | 0 | 2 |
| | pAPA01-060 | AP011127 | 1,815 | 29.1 | 58.6 | 0 | 0 | 2 |

The putative replication origin (*ori*) of the chromosome was predicted based on the GC skew (47) and the coding direction of genes (48) (Fig. 2a). There is a relatively strong correlation between the directions of transcription and replication. The replication terminal was not specifically determined, but may be located in the region (*ter* region) between 1.25 and 1.55 Mbp from the *ori*. The chromosome of IFO 3283-01 contains 2,628 protein-coding genes, 57 tRNA genes and five copies of ribosomal RNA operons and plasmids encode 422 protein-coding genes (Table 2, Fig.2b and details in Supplemental Table S2).

Specific functions were assigned to 75.1% (2291 genes) of the total 3050 protein-coding genes, and 24.9% were hypothetical genes. A remarkable number of genes, 285 (9.0%), were found to encode transposases (Fig. 2b and Supplemental Table S3). A total 1836 orthologous genes between chromosomes of *A. pasteurianus* IFO 3283-01 and *Gluconobacter oxydans* 621H (11) are linked but a number of fragmented short regions show synteny between these species (Fig. 2a). Locations of genes categorized into several groups are shown in Figure 2b. Notable biases of gene density were found in the groups of transport and phage related genes.




Fig 2. Genome map and structure of *A. pasteurianus* IFO 3283-01 chromosome

a) The origin of replication was determined based on three values from cumulus of the GC (blue) and keto (green) skews, and transcription directions (red), and shown as *ori* (47). The locations of orthologues are shown by linking between *A. pasteurianus* and *G. oxydans* (11). b) Genes in the chromosome are mapped after gene categorization into 9 groups such as cell structure, replication and so on. Transcription and stress are indicated in black and red, respectively. Genes in translation are separated into protein coding in blue, and 52 tRNA in green and 15 rRNA in red. Numbers next to the categories are the number of genes in the chromosome and whole genome on the left and right, respectively. Genes in "common" of metabolism means that its orthologues are found in both genomes of *Gluconobacter oxydans* (11) and *Granulibacter bethesdensis* (12). The mutation loci identified in this work are also mapped. Genome variation by 3 SNPs in red, 4 transposon insertions in blue and 5 HTRs in green, and terminal deletion in red of the 42°C evolved strain.

Metabolism

Acetobacter aceti IFO3281 utilizes glycerol as the best carbon source for growth (49), and glycerol was the best carbon substrate for *A. pasteurianus* as well under our experimental conditions (data not shown). Metabolic cascades of glycerol utilization was not unveiled yet by experimental methods, but it may be predicted by the results from *Acetobacter xylinus* (renamed to *Gluconacetobacter xylinus*) (50,51) and the gene contents of *A. pasteurianus* in this

work, and the predicted metabolic pathways were illustrated in the **Supplemental Figure S1**. There are generally two pathways to produce dihydroxyacetone (DHA) phosphate from glycerol, one is via DHA catalyzed by glycerol dehydrogenase and the other is via glycerol 3-phosphate by glycerol kinase. The DHA phosphate may be converted to D-glyceraldehyde 3-phosphate by triosephosphate isomerase and thus enter into the glycolysis and gluconeogenesis pathway. However *A. pasteurianus* genome (as well as *G. xylinus*, data not shown) contains no genes encoding membrane-bound glycerol dehydrogenase which is well characterized in *Gluconobacter* (52), the capability to produce both DHA and glycerol 3-phosphate by *G. xylinus* (50,51) were reported and *A. pasteurianus* genome includes the glycerol kinase gene (described later) and several genes for alcohol dehydrogenase genes, such as sorbitol dehydrogenase. Therefore *A. pasteurianus* might promote glycerol utilization in both pathways. *A. pasteurianus* genome does not carry two genes encoding glucose-6-phosphate isomerase and pyruvate ferredoxin oxidoreductase in the glycolysis pathway. It suggests that nucleotide sugars metabolism and glycolysis may be connected through pentose phosphate pathway (11). There are no genes encoding succinyl-CoA synthetase in the citrate cycle shown previously and the process might be substituted with the acetic acid resistance gene product, AarC (53). KEGG based metabolic maps may be postulated in my web site, <http://web.me.com/yoshinaoazuma>.

Lineage of genome variation.

Unlike chromosome variations in diploid eukaryotes, mutations in a bacterial chromosome may conserve the order, in which the mutations occur in chromosomes. Chromosome variations observed in *A. pasteurianus* genomes, 3 SNPs and 4 transposon insertions were analyzed to classify a generation order of variations and a strain lineage (**Table 1** and **Fig. 3a**). HTRs were not considered for this purpose because the repetitive numbers of HTRs were modified even in the process of PCR-directed DNA sequencing.

SNP01 at 746 nucleotide (nt) of *rpoA* gene (APA01-09550) shows alterations of T or A among the 32 isolates, which resulted in amino acid alterations of leucine or glutamine, respectively. Multiple alignment with other alpha-proteobacteria homologues suggested that leucine is the ancestral amino acid and thus T746 must be mutated to A. Similarly G144 at SNP02 in glycerol kinase gene (APA01-12552) might be an ancestral one and mutated to T. Alteration of nine and eight adenine at SNP03 generates different coding frames to encode 604 and 90 aa in downstream of the SNP03, respectively. However, no significantly similar proteins to both 604 and 90 aa were found in non-redundant database, the frame with 604 aa is more likely an ancient type because no overlapping genes were observed on the coding region and the coding sequence does not contain a significant amount of minor codons, such as AGG and CTA.

When the corresponding genes (alleles) at chromosome variations associated with transposon insertions (Gap03, Gap14, Gap22 and Gap18) were compared among the 32 strains, all transposons were inserted in a same site in each allele, and no insertion or deletion were observed in alleles without transposons. This suggests that each transposon insertion happened in a certain strain and inherited, and no removing occurred.

More assumptions were added for the lineage analysis of the chromosome variation, (1) no reversible mutation occurred, (2) no chromosome recombination happened between independent strains, and (3) when a wider range of strains with a certain mutation incorporates a smaller one with a different mutation, the former mutation may happen earlier than the latter one. For example, Gap22 is observed in IFO3238-01, 05, 09, 10, 13, 16, 07, 06, 11, 14 and 15, but Gap03 in only IFO3238-01, 05, 09, 10, 13 and 16. In this case, Gap22 was thought to happen earlier than Gap03 (**Table 1**).

Based on these assumptions, the 32 isolates were classified into 7 groups in the order shown in **Figure 3a**. One transposon insertion (Gap107) in the largest plasmid was excluded from the categorization of the genome variations, because plasmids are possibly transmittable among the independent cells. However, no transmission of the plasmids was observed between different genome groups in this work. One representative clone each was chosen from the 7 groups, i.e. IFO 3283-01, IFO 3283-03, IFO 3283-07, IFO 3283-12, IFO 3283-22, IFO 3283-26, and IFO 3283-32 (**Table 1**). IFO 3283-32 is the only strain without mutations, which may retain the original genotype, and may also show the original phenotype of *A. pasteurianus* NBRC 3283. The substrains carrying no (or a few) mutations were minor in the IFO 3283 cell complex.




Fig 3. Mutation accumulation within maintenance of *A. pasteurianus* IFO 3283 for 21 years.

a) Genotype lineage (chromosome) and phenotypes of 32 substrains of the IFO 3283 multi-phenotype complex is summarized as an illustration based on **Table 1**. S and R indicate smooth and rough in the colony textures. b) Profiles of neutral sugar content of the representing substrains measured in liquid and plate media. The top and bottom panels show the amounts of neutral sugar in liquid and plate culture, respectively. White and gray bars indicate the amounts of soluble and insoluble neutral sugar, respectively. Vertical error bars show 1 SD calculated from at least 4 independent experiments and * means significantly different in sugar production ($p < 0.02$).

Relationship between phenotypes and genotypes.

The remarkable phenotypic differences in the 32 clones are the colony textures on plate culture and pellicle (a liquid/air interface biofilm) formation in liquid culture (**Table 1**, **Fig. 1b** and **Fig. 3a**). Strains forming rough surface colonies produce more polysaccharide consisting of neutral sugars than the other (42,54). Polysaccharide productivities of the 32 clones were measured. Since the clones in each group showed a similar production of neutral sugar, the averages of the sugar production in each group were calculated and are shown in **Figure 3b**.

Groups represented by IFO 3283-32, -22, -26 and -03 showed significantly high production of neutral sugar on plate and in liquid culture. On the other hand, IFO 3283-07 and -12 groups produced less under both conditions. Interestingly, the IFO 3283-01 group produced soluble sugar in liquid culture as high as IFO3283-32 etc., but did less in the other conditions like the IFO 3283-07 and -12 groups (**Fig. 3b**). The constituents of the slime layer produced by smooth colony strains are possibly polysaccharides but chemically undetectable by the Phenol- H_2SO_4 method used in this study, and have never been determined.

Based on the genome lineage analysis, the single nucleotide nonsynonymous mutation in the glycerol kinase gene (SNP02) seems to relate to sugar production on plate and liquid media. However, at a terminal on the lineage, the high productivity of soluble polysaccharide of the IFO 3283-01 in liquid culture seems to be accompanied with disruption of *secB2* gene (Gap03). Colony textures and pellicle formation of all 32 isolates in media containing 0.5% glucose were same as without glucose (**Supplemental Fig. S2**), suggesting that the mutations, SNP02 and Gap03, affect polysaccharide production and the structural modification rather than glucose metabolism.

Glycerol kinase catalyzes the rate-limiting reaction in the pathway of glycerol utilization and the mutation is located in a beta sheet used to form the inter-homodimer Zn(II) binding site (55). Therefore, the SNP02 mutation in glycerol kinase gene might directly affect glucose control of glycerol utilization (56). But since the strain with SNP02 can grow on glycerol as a sole carbon source at the similar speed as the strain without the mutation, the glycerol utilization by glycerol kinase may be similar to each other strain. SecB is a molecular chaperone in the general secretion system (57). The *A. pasteurianus* genome as well as *G. oxydans* (11) and a zoonosis pathogen *Francisella tularensis* (58), contain a well conserved *secB1* (APA01-07990) and a divergent *secB2* (APA01-21762). The precise function of SecB2 and the relationship to polysaccharide synthesis are unknown. There are two genes related to glycerol metabolism (APA01-12570 and -12560) downstream of the glycerol kinase gene, and six genes (from APA01-21770 to -21820) in a predicted operon with *secB2* (**Supplemental Table S4**). It is possible that these gene products affect the

polysaccharide and pellicle production as well.

Fixation of mutant strains in population

A. pasteurianus IFO 3283 (NBRC 3283) was originally isolated from a pellicle. However, the strains of IFO 3283-07, -12 and -01 groups, which lost the ability to produce a pellicle, were fixed in this population and became a major portion of the cell complex (total 12 clones out of 32). It is a fundamental but difficult question to answer how mutant strains could be prominent during cell maintenance without special selective pressures.

The strains in the IFO 3283-07, -12 and -01 groups showed longer lag phase in the YPG medium containing 2.5% ethanol than the others, but in the absence of ethanol the former strains grew up to slightly but significantly higher density at stationary phase than the latter ones (data not shown). All strains showed no significant difference in viability in 5°C storage for 3 months (data not shown). The strains of IFO 3283-07, -12 and -01 groups form more smooth colonies to spread easily on the surface of solid medium. This could explain the fixation in the population.

No phenotypic differences were observed among the groups of IFO 3283-03, -22, -26 and -32, but the strains in the groups of IFO 3283-32 and IFO 3283-22 turned to minor groups (one each out of 32). Instead, strains in IFO 3283-03 and -26 turned to a major one (18/32) in the complex with two more mutations, SNP01 and SNP03. The SNP01 mutation in *rpoA* gene changes Leu to Gln at 149 aa, which locates at a hinge region between an N-terminal domain for initiation of subunit assembly and a C-terminal domain for transcriptional factor binding and DNA association (59). It is possible that the mutation affects the affinity of the alpha subunit to other transcriptional factors, and subsequently the transcriptional level of its regulon(s) modifies fitting ability to the given environment. A gene (APA01-17930) at SNP03 is hypothetical and probably monocistronic, thus the significance of SNP03 is unpredictable. The annotation of the genes related to SNP01 and other mutation sites are listed in **Supplemental Table S4**.

Transposons and plasmids

A. pasteurianus genome contains more than 280 transposons, approximately 9% of total genes in the genome. Phylogenetic analysis revealed that the transposases in those transposons construct several paralogous clusters and the transposase in IS1380, which was previously reported of *A. pasteurianus* (9), is most abundant and the total copy number is 74. The phylogenetic tree was shown in **Supplemental Figure 3**, and a list of transposases and the result of paralogous clustering were shown in **Supplemental Table S3**. Thirty-two genes assigned for specific functions are truncated by the transposons (**Supplemental Figure S3**) and the gene truncation may affect the metabolism or response to environment stimuli. For instance, a gene cluster similar to the nitrate assimilation gene cluster of *Granulibacter bethesdensis* CGDNIH1 (12) was identified in *A. pasteurianus* genome, but the assimilatory nitrate reductase catalytic subunit (APA01_42100) was truncated by a transposon while the genes encoding a assimilatory nitrate reductase electron-transfer subunit (APA01_42120) and a nitrate transporter (APA01_42150) contain a frameshift in each coding region. It suggests *A. pasteurianus* is not able to utilize nitrate as a source of nitrogen by the reduction of nitrate via nitrite to ammonium. Comparing simply with truncation of three genes in the 21-year maintenance, the truncation may occur within a few centuries, and no gene truncations in the previous era may be detected.

A. pasteurianus genome harbors 6 plasmids besides a number of transposons, which is 9% of total genes in the genome. Spiral bacteria *Borrelia* genomes harbor remarkable amount of plasmids (60), but transposons in *Borrelia* genomes account for only a few percentages of the total genes. Reversely, *Shigella* (61), *Xanthomonas* (62), and *Orientia* (63) contain a significant proportion of transposon but carry few plasmids in the genomes. The plot analysis of the proportion of transposase genes in the total genes and the fragment number including chromosomes and plasmids using genome sequences of 777 bacterial species in the public database, suggests that the combination of relatively large amounts of plasmids and transposons is one of the characteristics leading to the hyper-mutability of *A. pasteurianus* genome (**Fig. 4**).




Fig 4. Features of *A. pasteurianus* hyper-mutability.

Proportion of transposase genes in the total number of genes (vertical) and DNA fragment number of the genome including chromosomes and plasmids (horizontal) are plotted using each genome of 777 bacterial species with gene sequences in the public database (<ftp://ftp.ncbi.nlm.nih.gov/genbank/genomes/Bacteria>). Blue and purple spots indicate the proportion for *A. pasteurianus* IFO 3283-01 and other AAB, respectively. Remarkable bacterial in this plotting, such as *Borrelia* (60), *Shigella* (61), *Xanthomonas* (62), and *Orientia* (63), are indicated in orange, red, light green and green. Profiles for other bacteria are shown in black, and those close to *A. pasteurianus* were indicated by bars, such as *Aliivibrio*, *Acinetobacter* and *Lactococcus*.

Hyper-mutable tandem repeat (HTR)

Three HTRs that cause genome hyper-variation were detected through the genome DNA sequencing and its assembling. Tandem repeats (TRs) were extracted using the JSTRING software (40) to determine all of the HTRs from the genome DNA sequence of *A. pasteurianus*. Seven TRs (named as AP-TR01 to 07) in *A. pasteurianus* were found to have a score over 200 (64), including the 3 HTRs (AP-TR01, 04 and 07; Table 1). Six TRs, AP-TR01 and AP-TR03-07, are built from homogeneous repetitive sequences, and AP-TR02 is built from a heterogeneous one. All TRs are tracked into individual protein coding regions. Out of the 7 TRs, AP-TR01, 03, 04, 06 and 07 all showed hyper-mutability by PCR-directed DNA sequencing and therefore were designated as *A. pasteurianus* HTRs herein.

The repetitive sequence units and numbers of HTRs and genes including HTRs are listed in **Table 1**. In the case of AP-TR01 in the IFO 3283-01, a sequence unit and repetitive number are gcctgt and 14, respectively. While the repetitive number of each HTR was determined by the majority (approximately 90%) of DNA fragments, 5% of the one repeat shorter and 5% of longer were observed as contaminations (data not shown). It seems that HTRs could be expanded and contracted by repetitive units during DNA multiplication from a single cell to PCR directed colony sequencing, corresponding to several dozen generations (**Supplemental Fig. S2**).

The repetitive sequence units of AP-TR04 and AP-TR07 are penta (aggac) and octa (aggacaga) nucleotides, respectively. Therefore, the repetitive numbers alter the coding frames following the HTRs. When the repetitive number of AP-TR04 is multiple of three, the gene, APA01-18740, encodes a 1289 aa sequence consisting of three domains, a DNA helicase domain in the N-terminal, DRTGQ repeats at the TR and 283 aa in the C-terminal. In the other two cases of the repetitive numbers, the gene encodes 8 aa and 58 aa in the C-termini (**Fig. 5**). An interesting sequence was found in a draft DNA sequence of *G. xylinus* NBRC 3288 genome (unpublished work). The largest plasmid of *G. xylinus*, pGXY010, contains homogeneous TR (aggac)₂₈ in a DNA helicase gene similar to APA01-18740. The DNA helicase gene with the TR may be horizontally transferred between two independent genera by a plasmid and this horizontal gene/plasmid transfer might be another mechanism for genome instability of AAB.




Fig 5. A schematic illustration of the structures of gene products with hyper-mutable tandem repeat sequences.

Products of the genes with AP-TR04 are shown on top. Similar DNA helicases in the largest plasmid (BABN01000001) of *Gluconacetobacter xylinus* NBRC 3288, *Azotobacter vinelandii* DJ (65), *Pseudomonas syringae* pv. tomato str. DC3000 (NC-004578) and *Myxococcus xanthus* DK 1622 (NC-008095) are shown below. N-terminal regions in grey indicate a conserved DNA helicase domain. Tandem repeat regions are shown in green with strips for *A. pasteurianus* and *G. xylinus* and in white stripes for an *A. vinelandii* homologue. Amino acid sequences in C-terminal regions indicated in blue, yellow and red are encoded by same DNA sequences but different frame based on repetitive number of the AP-TR04 in *A. pasteurianus*. Amino acid sequences in C-terminal region in blue from 4 species are similar to each other. Gene locations, chromosome and plasmid, are indicated as *chr* and *pld*, respectively.

Genome DNA sequence of *Azotobacter vinelandii* DJ (65) was recently published and we found a high similarity between the two genes, *A. pasteurianus* APA01-18740 and *A. vinelandii* Avin_22890, which the authors assigned as a pseudogene (65). However *A. pasteurianus* and *A. vinelandii* belongs to different bacterial groups, alpha- and gamma-proteobacteria, respectively, the APA01-18740 and Avin_22890 contain TRs at a very similar site in each gene. Interestingly the repetitive sequence, (agcc)₄₀, in the Avin_22890 is different from one in the APA01-18740. However the primary habitats of present *A. vinelandii* and *A. pasteurianus* are different, both *A. vinelandii* and *A. pasteurianus* are an obligate aerobic bacteria and found in soil worldwide, and it is possible that *A. pasteurianus* and *A. vinelandii* have shared its habitation and exchanged their genome information in their evolutionary processes. It is still debatable whether the two genes are related to each other in their history or the existences of the two TRs in similar sites are just coincident. Whichever it is difficult to explain how the different repetitive sequence units were generated.

The most favorable explanation responsible for the mechanism of TR instability attributing to hereditary neurological disease such as polyglutamine expansion is a DNA strand slippage model (66,67), in which either the template or newly synthesized DNA strand forms partially stable secondary structures with repetitive sequences and overcoming the complex replication results in expansion or deletion of TRs. Several cellular mechanisms may underlay the process, such as general replication machinery including DNA polymerase III, helicases, recombination factors and DNA repair system (68-70). Where expansion and deletion of tandem repeats are thought to be enhanced by mutations that reduce the fidelity of replication (68), two genes with HTRs encoding a DNA helicase and DNA a polymerase III exonuclease epsilon subunit may be involved in the hyper-mutability of *A. pasteurianus*.

It is still unclear whether the involvement of translesion synthesis (TLS) or TLS polymerase is significant in the mechanism of expansions or deletions of TRs (69,71). Genome of *A. pasteurianus* contains genes encoding RecA, RecFOR, SSB, and DNA polymerase I, III and IV, but not RecBCD, or DNA polymerase II or V. Therefore it is possible that TLS is involved in *A. pasteurianus* hyper-mutability, if so, DNA polymerase IV, of which a mutant was shown to increase instability of TRs in *E. coli*, is the only error-prone DNA polymerase in *A. pasteurianus* (69).

Hyper-mutable TRs may play critical roles in the phenotypic modification of bacteria, such as phase variation and genetic diversity in pathogens. A TR is located in the promoter sequence of the *Neisseria meningitidis nadA* gene coding a pathogenic adhesion protein and the number of repeats influences the binding status of transcriptional regulators to the *nadA* promoter and the *nadA* gene expression (72). High-frequency phase variation of LPS structures, characteristic of *Haemophilus influenzae* requires *licA* gene regulation. Changes of the TR repeat number in the 5' region of *licA* result in a translational switch for the *licA* product involved in LPS biosynthesis (73,74). *A. pasteurianus* HTRs locate in the middle of protein coding region, thus the meaning of HTRs may be different from the cases for transcriptional and translational regulation.

Four other AAB genomes were analyzed using the JSTRING software, (40) to determine the prevalence of TRs in AAB. The *G. oxydans* 621H genome (11) also contains two homogeneous repeats. Genomes of other AAB, *G. bethesdensis* CGDNIH1 (12), *G. diazotrophicus* PAL5 and *A. cryptum* JF-5, include many heterogeneous repeats, such as a unique long repeating sequence (cccgtggcg)₂₀₅, but rarely a homogeneous one. As a result, TRs are prevalent in AAB, but sequences of the repeat units, repetitive numbers and genes with the TRs are varied. The TRs might be significant, but their roles are largely unknown.

Adaptation to unviable high-temperatures

To clarify whether *A. pasteurianus* has other systems for the genome instability and to show the utility of the genome instability to produce useful features, we performed an adaptation to unviable temperatures under a condition accelerating evolution in vitro. *A. pasteurianus* IFO 3283-01, which produces soluble polysaccharide in liquid culture and formed less aggregation, was used. *A. pasteurianus* can grow at 39°C, but the growth is unstable at 40°C. No growth was observed at 40.5°C or higher (data not shown). Then adaptation at high temperature was initiated by a long-term cultivation at 40°C to allow the cells grow stably at 40°C. Remarkable morphological changes were observed during this cultivation. The cells lengthened from approximate 2-3 µm up to 300 µm within the first 3 days, and 10 days later the cell length shortened to approximately 5 µm (**Supplemental Fig. S4**). The long cells contained multi-nucleoid bodies. It is not clear yet whether the long cells turned into shorter or that short cells present at beginning occupied a majority in the population.

After 27 days cultivation at 40°C, representing 5 times continuous dilution-growth culturing passages, the cells in 5th culture at 40°C (termed 40C(5)) could be successfully established to grow at 42°C (**Fig. 6a**). Growth at 42°C was tested for 365 days by serial dilution-growth cultivation (partially shown in **Fig. 6a**). In principle, the acquired resistance may result from physiological or genetic adaptation (75). When the acquired 42°C resistance resulted from a transient physiological adaptation similar to phenomenon of adaptive pH response in *Acetobacter* (8,76,77), the cells would rapidly lose the 42°C resistance upon cultivation under non-limiting conditions at 30°C culturing. The first evolved culture, 42C(1), and the parental strain IFO 3283-01 were grown on YPG plates at 30°C and 42°C for 3.5 days. The cells grown at 30°C were successively cultured on YPG plates at 30°C and 42°C twice a week. After 13 serial plating at 30°C, for total for 7 weeks representing 150 generations, the ability of growth at 42°C still remained (**Fig. 6b**), thus suggesting that the acquisition of growth ability at 42°C is based on mutation(s).

Genomic DNA prepared from a clone from 42C(30) and the parental strain IFO 3283-01 were compared by pulse field gel electrophoresis after digestion by restriction nucleases to clarify the genome-wide modifications. No remarkable differences were observed except a fragment probably corresponding to the smallest plasmid (data not shown). Plasmid contents were analyzed and it was revealed that proportion of the smallest plasmid in the genomic DNA was remarkably changed at the 11th culture 42C(11) and it was undetectable in the 42C(21) culture (**Supplemental Fig. S4**). Repetitive numbers of the 3 HTRs were modified in the cultivation at 42°C (**Supplemental Fig. S4**).

The clone, *A. pasteurianus* IFO 3283-01-42C, isolated from the first culture at 42°C may contain neither genome wide rearrangement nor plasmid depletion, and therefore mutation mapping was carried out by saturated genome shotgun sequencing with a Genome Analyzer (Illumina, San Diego, CA). Approximately 10 million reads of 35 bp raw data were obtained. Using the raw data, a high quality data set with only 1.4 million high quality reads was prepared, corresponding to 14 folds of *A. pasteurianus* genome (**Supplemental Fig. S4**). Mapping analysis was performed with the whole and high quality data sets using the MAQ software (31).

Analyses with the whole and high quality data sets indicated that there were 13 and 4 SNP candidates in chromosome, respectively, but no SNP candidates in the plasmids. Three SNP candidates were chosen from both analyses from the two data sets, thus DNA sequences at 14 SNP candidate loci were determined using PCR-direct DNA sequencing. Consequently the 3 SNP candidates, i.e. SNP42-01, SNP42-03, and SNP42-04, chosen with both data sets were shown as actual SNPs in IFO 3283-01-42C genome. Besides SNPs, sequence mapping indicated that there are clustered regions without mapped sequences at the predicted replication terminal region (**Fig. 6c**). The entire regions were thoroughly sequenced using PCR products, and it revealed that a region 92 kb in length involving APA01-13760 to APA01-14470 was truncated.




Fig 6. Adaptation to an unviable environment at high-temperatures.

a) Growth profiles of *A. pasteurianus* IFO 3283-01 in the successful case for adaptation. Green, blue and red curves indicate cell growth at 30°C, 40°C and 42°C, respectively. Three arrows, 40C(3), 40C(5) and 42C(1), indicate the cell cultures analyzed for mutations. b) A heritability analysis of growth at 42°C. The seed strain (IFO 3283-01), the bred substrain (IFO 3283-01/42°C) and a different cell type strain (IFO3283-02) were inoculated on same plate and tested for growth at 30°C and 42°C. Cells grown on a 30°C plate were then inoculated onto the next two plates for growth at 30°C and 42°C. c) A mapping analysis of sequencing reads on the chromosome of IFO 3283-01/42°C (top panel). Vertical and horizontal axes indicate chromosome location and frequency of high quality sequence reads in 1000 bases. PCR analyses were carried out for the large deletion region and positions of primer pairs for PCR analyses were shown in the middle. DNA was used from the clone IFO 3283-01-42C isolated from the culture 42C(1) indicated by the right arrow in the panel a. Grey zones and grey arrows in the middle panel indicate existence of the region in the genomic DNA, while white areas, no-mapped regions, and white arrow show the region deleted from the genome. PCR products analyzed in agarose gel electrophoresis were shown in the bottom. O and A on the top of each lane designate DNA templates for PCR prepared from the original (IFO 3283-01) and the adapted strain (IFO 3283-01-42C), respectively. d) A summary of the mutations found in the adapted strain (IFO 3283-01-42C). Original and mutated sequences are shown in capital and small letters, respectively.

To clarify the contributions of the SNPs and the 92 kb deletion region for adaptive evolution at high temperature, clones from 40C(3) and 40C(5) were tested for viability at 42°C and the genome variations shown in IFO 3283-01-42C. The 40C(5) clone was able to multiply at 42°C but not the 40C(3) one. No mutations were observed in the 40C(3) clone, but the SNP42-01 and 92 kb deletion were detected in 40C(5). This indicated that the SNP42-01 and/or 92 kb deletion might be involved in the adaptive evolution (Fig. 6d).

The SNP42-01 was located in APA01-42C-00810 coding a two-component hybrid sensor histidine kinase and regulator. The mutation, from C1187 to T, is non-synonymous, and resulted in amino acid substitution T396I in a conserved domain among histidine kinases. Therefore, the mutation may affect the signal transduction by an unknown environmental stimulation. The 92 kb deleted region includes 72 genes including a rRNA operon, 16 putative pseudogenes, 21 metabolism-related genes, 14 transposons, 13 transporter genes and 6 genes encoding unknown function proteins (Supplemental Table S2). It is very difficult to evaluate the association of the SNP42-01 gene and genes in the deleted region with the adaptive evolution. While shorter plasmids and mitochondrial genomes show faster replication (78), the 92 kb deletion corresponding to approximate 3% of the whole genome may affect growth speed or cell viability by decreasing the stress for proliferation in this stressful environment.

Concluding remarks.

Microorganisms utilized in academic and industrial fields must be maintained to conserve identical characteristics wherever and whenever used. Extraordinary consideration for storage and handling should be paid to hyper-mutable microorganisms isolated from nature. In the middle of the 20th century, it was believed that AAB should be maintained by serial passage to avoid phenotypic modification, but neither frozen nor maintained at 4°C, because of the extraordinarily low survival rate and high mutation frequency (79,80). Furthermore, colony isolation during the passage was prohibited to avoid cloning unanticipated mutants. Attempting to clarify the hyper-mutability of *A. pasteurianus*, known as a hyper-mutable bacterium (6,9,10), therefore required the genome DNA sequencing of cells

grown in two different culturing conditions.

First, *A. pasteurianus* IFO 3283 (NBRC 3283), which formed a multi-phenotype cell complex following a series of slant passages for 21 years, was subjected to “mixed” genome DNA sequencing. The genome analysis revealed that more than 280 transposons, approximately 9% of the genes, exist in the genome consisting of a 2.9 Mb chromosome and 6 plasmids. The *A. pasteurianus* genome may be characterized by a combination of a large number both transposons and plasmids. Actually 5 variations by transposon insertion, as well as 3 SNPs, were found in the IFO 3283 (NBRC 3283) complex. A plasmid of *G. xylinus* carries a gene specifically similar to one in the *A. pasteurianus* chromosome, and a plasmid of *A. pasteurianus* was depleted during high temperature adaptation. These findings may illustrate the involvement of plasmids in the genetic instability of AAB. The existence of HTRs in open reading frames is a characteristic of the *A. pasteurianus* genome. However, it seems to be involved in the feasibility of DNA maintenance, the physiological meaning is largely unknown.

Second, *A. pasteurianus* was used for the breeding of a strain proliferative at an unviable high-temperature (42°C). Genome analyses of the bred strain by saturated genome shotgun sequencing revealed that a 92 kb deletion (approximately 3% of the whole genome size) and three single nucleotide mutations occurred in the genome during the high-temperature adaptation. However, the generating mechanism and genetic meaning of the large deletion is unknown, it seems that the smaller genome yields a survival advantage in the highly stressful conditions, possibly because of faster replication or lower heat generation based on less burden in DNA replication.

Alpha-proteobacteria, including AAB, contains many intracellular symbionts and parasites such as bacteria in the families, Rhizobiaceae, Rickettsiaceae, and Brucellaceae, whose genomes are generally shorter and show increased evolution rates in comparison to those of closely related free-living bacteria (20). However, *A. pasteurianus* is assumed to be a free-living bacterium, it may have evolved to fit in competitive natural niches of seasonal fruits and flowers accompanied by the genetic flexibility and instability.

Acknowledgements

This study was supported by a Grant-in-Aid from the Program for Promotion of Basic Research Activities for Innovative Biosciences (PROBRAIN), Mizkan Group Corporation and Innovation Center of Yamaguchi University.

References

1. Lambert, B., Kersters, K., Gossele, F., Swings, J. and De Ley, J. (1981) Gluconobacters from honey bees. *Antonie Van Leeuwenhoek*, **47**, 147-157.
2. Ryu, J.H., Kim, S.H., Lee, H.Y., Bai, J.Y., Nam, Y.D., Bae, J.W., Lee, D.G., Shin, S.C., Ha, E.M. and Lee, W.J. (2008) Innate immune homeostasis by the homeobox gene caudal and commensal-gut mutualism in *Drosophila*. *Science*, **319**, 777-782.
3. Yamada, Y. and Yukphan, P. (2008) Genera and species in acetic acid bacteria. *Int J Food Microbiol*, **125**, 15-24.
4. Greenberg, D.E., Porcella, S.F., Stock, F., Wong, A., Conville, P.S., Murray, P.R., Holland, S.M. and Zelazny, A.M. (2006) *Granulibacter bethesdensis* gen. nov., sp. nov., a distinctive pathogenic acetic acid bacterium in the family Acetobacteraceae. *Int J Syst Evol Microbiol*, **56**, 2609-2616.
5. Adachi, O., Moonmangmee, D., Toyama, H., Yamada, M., Shinagawa, E. and Matsushita, K. (2003) New developments in oxidative fermentation. *Appl Microbiol Biotechnol*, **60**, 643-653.
6. Beppu, T. (1993) Genetic organization of *Acetobacter* for acetic acid fermentation. *Antonie Van Leeuwenhoek*, **64**, 121-135.
7. Sokollek, S.J., Hertel, C. and Hammes, W.P. (1998) Description of *Acetobacter oboediens* sp. nov. and *Acetobacter pomorum* sp. nov., two new species isolated from industrial vinegar fermentations. *Int J Syst Bacteriol*, **48 Pt 3**, 935-940.
8. Steiner, P. and Sauer, U. (2001) Proteins induced during adaptation of *Acetobacter aceti* to high acetate concentrations. *Appl Environ Microbiol*, **67**, 5474-5481.
9. Takemura, H., Horinouchi, S. and Beppu, T. (1991) Novel insertion sequence IS1380 from *Acetobacter*

- pasteurianus* is involved in loss of ethanol-oxidizing ability. *J Bacteriol*, **173**, 7070-7076.
10. Coucheron, D.H. (1991) An *Acetobacter xylinum* insertion sequence element associated with inactivation of cellulose production. *J Bacteriol*, **173**, 5723-5731.
 11. Prust, C., Hoffmeister, M., Liesegang, H., Wiezer, A., Fricke, W.F., Ehrenreich, A., Gottschalk, G. and Deppenmeier, U. (2005) Complete genome sequence of the acetic acid bacterium *Gluconobacter oxydans*. *Nat Biotechnol*, **23**, 195-200.
 12. Greenberg, D.E., Porcella, S.F., Zelazny, A.M., Virtaneva, K., Sturdevant, D.E., Kupko, J.J., 3rd, Barbian, K.D., Babar, A., Dorward, D.W. and Holland, S.M. (2007) Genome sequence analysis of the emerging human pathogenic acetic acid bacterium *Granulibacter bethesdensis*. *J Bacteriol*, **189**, 8727-8736.
 13. Sallstrom, B. and Andersson, S.G. (2005) Genome reduction in the alpha-Proteobacteria. *Curr Opin Microbiol*, **8**, 579-585.
 14. Nobusato, A., Uchiyama, I., Ohashi, S. and Kobayashi, I. (2000) Insertion with long target duplication: a mechanism for gene mobility suggested from comparison of two related bacterial genomes. *Gene*, **259**, 99-108.
 15. Finkel, S.E. and Kolter, R. (1999) Evolution of microbial diversity during prolonged starvation. *Proc Natl Acad Sci U S A*, **96**, 4023-4027.
 16. Bjedov, I., Tenaillon, O., Gerard, B., Souza, V., Denamur, E., Radman, M., Taddei, F. and Matic, I. (2003) Stress-induced mutagenesis in bacteria. *Science*, **300**, 1404-1409.
 17. Drake, J.W., Charlesworth, B., Charlesworth, D. and Crow, J.F. (1998) Rates of spontaneous mutation. *Genetics*, **148**, 1667-1686.
 18. Nilsson, A.I., Koskiniemi, S., Eriksson, S., Kugelberg, E., Hinton, J.C. and Andersson, D.I. (2005) Bacterial genome size reduction by experimental evolution. *Proc Natl Acad Sci U S A*, **102**, 12112-12116.
 19. Kim, W.S., Park, J.H., Ren, J., Su, P. and Dunn, N.W. (2001) Survival response and rearrangement of plasmid DNA of *Lactococcus lactis* during long-term starvation. *Appl Environ Microbiol*, **67**, 4594-4602.
 20. Andersson, S.G. and Kurland, C.G. (1998) Reductive evolution of resident genomes. *Trends Microbiol*, **6**, 263-268.
 21. Stephens, R.S., Kalman, S., Lammel, C., Fan, J., Marathe, R., Aravind, L., Mitchell, W., Olinger, L., Tatusov, R.L., Zhao, Q. *et al.* (1998) Genome sequence of an obligate intracellular pathogen of humans: *Chlamydia trachomatis*. *Science*, **282**, 754-759.
 22. Moran, N.A. (2002) Microbial minimalism: genome reduction in bacterial pathogens. *Cell*, **108**, 583-586.
 23. Azuma, Y., Hirakawa, H., Yamashita, A., Cai, Y., Rahman, M.A., Suzuki, H., Mitaku, S., Toh, H., Goto, S., Murakami, T. *et al.* (2006) Genome sequence of the cat pathogen, *Chlamydophila felis*. *DNA Res*, **13**, 15-23.
 24. Dufresne, A., Garczarek, L. and Partensky, F. (2005) Accelerated evolution associated with genome reduction in a free-living prokaryote. *Genome Biol*, **6**, R14.
 25. Moran, N.A. and Mira, A. (2001) The process of genome shrinkage in the obligate symbiont *Buchnera aphidicola*. *Genome Biol*, **2**, RESEARCH0054.
 26. Silva, F.J., Latorre, A. and Moya, A. (2001) Genome size reduction through multiple events of gene disintegration in *Buchnera* APS. *Trends Genet*, **17**, 615-618.
 27. Sekine, M., Tanikawa, S., Omata, S., Saito, M., Fujisawa, T., Tsukatani, N., Tajima, T., Sekigawa, T., Kosugi, H., Matsuo, Y. *et al.* (2006) Sequence analysis of three plasmids harboured in *Rhodococcus erythropolis* strain PR4. *Environ Microbiol*, **8**, 334-346.
 28. Gordon, D., Desmarais, C. and Green, P. (2001) Automated finishing with autofinish. *Genome Res*, **11**, 614-625.
 29. Ewing, B. and Green, P. (1998) Base-calling of automated sequencer traces using phred. II. Error probabilities. *Genome Res*, **8**, 186-194.
 30. Ewing, B., Hillier, L., Wendl, M.C. and Green, P. (1998) Base-calling of automated sequencer traces using phred. I. Accuracy assessment. *Genome Res*, **8**, 175-185.
 31. Li, H., Ruan, J. and Durbin, R. (2008) Mapping short DNA sequencing reads and calling variants using mapping quality scores. *Genome Res*.
 32. Altschul, S.F., Madden, T.L., Schaffer, A.A., Zhang, J., Zhang, Z., Miller, W. and Lipman, D.J. (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res*, **25**, 3389-3402.
 33. Lowe, T.M. and Eddy, S.R. (1997) tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res*, **25**, 955-964.

34. Sakiyama, T., Takami, H., Ogasawara, N., Kuhara, S., Kozuki, T., Doga, K., Ohyama, A. and Horikoshi, K. (2000) An automated system for genome analysis to support microbial whole-genome shotgun sequencing. *Biosci Biotechnol Biochem*, **64**, 670-673.
35. Yada, T., Nakao, M., Totoki, Y. and Nakai, K. (1999) Modeling and predicting transcriptional units of *Escherichia coli* genes using hidden Markov models. *Bioinformatics*, **15**, 987-993.
36. Delcher, A.L., Harmon, D., Kasif, S., White, O. and Salzberg, S.L. (1999) Improved microbial gene identification with GLIMMER. *Nucleic Acids Res*, **27**, 4636-4641.
37. Altschul, S.F., Gish, W., Miller, W., Myers, E.W. and Lipman, D.J. (1990) Basic local alignment search tool. *J Mol Biol*, **215**, 403-410.
38. Pearson, W.R. and Lipman, D.J. (1988) Improved tools for biological sequence comparison. *Proc Natl Acad Sci U S A*, **85**, 2444-2448.
39. Hirokawa, T., Boon-Chieng, S. and Mitaku, S. (1998) SOSUI: classification and secondary structure prediction system for membrane proteins. *Bioinformatics*, **14**, 378-379.
40. De Fonzo, V., Aluffi-Pentini, F. and Parisi, V. (2008) JSTRING: A Novel Java Tandem Repeats Searcher in Genomic Sequences with an Interactive Graphic Output. *Open Applied Informatics Journal*, **2**, 14-17.
41. Dubois, M., Gilles, K., Hamilton, J.K., Rebers, P.A. and Smith, F. (1951) A colorimetric method for the determination of sugars. *Nature*, **168**, 167.
42. Deeraksa, A., Moonmangmee, S., Toyama, H., Yamada, M., Adachi, O. and Matsushita, K. (2005) Characterization and spontaneous mutation of a novel gene, *polE*, involved in pellicle formation in *Acetobacter tropicalis* SKU1100. *Microbiology*, **151**, 4111-4120.
43. Kibota, T.T. and Lynch, M. (1996) Estimate of the genomic mutation rate deleterious to overall fitness in *E. coli*. *Nature*, **381**, 694-696.
44. Boe, L., Danielsen, M., Knudsen, S., Petersen, J.B., Maymann, J. and Jensen, P.R. (2000) The frequency of mutators in populations of *Escherichia coli*. *Mutat Res*, **448**, 47-55.
45. Drake, J.W. (1991) A constant rate of spontaneous mutation in DNA-based microbes. *Proc Natl Acad Sci U S A*, **88**, 7160-7164.
46. Loewe, L., Textor, V. and Scherer, S. (2003) High deleterious genomic mutation rate in stationary phase of *Escherichia coli*. *Science*, **302**, 1558-1560.
47. Grigoriev, A. (1998) Analyzing genomes with cumulative skew diagrams. *Nucleic Acids Res*, **26**, 2286-2290.
48. Nomura, M. and Morgan, E.A. (1977) Genetics of bacterial ribosomes. *Annu Rev Genet*, **11**, 297-347.
49. Kylma, A.K., Granstrom, T. and Leisola, M. (2004) Growth characteristics and oxidative capacity of *Acetobacter aceti* IFO 3281: implications for L-ribulose production. *Appl Microbiol Biotechnol*, **63**, 584-591.
50. Nabe, K., Izuo, N., Yamada, S. and Chibata, I. (1979) Conversion of Glycerol to Dihydroxyacetone by Immobilized Whole Cells of *Acetobacter xylinum*. *Appl Environ Microbiol*, **38**, 1056-1060.
51. Weinhouse, H. and Benziman, M. (1976) Phosphorylation of glycerol and dihydroxyacetone in *Acetobacter xylinum* and its possible regulatory role. *J Bacteriol*, **127**, 747-754.
52. Adachi, O., Ano, Y., Shinagawa, E. and Matsushita, K. (2008) Purification and properties of two different dihydroxyacetone reductases in *Gluconobacter suboxydans* grown on glycerol. *Biosci Biotechnol Biochem*, **72**, 2124-2132.
53. Mullins, E.A., Francois, J.A. and Kappock, T.J. (2008) A specialized citric acid cycle requiring succinyl-coenzyme A (CoA):acetate CoA-transferase (AarC) confers acetic acid resistance on the acidophile *Acetobacter aceti*. *J Bacteriol*, **190**, 4933-4940.
54. Moonmangmee, S., Kawabata, K., Tanaka, S., Toyama, H., Adachi, O. and Matsushita, K. (2002) A novel polysaccharide involved in the pellicle formation of *Acetobacter aceti*. *J Biosci Bioeng*, **93**, 192-200.
55. Pettigrew, D.W., Liu, W.Z., Holmes, C., Meadow, N.D. and Roseman, S. (1996) A single amino acid change in *Escherichia coli* glycerol kinase abolishes glucose control of glycerol utilization in vivo. *J Bacteriol*, **178**, 2846-2852.
56. Chagneau, C., Heyde, M., Alonso, S., Portulier, R. and Laloi, P. (2001) External-pH-dependent expression of the maltose regulon and *ompF* gene in *Escherichia coli* is affected by the level of glycerol kinase, encoded by *glpK*. *J Bacteriol*, **183**, 5675-5683.
57. van der Sluis, E.O. and Driessen, A.J. (2006) Stepwise evolution of the Sec machinery in Proteobacteria. *Trends*

- Microbiol*, **14**, 105-108.
58. Larsson, P., Oyston, P.C., Chain, P., Chu, M.C., Duffield, M., Fuxelius, H.H., Garcia, E., Halltorp, G., Johansson, D., Isherwood, K.E. *et al.* (2005) The complete genome sequence of *Francisella tularensis*, the causative agent of tularemia. *Nat Genet*, **37**, 153-159.
 59. Ishihama, A. (1992) Role of the RNA polymerase alpha subunit in transcription activation. *Mol Microbiol*, **6**, 3283-3288.
 60. Fraser, C.M., Casjens, S., Huang, W.M., Sutton, G.G., Clayton, R., Lathigra, R., White, O., Ketchum, K.A., Dodson, R., Hickey, E.K. *et al.* (1997) Genomic sequence of a Lyme disease spirochaete, *Borrelia burgdorferi*. *Nature*, **390**, 580-586.
 61. Jin, Q., Yuan, Z., Xu, J., Wang, Y., Shen, Y., Lu, W., Wang, J., Liu, H., Yang, J., Yang, F. *et al.* (2002) Genome sequence of *Shigella flexneri* 2a: insights into pathogenicity through comparison with genomes of *Escherichia coli* K12 and O157. *Nucleic Acids Res*, **30**, 4432-4441.
 62. da Silva, A.C., Ferro, J.A., Reinach, F.C., Farah, C.S., Furlan, L.R., Quaggio, R.B., Monteiro-Vitorello, C.B., Van Sluys, M.A., Almeida, N.F., Alves, L.M. *et al.* (2002) Comparison of the genomes of two *Xanthomonas* pathogens with differing host specificities. *Nature*, **417**, 459-463.
 63. Cho, N.H., Kim, H.R., Lee, J.H., Kim, S.Y., Kim, J., Cha, S., Kim, S.Y., Darby, A.C., Fuxelius, H.H., Yin, J. *et al.* (2007) The *Orientia tsutsugamushi* genome reveals massive proliferation of conjugative type IV secretion system and host-cell interaction genes. *Proc Natl Acad Sci U S A*, **104**, 7981-7986.
 64. Parisi, V., De Fonzo, V. and Aluffi-Pentini, F. (2003) STRING: finding tandem repeats in DNA sequences. *Bioinfo*, **19**, 1733-1738.
 65. Setubal, J.C., Dos Santos, P., Goldman, B.S., Ertesvag, H., Espin, G., Rubio, L.M., Valla, S., Almeida, N.F., Balasubramanian, D., Cromes, L. *et al.* (2009) The genome sequence of *Azotobacter vinelandii*, an obligate aerobic specialized to support diverse anaerobic metabolic processes. *J Bacteriol*. **191**, 4534-4545
 66. Wells, R.D., Parniewski, P., Pluciennik, A., Bacolla, A., Gellibolian, R. and Jaworski, A. (1998) Small slipped register genetic instabilities in *Escherichia coli* in triplet repeat sequences associated with hereditary neurological diseases. *J Biol Chem*, **273**, 19532-19541.
 67. Lechner, R.L., Engler, M.J. and Richardson, C.C. (1983) Characterization of strand displacement synthesis catalyzed by bacteriophage T7 DNA polymerase. *J Biol Chem*, **258**, 11174-11184.
 68. Iyer, R.R., Pluciennik, A., Rosche, W.A., Sinden, R.R. and Wells, R.D. (2000) DNA polymerase III proofreading mutants enhance the expansion and deletion of triplet repeat sequences in *Escherichia coli*. *J Biol Chem*, **275**, 2174-2184.
 69. Jacob, K.D. and Eckert, K.A. (2007) *Escherichia coli* DNA polymerase IV contributes to spontaneous mutagenesis at coding sequences but not microsatellite alleles. *Mutat Res*, **619**, 93-103.
 70. Delagoutte, E., Goellner, G.M., Guo, J., Baldacci, G. and McMurray, C.T. (2008) Single-stranded DNA-binding protein in vitro eliminates the orientation-dependent impediment to polymerase passage on CAG/CTG repeats. *J Biol Chem*, **283**, 13341-13356.
 71. Bichara, M., Wagner, J. and Lambert, I.B. (2006) Mechanisms of tandem repeat instability in bacteria. *Mutat Res*, **598**, 144-163.
 72. Martin, P., Makepeace, K., Hill, S.A., Hood, D.W. and Moxon, E.R. (2005) Microsatellite instability regulates transcription factor binding and gene expression. *Proc Natl Acad Sci U S A*, **102**, 3800-3804.
 73. Weiser, J.N., Shchepetov, M. and Chong, S.T. (1997) Decoration of lipopolysaccharide with phosphorylcholine: a phase-variable characteristic of *Haemophilus influenzae*. *Infect Immun*, **65**, 943-950.
 74. Weiser, J.N., Love, J.M. and Moxon, E.R. (1989) The molecular mechanism of phase variation of *H. influenzae* lipopolysaccharide. *Cell*, **59**, 657-665.
 75. Sauer, U. (2001) Evolutionary engineering of industrially important microbial phenotypes. *Adv Biochem Eng Biotechnol*, **73**, 129-169.
 76. Bearson, S., Bearson, B. and Foster, J.W. (1997) Acid stress responses in enterobacteria. *FEMS Microbiol Lett*, **147**, 173-180.
 77. Steiner, P. and Sauer, U. (2003) Long-term continuous evolution of acetate resistant *Acetobacter aceti*. *Biotechnol Bioeng*, **84**, 40-44.
 78. Casane, D., Dennebouy, N., de Rochambeau, H., Mounolou, J.C. and Monnerot, M. (1994) Genetic analysis of

- systematic mitochondrial heteroplasmy in rabbits. *Genetics*, **138**, 471-480.
79. Lopez, A., Johnson, L.W. and Wood, C.B. (1961) Observations on a laboratory method for submerged acetic fermentation. *Appl Microbiol*, **9**, 425-433.
80. Sourek, J. and Kulhanek, M. (1969) Long-term storage of acetic acid bacteria by means of lyophilization. *Zentralbl Bakteriol Parasitenkd Infektionskr Hyg*, **123**, 580-585.

1 **Table 1. Summary of phenotypes and mutation loci in the IFO 3283 (NBRC 3283) multi-phenotype complex.**

| | | Strain represent | NBRC3283-01 | NBRC3283-07 | NBRC3283-12 | NBRC3283-32 | NBRC3283-22 | NBRC3283-26 | NBRC3283-03 | | | | | |
|------------|-----------------|--|---------------------|---------------------|----------------------------|---------------------|---------------------|---------------------|--------------------------------|-------------|-------------|-------------|-------------|----|
| | | NBRC No. | NBRC 105184 | NBRC 105187 | NBRC 105191 | NBRC 105190 | NBRC 105188 | NBRC 105189 | NBRC 105186 | | | | | |
| | | DDJ accession No. | AP011121 – AP011127 | AP011135 – AP011141 | AP011170 – AP011176 | AP011156 – AP011162 | AP011142 – AP011148 | AP011149 – AP011155 | AP011128 – AP011134 | | | | | |
| | | Clones | 1,5,9,10,13,16 | 6,7,11,14,15 | 12 | 32 | 22 | 24,26,27 | 2-4,8,17-21,23, 25,28,29,30,31 | | | | | |
| Phenotypes | | Colony texture | | | Surface | Smooth & mucoid | Smooth & mucoid | Smooth & mucoid | Rough | Rough | Rough | Rough | | |
| | | Neutral sugar production and biofilm formation in NCG liquid culture | | | Transparency | Transparent | Transparent | Transparent | Dark | Dark | Dark | Dark | | |
| | | | | | Pellicle | - | - | - | + | + | + | + | | |
| | | | | | Total culture | High | Low | Low | High | High | High | High | | |
| | | | | | Retained on cells | Low | Low | Low | High | High | High | High | | |
| Genotypes | Chromosome | Single nucleotide mutation (SNP) | Variation | Gene | Annotation | Mutation | NBRC3283-01 | NBRC3283-07 | NBRC3283-12 | NBRC3283-32 | NBRC3283-22 | NBRC3283-26 | NBRC3283-03 | |
| | | | SNP01 | APA01_09550 | RNA pol alpha, RpsA | T > u | T | T | T | T | T | T | a | a |
| | | | SNP02 | APA01_12552 | Glycerol kinase | G > t | t | t | t | G | G | G | G | G |
| | | SNP03 | APA01_17930 | Hypothetical | No of A residues | 9 | 9 | 9 | 9 | 9 | 9 | 9 | 9 | 8 |
| | | Transposon insertion (Tp) | Gap03 | APA01_21762 | Protein translocase, SecB2 | insertion | + | - | - | - | - | - | - | - |
| | | | Gap14 | APA01_10210 | histidine kinase | insertion | - | - | - | - | + | + | + | + |
| | | | Gap18 | intergenic | (APA01_18050 /18060) | insertion | - | - | - | - | + | + | + | + |
| | | | Gap22 | APA01_24442 | Glycosyl transferase | insertion | + | + | - | - | - | - | - | - |
| | | Hyper-mutable tandem repeat (HTR) | AP-TR01 | APA01_02732 | Hypothetical | ggcgt | 14 | 13 | 13 | 13 | 13 | 13 | 13 | 13 |
| | | | AP-TR03 | APA01_16988 | Transcriptional regulator | tgcc | 24 | 22 | 22 | 20 | 19 | 21 | 20 | |
| | | | AP-TR04 | APA01_18740 | DNA helicase | ggaca | 36 | 36 | 35 | 41 | 30 | 36 | 33 | |
| | | | AP-TR06 | APA01_20312 | DNA pol III epsilon | aaggcgar | 10 | 8 | 8 | 8 | 8 | 8 | 8 | 8 |
| | Largest plasmid | Tp | Gap107 | APA01_40012 | Transposase | insertion | + | - | - | - | - | - | - | |
| | | HTR | AP-TR07 | APA01_40320 | Hypothetical | aggacaga | 14 | 14 | 12 | 16 | 14 | 13 | 14 | |

2 *1. High and low sugar production were determined based on the data of Fig. 3b.

3 *2. Repetitive numbers of the representative substrains are indicated.

4

ゲノム情報解析に基づく酢酸菌の遺伝的不安定性要因の解析

東 慶直

要旨

酢酸菌は α -proteobacteriaに属する細菌群で、食酢醸造槽から主要発酵菌として分離される。自然界では糖が蓄積するニッチ(花や果実など)に酵母などと共棲し広く分布する。一般にアルコールや酢酸に対する高い耐性、多様な有用二次代謝産物の生産、バイオポリマー生成など非常に多彩な特徴を持つ。一方で、酢酸菌の形質は継代培養によって変化しやすく、発酵産業や研究施設においては有用形質を維持するために経験的な対策が取られてきた。我々は酢酸菌が有する多様な有用代謝経路を明らかにするとともに、その多様性を生み出す遺伝的背景を明らかにすることを目的として、本邦伝統的酢酸発酵菌*Acetobacter pasteurianus*の全ゲノムDNA配列を詳細に解読した。ゲノム解析の試料には20年にも及ぶ継代培養により複数の形質を示す変異体複合菌系を用いた。そのゲノム解析からその複合菌系に共通する特徴として、2.9 Mbpの環状ゲノムDNAと6つの環状プラスミドを有し、合計3107遺伝子をコードすることが明らかとなった。形質の変化しやすさ、つまりゲノムの易変異性はそのプラスミドの多さに加え、トランスポゾンが全遺伝子の9%にも及ぶ280存在することが原因すると考えられる。さらに、細菌ゲノムとしては稀な遺伝子内部に存在するマイクロサテライトが5ヶ所発見された。一方、複合菌系に存在する差異を明らかにするために複合菌系から32株を分離し詳細に解析したところ、少なくとも3ヶ所の一塩基多型とトランスポゾンの転移による遺伝子の破壊が5ヶ所観察された。これは20年間の継代培養から考慮するとき、酢酸菌での変異発生率は大腸菌で報告されている変異発生率よりも有意に高いことが示された。酢酸菌のマイクロサテライトの伸長・短縮は20世代くらいで1反復配列程度を示した。酢酸菌が有する易変異性を利用し通常では生育不可能な高温(42°C)で増殖が可能となる菌株の育種を試みた。40°Cでの長期の培養の末、42°Cで増殖する株の獲得に成功した。その育種株の高温耐性形質が遺伝することを確認の後、全ゲノムDNA配列を次世代シーケンサーによって決定した。3カ所の一塩基置換と複製の集結店付近における92 kbpにおよぶゲノムの欠落が観察された。育種途中の株の解析から高温耐性にはその92 kbpの欠落のみが関与することが明らかとなった。その欠落領域には72遺伝子が存在し、そのいずれかの遺伝子の欠落によって生育限界を超えた高い温度で増殖が可能となったと考えられる。一方、欠落自体によりゲノムDNA複製のストレスが軽減され高い温度でも増殖が可能となった可能性もある。今後の解析を待つ必要がある。酢酸菌が属する α -proteobacteriaにはリケッチア菌群や根粒細菌群が含まれる。これらリケッチア菌群や根粒細菌群は動物や植物の細胞中をその増殖のニッチとして選択した菌群であり、顕著なゲノム縮重を進化の特徴としている。酢酸菌は環境中で自由に増殖が可能であるが、リケッチア菌群や根粒細菌群と同様に高い易変異性や大胆なゲノムDNAの欠落を許容しうる機構を有するのかもしれない。