

1 **Appeared as:** Smith, J.T., Sasina, N.V., Kryshev, A.I., Belova, N.V., Kudelsky, A.V. (2009,). A
2 review and blind test of predictive models for the bioaccumulation of radiostrontium in fish. *Journal*
3 *of Environmental Radioactivity*, 100, 950-954.
4

5 A review and test of predictive models for the bioaccumulation of
6 radiostrontium in fish.

7
8 J.T. Smith^{a*}, N.V. Sasina^b, A.I. Kryshev^c, N.V. Belova^d, A.V. Kudelsky^b
9

10 a. School of Earth and Environmental Sciences, University of Portsmouth, Burnaby
11 Building, Burnaby Road, Portsmouth PO1 3QL. E-mail: jim.smith@port.ac.uk
12 Tel : 02392-842416; Fax: 023 9284 2244.

13 b. Belarussian Geological Prospecting Research Institute, Ministry of Natural Resources
14 and Protection of Environment. Academician Kuprevich str. 7. 220141 Minsk, Belarus

15 c. Scientific and Production Association “Typhoon”, 82 Lenin Ave., Obninsk, Kaluga
16 Region, 249038, Russia.

17 d. Biology Faculty, Moscow State University, Moscow, Russia.

18 * Corresponding author.
19

20 **Keywords:** Radiostrontium, ⁹⁰Sr; ⁸⁹Sr; fish, bioaccumulation, concentration factor, model,
21 size effect.
22

23 **Abstract**
24

25 Empirical relations between the ⁹⁰Sr concentration factor (*CF*) and the calcium concentration
26 in freshwater aquatic systems have previously been determined in studies based on data
27 obtained prior to the Chernobyl accident. The purpose of the present research is to review and
28 compare these models, and to test them against a database of post-Chernobyl measurements
29 from rivers and lakes in Ukraine, Russia, Belarus and Finland. It was found that two
30 independently developed models, based on pre-Chernobyl empirical data, are in close
31 agreement with each other, and with empirical data. Testing of both models against new data
32 obtained after the Chernobyl accident confirms the models' predictive ability. An
33 investigation of the influence of fish size on ⁹⁰Sr accumulation showed no significant
34 relationship, though the data set was somewhat limited.

1. Introduction

For many radionuclides, only single “best estimate” fish-water concentration factors (*CF*s) are available for dose assessment models (e.g. IAEA, 1994). For some radionuclides, however, estimates of the *CF* may be improved using empirical models which account for different ambient concentrations of their stable isotope or stable element analogue (e.g., for radiocaesium, Blaylock, 1982; Smith et al., 2000) The fish-water concentration factor (*CF*) of ⁹⁰Sr has been shown to vary as an inverse function of the concentration of calcium [*Ca*] (a stable analogue of radiostrontium) in the surrounding water (Vanderploeg et al., 1975; Blaylock, 1982; Kryshev, 2003, 2006). The application of the concentration factor approach usually implies that the uptake of the radioisotope to a fish population has reached equilibrium. Due to the changing ⁹⁰Sr activity concentration in water (e.g. Cross et al., 2002) and the deposit of ⁹⁰Sr in bone, this dynamic accumulation process is to an extent dependent on fish lifespan, but equilibrium is generally considered to have been achieved 8-12 years after radioisotope fallout (Kryshev, 2003). For estimating of ⁹⁰Sr accumulation in fish under non-equilibrium conditions, a number of dynamic models are available, (Sazykina, 2000; Kryshev and Ryabov, 2000, Kryshev, 2003, 2006; Smith et. al., 2005a) though all of these dynamic approaches use estimates of the equilibrium *CF* as one of their input parameters.

The processes which determine the accumulation of radioisotopes in fish are dependent on environmental and biological factors such as water chemistry, trophic level of fish species (predatory or non-predatory fish), fish type and size. ⁹⁰Sr can be accumulated through gills of fish from water (Chowdhury and Blust, 2001) and through the food pathway (Kryshev, 2003). Under low concentrations of the isotope in water, the food pathway is believed to be the more important of the two uptake routes (Michalusev et al, 1997).

Due to its similar bioaccumulation to calcium, approximately 95% of ⁹⁰Sr is found in the bony parts of fish (skeleton, fins, skin) and only 5% in the soft tissues or muscles of a fish (Vanderploeg et al., 1975; Blaylock, 1982). Smith et al. (2005a) assumed that an average of 80% of the wet weight of the fish is composed of soft tissue whilst 20% of the wet weight is bony parts. Another estimation (Shekhanova, 1983) gives an average of 77% of wet weight as soft tissues and 23% as bones. Such differences can influence the accuracy of *CF* calculations where data is presented as separate measurements of ⁹⁰Sr concentration in soft tissues and/or bones. This issue is considered further below.

69

70 The “size effect” of radioisotope accumulation in fish can result in an increasing activity
71 concentration (per unit weight of fish) with increasing fish size (Elliott et al., 1992; Koulikov
72 and Ryabov, 1992; Kryshev and Ryabov, 2000). For radiocaesium, a “size effect” was
73 observed in predatory fish such as perch and pike but no clear dependence was observed for
74 non-predatory fish (roach) (IAEA, 2000; Smith, 2005b). There is, however, less available
75 information on the size effect for radiostrontium: where possible, the database developed in
76 this research will be used to address this question.

77

78 *Previous modelling approaches*

79

80 Two models (Vanderploeg et al., 1975; Kryshev, 2006), based on different empirical data sets
81 have been developed relating the ^{90}Sr CF (in l kg^{-1} , fresh weight) to the water calcium content
82 $[Ca]$ (mg l^{-1}). The inter-comparison and testing (against new empirical data) of these
83 independently-developed models, carried out here, represents a strong test of both models.

84

85 Based on 34 measurements of fish bone-water CF and 19 measurements of fish muscle-water
86 CF , Vanderploeg et al. (1975) (also presented in IAEA, 1994) determined the following
87 relations to estimate the CF of fish (numbers in brackets show uncertainty range):

88

$$89 \quad CF(\text{Muscle}) = \frac{181 (59 - 540)}{[Ca]^{1.6(0.8-1.6)}} \quad (1)$$

90

$$91 \quad CF(\text{Bone}) = \frac{16317}{[Ca]^{1.6(0.8-1.6)}} \quad (2)$$

92

93 Assuming that 23% of wet weight of a fish is composed of bony parts (Shekhanova, 1983),
94 then the whole fish CF is given by:

95

$$96 \quad CF(\text{Whole Fish}) = \frac{3850}{[Ca]^{1.6}} \quad (3)$$

97

98 Using a similar linear regression approach Kryshev (2006) analysed 115 values of the CF at
99 different environmental concentrations of calcium $[Ca^{2+}]$, to obtain the following relationship

100 for whole fish:

$$101 \quad CF(\text{Whole Fish}) = \frac{3940(1770 - 6110)}{[Ca]} \quad (4)$$

102

103 where numbers in brackets show the uncertainty range. An assessment was made (Kryshev,
104 2006) for predatory and non-predatory fish separately and different parameter values were
105 obtained for the two types. The average parameter value was found to be 40 % higher for non-
106 predatory fish than for predatory species, although the confidence intervals overlapped.

107

108 The two models for whole fish CF are very similar: the main difference being in the slope of
109 the inverse power law relationship.

110

111 **2. Methods**

112

113 *Modelling*

114 As in the previous studies, we will model the fish-water CF of radiostrontium as an inverse
115 function of the calcium concentration of the water body (e.g. Blaylock, 1982):

116

$$117 \quad CF = A_1 [Ca]^{-B} \quad (5)$$

118

119 where A_1 and B are parameters to be determined empirically. This model is here called Model
120 1 and the simpler special case of Eq (5) in which B equals 1:

121

$$122 \quad CF = A_2 / [Ca] \quad (6)$$

123

124 will be called Model 2.

125

126 Equations 5 and 6 were fitted to the empirical data using the SAS statistical analysis package
127 (SAS 2002). Prior to fitting, CF and $[Ca]$ data were log-transformed (to give a distribution
128 closer to the normal distribution, and to linearise the relationships) and results back-
129 transformed for presentation. The SAS software (SAS, 2002) gives as output the best-fit
130 model parameters and estimates of 95% confidence interval in those parameter values.

131

132 *Use of previously developed databases*

133

134 The measurements used by Vanderploeg et al. (1975) consisted of 34 measurements of fish

135 bone-water *CF* and 19 measurements of fish muscle-water, together with measurements of
136 [*Ca*]. The later study of Kryshev (2006) consisted of 115 measurements of *CF* in whole fish,
137 but because of the risk of overlap between data sets, 16 measurements obtained by Kryshev
138 (2006) were not used, reducing the data set to 99 measurements. Obviously, all of the
139 Vanderploeg et al. (1975) measurements were pre-Chernobyl and the Kryshev (2006) study
140 used only two data points from freshwater systems contaminated by the Chernobyl accident.
141 For re-analysis, in order to separate the pre- and post-Chernobyl data, these two data points
142 were removed from the Kryshev (2006) data set, leading to a total of 97 data points.

143

144 *Post-Chernobyl CF database for ⁹⁰Sr*

145

146 Post-Chernobyl datasets, from the period 1994-2004, were collected from a literature review
147 (Table 1). They contain observations of ⁹⁰Sr activity concentration in various species of whole
148 fish as well as soft tissues and bones but sometimes only separate measurements in muscle or
149 bony tissue were available. In this case, following Kryshev (2006), whole fish activity
150 concentrations were estimated assuming that 77 % of the weight is soft tissues and 23% in
151 bony tissue. The dataset includes both predatory (pike (*Esox lucius*), perch (*Perca fluviatilis*),
152 pike-perch (*Sander lucioperca*), cat-fish (*Ictalurus punctatus*)) and non-predatory species
153 (roach (*Rutilus rutilus*), tench (*Tinca tinca*), bream (*Abramis brama*), carp (*Cyprinus carpio*),
154 goldfish (*Carassius auratus gibelio*), ruffe (*Gymnocephalus cernuus*)).

155

156 The post-Chernobyl datasets were used as predictive tests of the previously developed
157 models. Due to the time required for equilibration of the ⁹⁰Sr uptake process, only post-1994
158 data were used to test the *CF* models.

159

160 **3. Results and discussion**

161

162 *Comparison of models for ⁹⁰Sr CF in fish*

163

164 Both of the models (Model 1 and its special case, Model 2) were fitted to the available data of
165 *CF* vs. [*Ca*] and estimated parameter values are shown in Table 2. In interpreting these
166 parameter values, note that the value of the *A* parameters depends on the endpoint measured
167 (bone, muscle or tissue), but the value of the *B* parameter is expected to be independent of
168 endpoint measured. Model fits to the empirical data are shown in Figure 1.

169

170 Both models explained a large proportion of the variation in CF values (R^2 values were from
171 65 – 89%). There is some evidence in Table 2 for an inverse power law relationship of slope
172 (“ B ” in Eq. 5) greater than 1, since the analysis of the whole data set estimated B to be 1.11
173 with confidence intervals in the range 1.02 – 1.20. However, the R^2 values of the simpler
174 inverse model (Model 2: $B = 1$) are very close to those of Model 1, so the model which allows
175 “ B ” to be varied offers no major improvement over the simple inverse relationship of Model
176 2.

177

178 *Model testing*

179

180 The two models (Vanderploeg et al., 1975; Kryshev, 2006) were used to predict the whole
181 fish - water CF and muscle-water CF of ^{90}Sr in rivers and lakes impacted by the Chernobyl
182 accident. As shown in Figure 2 (a) and (b) both models generally performed well, showing
183 good agreement with the empirical data. In two out of 12 cases (Braginka River – whole fish;
184 Sozh River – fish muscle), the model predictions were significantly outside the error bars (± 2
185 S.D.) in the empirical data. This may in part have been due to poor estimation of the
186 uncertainty in empirical data since there were in some cases relatively few measurements. In
187 addition, it was not possible, with the available data, to estimate the uncertainty in
188 measurements of the water ^{90}Sr activity concentration.

189

190 It can be seen (Figure 2) that the low calcium Lake Saamia in Finland has significantly higher
191 bioaccumulation of ^{90}Sr than the significantly higher calcium waterbodies in Belarus, Russia
192 and Ukraine.

193

194 *Ratio of ^{90}Sr in bone:muscle tissue*

195

196 The re-analysis of the Vanderploeg et al. (1975) data set gives a best-estimate ratio of 73.3
197 (bone÷muscle activity concentration). This was calculated from the ratio of A_2 values
198 (bone÷muscle) in Table 2. A previous study by Saxén and Koskelainen (2002) measured a
199 significantly higher bone-muscle ratio of 248 (± 59 ; 1 S.D.). In this latter study, all bones,
200 large and small were separated very carefully from the muscle. The lower ratio of
201 Vanderploeg et al. (1975) may be due to inclusion of small bones, skin and/or fins in the
202 “muscle” sample. The higher ratio observed by Saxén and Koskelainen (2002) is likely to be

203 more accurate, but for practical purposes of radiation protection, this lower ratio (i.e. higher
204 predicted activity concentration in “muscle”) may better reflect the “edible” parts of the fish
205 which are typically consumed.

206

207 *“Size effect” on ⁹⁰Sr accumulation by fish*

208

209 The influence of fish size on ⁹⁰Sr accumulation in fish has been studied for the Pripyat and
210 Sozh Rivers and for the Chernobyl Cooling Pond. ⁹⁰Sr activity concentrations in whole fish
211 were plotted as a function of wet weight of fish for each of these systems (Figure 3). Contrary
212 to observations for radiocaesium (Hadderingh et al., 1997; Smith et al., 2002), there was no
213 evidence of a clear “size effect” of increasing ⁹⁰Sr activity concentration with increasing fish
214 weight. None of the relationships observed in Figure 3 showed a statistically significant
215 correlation between ⁹⁰Sr activity concentration and fish weight. It should be noted, however,
216 that the sample sizes were not large, so the ability of the data set to test for a weak size effect
217 relationship is limited. Further, since the measurements we have analysed were made some
218 years after the Chernobyl accident, any effects of differential uptake rates in small and large
219 fish may have been missed. Rapidly changing water activity concentrations in the months and
220 years after the accident could have led at that time to different observed *CF* values in different
221 fish sizes if, for example, equilibrium was more rapidly achieved in small fish than large.

222

223 **3. Conclusions**

224

- 225 1. The previously determined empirical models between *CF* for ⁹⁰Sr accumulation in fish
226 and [Ca²⁺] (mg l⁻¹) in the water are shown to be in good agreement with new
227 measurements for the water bodies affected by the Chernobyl contamination;
- 228 2. On the basis of the available data, no significant relationships between fish size and
229 ⁹⁰Sr activity concentration in fish (the “size effect”) were determined;
- 230 3. Remaining variation in *CF* not explained by an inverse relationship with [Ca²⁺] is
231 significant, but is likely to be due to a number of factors such as fish feeding
232 behaviour, recruitment and population age which may be difficult to predict using
233 general (as opposed to lake- or river- specific) models.

Acknowledgments

Nataliya Sasina would like to acknowledge the support of INTAS – 2005 Young Scientist Fellowship (Ref. number 05-109-4221). The study was also supported in part by project RFBR no. 07-04-00002.

References

- Blaylock, B.G., 1982 Radionuclide data bases available for bioaccumulation factors for freshwater biota. *Nucl. Safety*. 23, 427-438.
- Chowdhury M.J., Blust, R., 2001. A mechanistic model for the uptake of waterborne strontium in the common carp (*Cyprinus carpio* L). *Environ. Sci. Technol.* 35, 669-75.
- Cross, M.A., Smith, J.T., Saxén, R., Timms, D.N., 2002. An analysis of the time dependent environmental mobility of radiostrontium in Finnish river catchments. *J. Environ. Radioactivity*. 60, 149-163.
- Elliott, J.M., Hilton, J., Rigg, E., Tullett, P.A., Swift, D.J., Leonard, D.R.P., 1992. Sources of variation in post-Chernobyl radiocaesium in fish from two Cumbrian lakes (north-west England). *J. Appl. Ecol.* 29, 108-119.
- Haddingh, R. H., van Aerssen, G. H. F. M., Ryabov, I. N., Koulikov, O. A., Belova, N., 1997. Contamination of fish with ^{137}Cs in Kiev reservoir and old river bed of Pripjat near Chernobyl, in Desmet, G., Blust, R., Comans, R.N.J., Fernandez, J., Hilton, J., de Bettencourt, A. (Eds.), *Freshwater and estuarine radioecology*. Elsevier, The Netherlands, 339-351.
- International Atomic Energy Agency, 1994. Handbook of parameter values for the prediction of radionuclide transfer in temperate environments. IAEA Technical Reports Series No. 364, International Atomic Energy Agency, Vienna.
- International Atomic Energy Agency, 2000. Modelling of the transfer of radiocaesium from deposition to lake ecosystems. IAEA-TECDOC-1143. International Atomic Energy Agency, Vienna.
- Koulikov, A.O, Ryabov, I.N., 1992. Specific cesium activity in freshwater fish and size effect. *Sci. Tot. Environ.* 112, 125-142.
- Kryshev, A.I., Ryabov, I.N., 2000. A dynamic model of ^{137}Cs accumulation by fish of different age classes. *J. Environ. Radioactivity*. 50, 221 – 233.
- Kryshev, A.I., 2003. Model reconstruction of ^{90}Sr concentrations in fish from 16 Ural lakes contaminated by the Kyshtym accident of 1957. *J. Environ. Radioact.* 64, 67-84.
- Kryshev, A.I., 2006. ^{90}Sr in fish: a review of data and possible model approach. *Sci. Tot Environ.* 370, 182-189.
- Kryshev, I.I., 1991. I.I. Kryshev, (Ed.) *Radioecological Consequences of the Chernobyl Accident*. Kurchatov Institute of Atomic Energy, Moscow.
- Michalusev V.I., Gulakov A.V., Tolkachev A.P., Prochorchik A.P. Uptake of radionuclides by freshwater fishes after the Chernobyl accident. Collection of papers of the scientific-research association "Fauna Polecie". Issue 2. Gomel (Belarus). 1997. p.83-90 (in Russian).

- Mück, K., Pröhl, G., Likhtarev, I., Kovgan, L., Meckbach, R., Golikov, V. A., 2002. A consistent radionuclide vector after the Chernobyl accident. *Health Phys.* 82, 141-151.
- Pankov I.V., Volkova E.N., Shirokaya Z.O. (1993) Radioecological investigations within littoral zone of the Kiev reservoir before and after the Ch NPP accident. *Hydrobiology Journal*, 29(3), p.100-109 (in Russian).
- SAS, 2002. SAS 9.1.3 Help and Documentation. SAS Institute Inc., Cary, NC.
- Sasina, N.V., Smith, J.T., Kudelsky, A.V., Wright, S.M., 2007. “Blind” testing of models for predicting the ^{90}Sr activity concentration in river systems using post-Chernobyl monitoring data. *J. Environ. Radioactivity.* 92; 63-71.
- Saxén, R., Koskelainen, U., 2000. Distribution of ^{137}Cs and ^{90}Sr in various tissues and organs of freshwater fish in Finnish lakes. *Boreal Environ. Res.* 7, 105-112.
- Saxén, R., Koskelainen, U., 2001. Effect of site-specific parameters on the transfer of ^{137}Cs and ^{90}Sr into freshwater fishes. *Radiochem.* 43, 487-491.
- Sazykina, T.G., 2000. ECOMOD – An ecological approach to radio-ecological modelling. *J. Environ. Radioactivity.* 50, 207-20.
- Shekhanova, I.A., 1983. Radioecology of fish. Food Industry: Moscow: in Russian.
- Smith, J.T., Kudelsky, A.V., Ryabov, I.N., Hadderingh, R.H., 2000. Radiocaesium concentration factors of Chernobyl-contaminated fish: a study of the influence of potassium and “blind” testing of a previously developed model. *J. Environ. Radioactivity.* 48, 359-369.
- Smith, J.T., Kudelsky, A.V., Ryabov, I.N., Daire, S.E., Boyer, L., Blust, R.J., Fernandez, J.A., Hadderingh, R.H., Voitsekhovitch, O.V., 2002. Uptake and elimination of radiocaesium in fish and the “size effect”. *J. Environ. Radioactivity.* 62, 145-164.
- Smith, J.T., Belova, N.V., Bulgakov, A.A., Comans, R.N.J., Konoplev, A.V., Kudelsky, A.V., Madruga, M.J., Voitsekhovich, O.V., Zibold, G., 2005a. The “AQUASCOPE” simplified model for predicting $^{89,90}\text{Sr}$, ^{131}I , and $^{134,137}\text{Cs}$ in surface waters after a large-scale radioactive fallout. *Health Phys.* 89, 628-644.
- Smith, J.T., Beresford, N.A., 2005b. Chernobyl – catastrophe and consequences. Springer-Praxis, Berlin.
- Vanderploeg, H. A., Parzyck, D.C., Wilcox, W. H., Kerchner, J.R., Kaye, S. V., 1975. Bioaccumulation Factors for Radionuclides in Freshwater Biota. ORNL-5002, Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Figure captions

Figure 1. Fit of Model 2 to (a) Vanderploeg et al. (1975) data; (b) Kryshev (2006) data; (c) combined data set.

Figure 2. Test of Vanderploeg et al. (1975) and Kryshev (2006) models against post-Chernobyl data for (a) whole fish (Vanderploeg model: $CF=3850 \times [Ca]^{-1.2}$; Kryshev model: $CF=3610 \times [Ca]^{-1}$) and; (b) fish muscle (Vanderploeg model: $CF=181 \times [Ca]^{-1.2}$).

Figure 3. Relationships between ^{90}Sr concentration factor and fish weight for various predatory and non-predatory species in (a) Pripyat River (Choiniki); (b) Sozh River (Gomel); (c) Chernobyl Cooling Pond.

Table 1. Measurements of ^{90}Sr concentration factors in whole fish after Chernobyl

Water body	Number of observations	Sampling date	Calcium concentration in water, mg l^{-1}	<i>CF</i> Muscle	<i>CF</i> Whole fish	References
River Dnieper (Bragin)	4	1994-95	40.3	-	36.3	(1)
River Sozh (Gomel)	17	1994-95	48.9	2.59	159	(1)
River Braginka (Bragin)	12	1994-95	55.7	5.11	7.96	(1)
Lake Perstok	6	2002	26.45	-	250	(2)
Lake Kozhanovskoe	*	1994	33	-	173	(3)
Chernobyl Cooling Pond	27	2002-04	50.5	2.88	38.6	(3)
Lake Glubokoye	10	2003-04	27.2	5.52	202	(3)
Lake Saimaa, Finland	4	1994	3	-	455	(4)

1. Michalusev et al., 1997; 2. Smith et al., 2005b; 3. Belova N.V., Severtsov Institute, Moscow, unpubl. res.; 4. Saxén R., Koskelainen U., 2001. * This given as a mean value, but the no. of observations is not known.


Table 2. Parameter values determined by fitting Models 1 and 2 to the data sets of Vanderploeg et al. (1975) and Kryshev (2006).

Data set	No. of obs.	Model 1 CF = $A_1[Ca]^{-B}$			Model 2 CF = $A_2[Ca]^{-1}$	
		A_1	B	R^2	A_2	R^2
Vanderploeg (bone)	35	13430 (8913 – 20000)	1.12 (1.26 – 0.99)	89%	9750 (8110 – 11700)	88%
Vanderploeg (muscle)	19	204 (57.1 – 727)	1.16 (1.59 – 0.73)	66%	133 (77.6 – 231)	65%
Kryshev (whole fish)	97	6412 (4450 – 9238)	1.18 (1.29 – 1.07)	82%	3610 (3231 – 4034)	80%
All data (whole fish)	132	4511 (3385 – 6013)	1.11 (1.20 – 1.02)	82%	3224 (2923 – 3557)	81%


Table 3. Test for relationships between ^{90}Sr activity concentration (whole fish, f.w.) and fish weight. None of the correlations was statistically significant.

Water body (period of study)	Fish Type	R^2 value
River Pripyat (1994-5)	Predatory	0.016 (n=13)
River Sozh (1994-5)	Predatory	0.042 (n = 7)
Chernobyl Cooling Pond (2002-4)	Predatory	0.24 (n = 8)
River Pripyat (1994-5)	Non-Predatory	0.094 (n = 11)
River Sozh (1994-5)	Non-Predatory	0.121 (n = 10)
Chernobyl Cooling Pond (2002-4)	Non-Predatory	0.026 (n = 19)

(a) Vanderploeg et al. (1975) data



(b) Kryshev (2006) data



(c) Complete data set




Figure 1





Figure 2




Figure 3