

**Keywords**Pesticides residues, GC-MS/MS,  
Food Safety, Environmental  
contaminants**CORRESPONDING AUTHOR****Giuseppe Labella**  
giuseppe.labella@unimi.it**JOURNAL HOME PAGE**[riviste.unimi.it/index.php/haf](http://riviste.unimi.it/index.php/haf)UNIVERSITÀ DEGLI STUDI DI MILANO  
DIPARTIMENTO DI SCIENZE VETERINARIE  
PER LA SALUTE, LA PRODUZIONE ANIMALE  
E LA SICUREZZA ALIMENTARE

## Analyses of organochlorine pesticides residues in eels (*Anguilla anguilla*) from Lake Garda using Gas chromatography coupled with Tandem Mass Spectrometry (GC-MS/MS)

Giuseppe F. Labella<sup>1\*</sup>, Luca M. Chiesa<sup>1</sup>, Sara Panseri<sup>1</sup>,  
Francesco Arioli<sup>1</sup><sup>1</sup>University of Milan, Department of Health, Animal Science and Food Safety, Italy

### Abstract

Lake Garda is located in one of the most populated and industrialized area of Italy (Camusso *et al.*, 2001). Therefore, the Lake water, and also the fish species present, could be affected by environmental contamination. European eels (*Anguilla anguilla*) are considered as suitable matrix for biomonitoring environmental contaminants in European water (Belpaire *et al.*, 2007), being widespread in many European waters and highly contaminated by lipophilic compounds, due to the high lipid content (up to 40%) (Larsson *et al.*, 1991). Moreover, eel is an edible species (its farming currently supplies approximately 45,000 tons/year) (Nielsen *et al.*, 2008), so it also represents a public health issue. Based on these considerations, the aim of this study was to evaluate the occurrence of fourteen organochlorine pesticides (OCs) in forty-five eels (*Anguilla anguilla*) from Lake Garda, using Accelerated Solvent Extraction (ASE) procedure for the analytes extraction and Gas chromatography coupled with Tandem Mass Spectrometry (GC-MS/MS) for the analysis of OCs. GC-MS/MS analysis was developed and validated according to the SANTE/11945/2015 guidelines. Uncontaminated eel sample (previously checked for the presence of OCs and considered blank with a concentration of compounds < Limit of Detection) were used for all procedure's optimization steps. For all the OCs, satisfactory results were achieved in terms of linearity ( $R_2$  higher than 0.985); recovery (ranging between 70–120 %) and repeatability (coefficient of variation % lower than 20 %). The results met the validation criteria required by EU guidelines. Regarding eel samples, several pesticides were detected, but DDT and its metabolites were found with the highest prevalence (92 %). The concentration range was from not detected (n.d.) to 19000 ng g<sup>-1</sup>. Although DDTs levels in the environment are declining (Albaiges *et al.*, 2011), they continue to bioaccumulate in tissues of human and animal and biomagnify in food chains.

## References

Camusso, M., Balestrin, R., Binelli, A. 2001. Use of zebra mussel (*Dreissena polymorpha*) to assess trace metal contamination in the largest Italian subalpine lakes. *Chemosphere*. 44, 263-270.

Belpaire, C., Goemans, G. 2007. The European eel *Anguilla anguilla*, a rapporteur of the chemical status for the water framework directive? *Vie Milieu*. 57, 235–252.

Larsson, P., Hamrin, S., Okla, L. 1991. Factors determining the uptake of persistent pollutants in a eel population (*Anguilla anguilla* L.). *Environ Pollut*. 69, 39-50.

Nielsen, T., Prouzet, P. 2008. Capture-based aquaculture of the wild European eel (*Anguilla anguilla*). In A. Lovatelli & P. F. Holthus (Eds.), *Capture-based aquaculture. Global overview*. FAO Fisheries Technical Paper. No. 508 (pp. 141–168). Rome: FAO.

Albaiges, J., Murciano, C., Pon, J., 2011. Hazardous substances in the Mediterranean: a spatial and temporal assessment. In: UNEP/MAP, *Consultation Meeting to Review MED POL Monitoring Activities Athens, 22 e 23 November 2011*, p. 106. Annex II.