

DIAS-STP-04-02

Two Order Parameters in Quantum XZ Spin Models with Gibbsian Ground States

T. C. Dorlas

*Dublin Institute for Advanced Studies
School of Theoretical Physics
10 Burlington Road, Dublin 4, Ireland.
Email: dorlas@stp.dias.ie*

W. Skrypnik

*Institute of Mathematics of NANU
Kyiv, Ukraine.
Email: skrypnik@imath.kiev.ua*

March 8, 2004

Abstract

We describe a family of quantum spin models which are generators of a discrete Markovian process. We show that there exists an explicit expression for the ground state of such models and give a simple argument for the existence of two types of long-range order in such systems. Two special examples of these systems are analysed in detail.

1 Introduction

The existence of long-range order for order parameters in quantum many-body systems is an important problem which is the first step towards a complete description of the phase diagram.

This problem has been solved for a large class of quantum spin systems of the mean-field type. These models include the Vonsovsky-Zener type fermion-spin systems [1] explaining the occurrence of superconductivity and of ferromagnetism at non-zero temperatures. The first rigorous analysis [1–3] of such systems made use of the so-called approximating Hamiltonian method. Other methods include large-deviation theory combined with group representations [4–7] and C*-algebra analysis [8–10]. Note also that the approximating Hamiltonian method has been extended to boson systems in [11] and [12].

Tian [21] formulated a sufficient condition for the coexistence of two independent order parameters with long-range order in the ground state of some boson and fermion systems. For the Hubbard model this condition coincides with the RVB (resonating valence bond) long-range order and on-site-pairing long-range order. Macris and Piguet [20] proved the existence of two order parameters for lattice boson-fermion systems at a non-zero temperature by generalizing [19] the Tian technique in and the Lieb-Simon reflection-positivity technique.

In this paper we formulate a special class of quantum spin XZ models on the hypercubic lattice \mathbb{Z}^d with a Gibbsian ground state in which long-range order occurs for the spin operators S^1 and S^3 in dimensions greater than one. (In one-dimensional systems ferromagnetic long-range order for S^1 is easy to prove.)

Our systems differ from the XZ spin $\frac{1}{2}$ systems which admit Gibbsian ground states considered in [15]. There, the classical Gibbsian system which generates the ground state is in fact quite complicated. Kirkwood and Thomas proved that there is ferromagnetic long-range order for S^3 in the ground state in some of their ferromagnetic systems. Our proof of the S^1 -long-range order is analogous to theirs. In [16] the Kirkwood-Thomas analysis is formulated as a fixed-point problem and applied to find quasi-particle states. The method has been further generalised by Yarotsky [17]. Our analysis is less general but has the advantage of simplicity.

In [18], Matsui showed that in one dimension, classical Gibbsian systems are associated with quantum Potts systems. The structure of the Matsui Hamiltonians are a special case of the Hamiltonians of XZ spin systems considered here, which can be represented as a sum of a diagonal part of a specific form and an Ising-type non-diagonal part.

Our Hamiltonians are expressed in terms of the Pauli matrices

$$S^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad S^2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \text{and } S^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (1.1)$$

Given a finite subset $\Lambda \subset \mathbb{Z}^d$ with cardinality $|\Lambda|$ let S_x^1 etc. be the corresponding operators on $\mathbb{E}_\Lambda = (\mathbb{C}^2)^\Lambda$ acting on the factor for the point $x \in \Lambda$. If we denote for $s_\Lambda \in \{-1, 1\}^\Lambda$,

$$\Psi_\Lambda^0(s_\Lambda) = \otimes_{x \in \Lambda} \psi_0(s_x), \quad \text{where } \psi_0(1) = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \psi_0(-1) = \begin{pmatrix} 0 \\ 1 \end{pmatrix},$$

then this can be written as

$$S_x^1 \Psi_\Lambda^0(s_\Lambda) = \Psi_\Lambda^0(s_\Lambda^{\{x\}}), \quad S_x^3 \Psi_\Lambda^0(s_\Lambda) = s_x \Psi_\Lambda^0(s_\Lambda), \quad (1.2)$$

where, for any subset $A \subset \Lambda$, s_Λ^A is the configuration s_Λ with the spins in A flipped. (Note that the states $\Psi_\Lambda^0(s_\Lambda)$ form an orthonormal basis for $(\mathbb{C}^2)^\Lambda$. In particular,

$$\langle \Psi_\Lambda^0(s_\Lambda) | \Psi_\Lambda^0(s'_\Lambda) \rangle = \delta(s_\Lambda; s'_\Lambda) = \prod_{x \in \Lambda} \delta_{s_x, s'_x},$$

where δ_{s_x, s'_x} is the Kronecker symbol.)

We now define the operators

$$P_A = S_A^1 - e^{-\frac{\alpha}{2} W_A(S_A^3)}, \quad S_A^1 = \prod_{x \in A} S_x^1, \quad (1.3)$$

where

$$W_A(s_\Lambda) = U_0(s_\Lambda^A) - U_0(s_\Lambda), \quad U_0(s_\Lambda^A) = U_0(s_{\Lambda \setminus A}, -s_A). \quad (1.4)$$

Our main results concern Hamiltonians of the form

$$H_\Lambda = \sum_{A \subset \Lambda} J_A P_A, \quad J_A \leq 0 \quad (1.5)$$

In Theorem 2.1 below, we show that their ground state is given by

$$\Psi_\Lambda = \sum_{s_\Lambda} e^{-\frac{\alpha}{2} U_0(s_\Lambda)} \Psi_\Lambda^0(s_\Lambda), \quad \alpha \in \mathbb{R}^+. \quad (1.6)$$

In the proof we establish that the Hamiltonian (1.5) is the generator of a discrete Markovian process. The spectral structure for such generators in

the simplest case ($|A| = 1$) was established in [22]. In Theorem 2.2, we formulate conditions on J_A for which this ground state is unique. As a simple consequence, we show in Theorem 2.3 that in dimensions $d > 1$, there are two types of long-range order in these systems.

In the third section we calculate explicit expressions for the Hamiltonians in the case $J_A = 0, |A| > 2$ and with the simplest choice of a ferromagnetic U_0 . The Hamiltonian corresponding to the case $d = 1, J_A = 0, |A| > 1$ already appeared in [Ma]. The case $J_A = 0, |A| \neq 2$ is interesting since our Hamiltonian is expressed as a perturbation of the simple ferromagnetic Hamiltonian

$$H_\Lambda = J \sum_{\langle x,y \rangle \in \Lambda} (S_x^1 S_y^1 + \gamma S_x^3 S_y^3), \quad J < 0,$$

where $\gamma = 4d(\cosh \alpha)^{4d-3} \sinh \alpha$. Our condition of uniqueness of the ground state does not apply to this case since it does not hold if $J_A = 0$ for all A with $|A| \neq 2$. However, see Remark 2.2.

Remark. The class of Hamiltonians for which (1.6) is a ground state can be generalised to

$$H_\Lambda = \sum_{A_1, \dots, A_l \subset \Lambda} J_{A^{(l)}} (P_{A_1} \dots P_{A_l} + P_{A_l} \dots P_{A_1}), \quad A^{(l)} = (A_1, \dots, A_l), \quad (1.7)$$

where the summation is over families of disjoint non-empty subsets of Λ . This follows from the following equality for an arbitrary A

$$P_A \Psi_\Lambda = 0. \quad (1.8)$$

2 Main results

We first prove that (1.6) is a ground state with eigenvalue zero for the Hamiltonian (1.5):

Theorem 2.1 *The Hamiltonian (1.5) is a positive self-adjoint operator on $(\mathbb{C}^2)^\Lambda$ and the state Ψ_Λ , given by (1.6), is a ground state with eigenvalue zero.*

We begin by proving (1.8). This shows that Ψ_Λ is an eigenfunction of the Hamiltonian (1.5) with eigenvalue zero. The identity (1.8) follows easily by

changing signs of the spin variables s_A in the first term:

$$\begin{aligned}
P_A \Psi_\Lambda &= \sum_{s_\Lambda} (\Psi_\Lambda^0(s_\Lambda^A) - e^{-\frac{\alpha}{2}W_A(s_\Lambda)} \Psi_\Lambda^0(s_\Lambda)) e^{-\frac{\alpha}{2}U_0(s_\Lambda)} \\
&= \sum_{s_\Lambda} (\Psi_\Lambda^0(s_\Lambda^A) e^{-\frac{\alpha}{2}U_0(s_\Lambda)} - \Psi_\Lambda^0(s_\Lambda) e^{-\frac{\alpha}{2}U_0(s_\Lambda^A)}) \\
&= \sum_{s_\Lambda} \left(e^{-\frac{\alpha}{2}U_0(s_\Lambda^A)} - e^{-\frac{\alpha}{2}U_0(s_\Lambda)} \right) \Psi_\Lambda^0(s_\Lambda) = 0.
\end{aligned}$$

Next we prove that the Hamiltonian is a positive operator. For this purpose, we define two further operators

$$H_\Lambda^+ = e^{\frac{\alpha}{2}U_0(S_\Lambda^3)} H_\Lambda e^{-\frac{\alpha}{2}U_0(S_\Lambda^3)}, \quad H_\Lambda^- = e^{-\frac{\alpha}{2}U_0(S_\Lambda^3)} H_\Lambda e^{\frac{\alpha}{2}U_0(S_\Lambda^3)}. \quad (2.9)$$

It is clear that

$$(H_\Lambda^+)^* = H_\Lambda^-, \quad H_\Lambda^- = e^{-\alpha U_0(S_\Lambda^3)} H_\Lambda^+ e^{\alpha U_0(S_\Lambda^3)}. \quad (2.10)$$

where the star denotes the adjoint in the Hilbert space $\mathbb{E}_\Lambda = (\mathbb{C}^2)^\Lambda$.

A straightforward calculation on the basis Ψ_Λ^0 shows that

$$H_\Lambda^+ = \sum_{A \subseteq \Lambda} J_A e^{-\frac{\alpha}{2}W_A(S_\Lambda^3)} (S_A^1 - I), \quad (2.11)$$

where I is the unit operator. This operator is symmetric with respect to the new scalar product

$$\langle F' | F \rangle_{U_0} = \langle F' | e^{-\alpha U_0(S_\Lambda^3)} F \rangle. \quad (2.12)$$

Indeed,

$$\begin{aligned}
\langle F' | H_\Lambda^+ F \rangle_{U_0} &= \langle F' | e^{-\alpha U_0(S_\Lambda^3)} H_\Lambda^+ F \rangle \\
&= \sum_{A \subseteq \Lambda} J_A \langle F' | e^{-\frac{\alpha}{2}[U_0(S_\Lambda^3) + U_0(S_\Lambda^{3A})]} (S_A^1 - I) F \rangle \\
&= \sum_{A \subseteq \Lambda} J_A \langle (S_A^1 - I) F' | e^{-\frac{\alpha}{2}[U_0(S_\Lambda^3) + U_0(S_\Lambda^{3A})]} F \rangle \\
&= \langle H_\Lambda^+ F' | F \rangle_{U_0}.
\end{aligned}$$

Here we used the equalities

$$e^{-\frac{\alpha}{2}U_0(S_\Lambda^3)} S_A^1 = S_A^1 e^{-\frac{\alpha}{2}U_0(S_\Lambda^{3A})}, \quad e^{-\frac{\alpha}{2}U_0(S_\Lambda^{3A})} S_A^1 = S_A^1 e^{-\frac{\alpha}{2}U_0(S_\Lambda^3)} \quad (2.13)$$

From these inequalities we derive, also,

$$\langle F' | H_\Lambda^+ F \rangle_{U_0} = \langle e^{-\frac{\alpha}{2}U_0(S_\Lambda^3)} F' | H_\Lambda e^{-\frac{\alpha}{2}U_0(S_\Lambda^3)} F \rangle. \quad (2.14)$$

This shows that it suffices to prove that H_Λ^+ is a positive operator for the new scalar product (2.12). Let

$$F = \sum_{s_\Lambda} F(s_\Lambda) \Psi_\Lambda^0(s_\Lambda);$$

then

$$(H_\Lambda^+ F)(s_\Lambda) = - \sum_{A \subseteq \Lambda} J_A e^{-\frac{\alpha}{2} W_A(s_\Lambda)} (F(s_\Lambda) - F(s_\Lambda^A)). \quad (2.15)$$

In deriving this equality one has to once again change the signs of the spins s_A in the expansion of $H_\Lambda^+ F$ on the basis Ψ_Λ^0 .

This means that

$$\begin{aligned} \langle F | H_\Lambda^+ F \rangle_{U_0} &= - \sum_{A \subseteq \Lambda} J_A \sum_{s_\Lambda} e^{-\frac{\alpha}{2} [U_0(s_\Lambda) + U_0(s_\Lambda^A)]} (F(s_\Lambda) - F(s_\Lambda^A)) F(s_\Lambda) \\ &= - \frac{1}{2} \sum_{A \subseteq \Lambda} J_A \sum_{s_\Lambda} e^{-\frac{\alpha}{2} [U_0(s_\Lambda) + U_0(s_\Lambda^A)]} (F(s_\Lambda) - F(s_\Lambda^A))^2 \geq 0. \end{aligned} \quad (2.16)$$

Here we used the fact that the exponential weight in the sum is invariant under changing signs of spin variables s_A . It now follows that H_Λ is positive definite.

Remark 2.1 The operator H_Λ^+ is an analog of the operator generated by the Dirichlet form for continuous spins [23]. Its exponent $e^{-tH_\Lambda^+}$ generates a discrete Markov process which can be called a generalized spin-flip process. For its adjoint the following relations are valid

$$(H_\Lambda^- F)(s_\Lambda) = \sum_{A \subseteq \Lambda} J_A [e^{\frac{\alpha}{2} W_A(s_\Lambda)} F(s_\Lambda^A) - e^{-\frac{\alpha}{2} W_A(s_\Lambda)} F(s_\Lambda)], \quad \sum_{s_\Lambda} (H_\Lambda^- F)(s_\Lambda) = 0.$$

The last equality implies the validity of the law of conservation of probability and is derived after changing signs of spins s_A in the first term of the first equality ($W_A(s_\Lambda^A) = -W_A(s_\Lambda)$).

Uniqueness of the ground state will be derived from the Perron-Frobenius Theorem [13, 14]:

Theorem *Let the square matrix B be non-negative and irreducible. Then the spectral radius $\rho(B)$ is a simple eigenvalue of B and $\rho(B) > 0$. Moreover, the components of the associated eigenvector are all strictly positive.*

We recall that a matrix is non-negative if all its matrix elements are non-negative, and an $n \times n$ -matrix B is irreducible if there does not exist a subset $I \subset \{1, \dots, n\}$ such that for all $(i, j) \in I \times I^c$, the matrix elements $B_{i,j} = 0$.

We use this theorem to derive two alternative conditions for uniqueness of the ground state:

Theorem 2.2 *The ground state Ψ_Λ of H_Λ is unique if one of the following conditions is satisfied:*

1. $J_{\{x\}} < 0$ for all $x \in \Lambda$; or
2. For every pair of points $x, y \in \Lambda$ there exists a chain $x_0 = x, x_1, \dots, x_n = y$ of points in Λ such that $J_{\{x_i, x_{i+1}\}} < 0$ and there is set $A \subset \Lambda$ with $J_A < 0$ and $|A|$ odd.

Proof. We apply the Perron-Frobenius Theorem to the operator $-H_\Lambda + aI$, where I is the identity operator (matrix) and a is a constant given by

$$a = \sum_{A \subset \Lambda} J_A e^{-\frac{a}{2} W_A(s_\Lambda)}. \quad (2.17)$$

Consider first the case $J_{\{x\}} < 0$ for all $x \in \Lambda$. Suppose that $I \subset \{-1, 1\}^\Lambda$ is such that

$$\langle \Psi_\Lambda^0(s'_\Lambda) | (-H_\Lambda + aI) \Psi_\Lambda^0(s_\Lambda) \rangle = - \sum_{A \subset \Lambda} J_A \langle \Psi_\Lambda^0(s'_\Lambda) | S_A^1 \Psi_\Lambda^0(s_\Lambda) \rangle = 0 \quad \forall s_\Lambda \in I, s'_\Lambda \in I^c. \quad (2.18)$$

Since $I \neq \{-1, 1\}^\Lambda$, there exists $s_\Lambda \in I$ and $x \in \Lambda$ such that $s'_\Lambda := S_x^1 \Psi_\Lambda^0(s_\Lambda) = \Psi_\Lambda^0(s_\Lambda^{\{x\}}) \notin I$. This contradicts (2.18) since all $J_A \leq 0$ and $J_{\{x\}} < 0$.

Next consider case 2, and assume again that (2.18) holds. Similar to the previous case, if $s_\Lambda \in I$ and $x, y \in \Lambda$ such that $J_{\{x,y\}} < 0$ then $s_\Lambda^{\{x,y\}} \in I$. By flipping pairs of spins in a chain as in the hypothesis, it then follows that we can flip any pair of spins in s_Λ . We conclude that I must contain all configurations with an even number of spins $s_x = -1$ or all configurations with an odd number of minus-spins. However, it is also assumed that there is a set $A \subset \Lambda$ with $|A|$ odd and $J_A < 0$. Flipping the spins in A converts a configuration with an odd number of spins $s_x = -1$ to one with an even number and vice versa. It follows that I must contain all configurations.

Remark 2.2 The second condition in case 2 is not superfluous: it follows from the proof that even if $J_A < 0$ for all A with $|A| = 2$, there does exist

a nontrivial set I satisfying (2.18). Indeed, in this case the spaces spanned by $\Psi_\Lambda^0(s_\Lambda)$ where $\#\{x : s_x = -1\}$ is odd resp. even are invariant, and the ground state is two-fold degenerate.

One of the most interesting features of the models considered is that they have two order parameters with long-range order. This is now surprisingly easy to prove:

Define, for finite subsets $A \subset \mathbb{Z}^d$, and operators F_A depending on S_x^1 , S_x^2 and S_x^3 with $x \in A$,

$$\langle F_A \rangle = \lim_{\Lambda \rightarrow \mathbb{Z}^d} \langle F_A \rangle_\Lambda, \quad \langle F_A \rangle_\Lambda = \frac{\langle \Psi_\Lambda | F_A \Psi_\Lambda \rangle}{\langle \Psi_\Lambda, \Psi_\Lambda \rangle}, \quad (2.19)$$

where Ψ_Λ is the ground state. The Gibbsian nature of the ground state then immediately yields the following theorem.

Theorem 2.3 *Suppose that the Hamiltonian H_Λ of a quantum spin system on finite subsets of the lattice \mathbb{Z}^d is given by (1.5) and that $\lim_{\Lambda \rightarrow \mathbb{Z}^d} W_A(s_\Lambda)$ exists for all finite $A \subset \mathbb{Z}^d$. Suppose moreover that the limit is bounded if $|A| = 2$. Then, for $d \geq 1$, there is ferromagnetic long-range order for S^1 . Moreover, if there is long-range order in the corresponding classical spin system with the potential energy U_0 then such long-range order occurs also for S^3 in the ground state of the quantum system.*

Proof. We have to prove that

$$\langle S_x^1 S_y^1 \rangle > a, \text{ for } a > 0. \quad (2.20)$$

Writing

$$Z_\Lambda = \langle \Psi_\Lambda | \Psi_\Lambda \rangle = \sum_{s_\Lambda} e^{-\frac{\alpha}{2} U_0(s_\Lambda)}.$$

we have

$$\langle S_x^1 S_y^1 \rangle_\Lambda = Z_\Lambda^{-1} \sum_{s_\Lambda} e^{-\frac{\alpha}{2} U_0(s_\Lambda)} e^{-\frac{\alpha}{2} W_{x,y}(s_\Lambda)} \geq \inf_{s_\Lambda, x, y} e^{-\frac{\alpha}{2} W_{x,y}(s_\Lambda)} < +\infty.$$

This proves (2.20).

Since S^3 is a diagonal matrix, the ground state expectation value of a function of S_x^3 equals the classical Gibbsian expectation value of the function depending on classical spins. This proves the last statement of the theorem.

Remark 2.3 For short range interactions the condition for $W_{x,y}$ of the theorem is always satisfied. It is well-known that for a ferromagnetic nearest-neighbour pair interaction

$$U_0(s_\Lambda) = -g \sum_{\langle x,y \rangle \subseteq \Lambda} s_x s_y \quad (g > 0), \quad (2.21)$$

there is ferromagnetic long-range order in the classical system at sufficiently low temperatures.

3 Examples

In this section we show that some of the the Hamiltonians considered in the previous section have the following form

$$H_\Lambda = \tilde{H}_\Lambda + H_{\partial\Lambda} + |\Lambda|\alpha_0, \quad (3.22)$$

where \tilde{H}_Λ is a polynomial in S_x^1 and S_x^3 , $H_{\partial\Lambda}$ is a boundary term, and α_0 is a constant.

We consider two specific examples.

3.1 Example 1

Put $J_x = -1$; $J_{x_1, \dots, x_k} = 0, k > 1$ and

$$U_0(s_\Lambda) = - \sum_{\langle x,y \rangle \in \Lambda} s_x s_y. \quad (3.23)$$

Then

$$W_x(s_\Lambda) = 2s_x \sum_{y \in \Lambda, |y-x|=1} s_y. \quad (3.24)$$

Let n_x be the number of nearest neighbours of x . Then from the simple equality

$$e^{-\alpha S} = \cosh \alpha - S \sinh \alpha, \quad S^2 = I, \quad (3.25)$$

it follows that $(Y_k = (y_1, \dots, y_k))$

$$\begin{aligned} e^{-\frac{\alpha}{2} W_x(S_\Lambda^3)} &= \prod_{y \in \Lambda, |y-x|=1} e^{-\alpha S_x^3 S_y^3} \\ &= \prod_{y \in \Lambda, |y-x|=1} (\cosh \alpha - S_x^3 S_y^3 \sinh \alpha) \end{aligned}$$

$$\begin{aligned}
&= \sum_{k=1}^{\lfloor \frac{n_x}{2} \rfloor} (\sinh \alpha)^{2k} (\cosh \alpha)^{n_x - 2k} \sum_{Y_{2k} \subset \Lambda, |y_j - x| = 1} S_{[Y_{2k}]}^3 \\
&\quad - S_x^3 \sum_{k=0}^{\lfloor \frac{n_x - 1}{2} \rfloor} (\sinh \alpha)^{2k+1} (\cosh \alpha)^{n_x - 2k - 1} \sum_{Y_{2k+1} \subset \Lambda, |y_j - x| = 1} S_{[Y_{2k+1}]}^3 \\
&\quad + (\cosh \alpha)^{n_x},
\end{aligned}$$

where $[n]$ is the integer part of the number n . The Hamiltonian can therefore be written as

$$\begin{aligned}
H_\Lambda &= - \sum_{x \in \Lambda} \left\{ S_x^1 - \sum_{k=1}^{\lfloor \frac{n_x}{2} \rfloor} \alpha_k(n_x) \sum_{Y_{2k} \subset \Lambda, |y_j - x| = 1} S_{[Y_{2k}]}^3 + \right. \\
&\quad \left. + \sum_{k=0}^{\lfloor \frac{n_x - 1}{2} \rfloor} \beta_k(n_x) \sum_{Y_{2k+1} \subset \Lambda, |y_j - x| = 1} S_x^3 S_{[Y_{2k+1}]}^3 \right\} + (\cosh \alpha)^{2d} |\Lambda| - c_{\partial\Lambda},
\end{aligned}$$

where

$$\alpha_k(n) = (\sinh \alpha)^{2k} (\cosh \alpha)^{n - 2k},$$

and

$$\beta_k(n) = (\sinh \alpha)^{2k+1} (\cosh \alpha)^{n - 2k - 1},$$

and

$$c_{\partial\Lambda} \leq (\cosh \alpha)^d (\cosh^d \alpha - 1) |\partial\Lambda|,$$

is a boundary term.

It is now evident that (3.22) holds with $\alpha_0 = (\cosh \alpha)^{2d}$ and

$$\begin{aligned}
\tilde{H}_\Lambda &= - \sum_{x \in \Lambda} S_x^1 - 2d\beta_0(2d) \sum_{\langle x, y \rangle \in \Lambda} S_x^3 S_y^3 \\
&\quad + \alpha_1(2d) \sum_{x \in \Lambda} \sum_{Y_2 \subset \Lambda, |y_j - x| = 1} S_{y_1}^3 S_{y_2}^3 + \\
&\quad + \sum_{k=2}^d \left[\alpha_k(2d) \sum_{x \in \Lambda} \sum_{Y_{2k} \subset \Lambda, |y_j - x| = 1} S_{[Y_{2k}]}^3 \right. \\
&\quad \left. - \beta_{k-1}(2d) \sum_{x \in \Lambda} \sum_{Y_{2k-1} \subset \Lambda, |y_j - x| = 1} S_x^3 S_{[Y_{2k-1}]}^3 \right]. \quad (3.26)
\end{aligned}$$

In the case $d = 1$ one has in particular, for $\Lambda = [-L, L]$,

$$\tilde{H}_\Lambda = - \sum_{x \in \Lambda} S_x^1 - (\sinh 2\alpha) \sum_{\langle x, y \rangle \in \Lambda} S_x^3 S_y^3 + (\sinh \alpha)^2 \sum_{x, y \in \Lambda, |x - y| = 2} S_x^3 S_y^3, \quad (3.27)$$

with boundary term

$$H_{\partial\Lambda} = \sinh \alpha (1 - \cosh \alpha) (S_{-L}^3 S_{-L+1}^3 + S_{L-1}^3 S_L^3) + 2 \cosh \alpha (1 - \cosh \alpha).$$

\tilde{H}_Λ is essentially the Hamiltonian introduced by Matsui in [18]. Notice that U_0 is of the form (2.21) so that in dimensions $d \geq 2$ there is long-range order of two different kinds by Theorem 2.3

3.2 Example 2

Put $J_x = 0$, $J_{x,y} = -1$, $|x - y| = 1$; $J_{x,y} = 0$, $|x - y| > 1$ and let U_0 be given by (3.23).

We first consider the one-dimensional case $d = 1$.

Since $J_A = 0$ unless A is a pair of nearest neighbour sites, we only need to compute $W_{\{x,x+1\}}$. It is given by the formula ($\Lambda = [-L, L]$)

$$W_{x,x+1}(s_\Lambda) = 2((1 - \delta_{-L,x})s_{x-1}s_x + (1 - \delta_{L,x})s_{x+1}s_{x+2}). \quad (3.28)$$

If $-L + 1 \leq x \leq L - 2$ then an application of (3.25) yields

$$\begin{aligned} e^{-\frac{\alpha}{2}W_{x,x+1}(S_\Lambda^3)} &= (\cosh \alpha - S_{x-1}^3 S_x^3 \sinh \alpha)(\cosh \alpha - S_{x+1}^3 S_{x+2}^3 \sinh \alpha) \\ &= -(\cosh \alpha)(\sinh \alpha)(S_{x-1}^3 S_x^3 + S_{x+1}^3 S_{x+2}^3) \\ &\quad + (\sinh \alpha)^2 S_{x-1}^3 S_x^3 S_{x+1}^3 S_{x+2}^3 + (\cosh \alpha)^2. \end{aligned}$$

We also have,

$$e^{-\frac{\alpha}{2}W_{-L,-L+1}(S_\Lambda^3)} = \cosh \alpha - S_{-L+1}^3 S_{-L+2}^3 \sinh \alpha$$

and

$$e^{-\frac{\alpha}{2}W_{L-1,L}(S_\Lambda^3)} = \cosh \alpha - S_{L-2}^3 S_{L-1}^3 \sinh \alpha$$

We thus obtain the following expression for the Hamiltonian:

$$\begin{aligned} H_\Lambda &= - \sum_{-L \leq x \leq L-1} S_x^1 S_{x+1}^1 - (\cosh \alpha)(\sinh \alpha) \sum_{-L+1 \leq x \leq L-2} (S_{x-1}^3 S_x^3 + S_{x+1}^3 S_{x+2}^3) \\ &\quad + (\sinh \alpha)^2 \sum_{-L+1 \leq x \leq L-2} S_{[(x-1, \dots, x+2)]}^3 - \sinh \alpha (S_{-L+1}^3 S_{-L+2}^3 + S_{L-2}^3 S_{L-1}^3) \\ &\quad + (2L - 2)(\cosh \alpha)^2 + 2 \cosh \alpha. \end{aligned} \quad (3.29)$$

This is obviously of the form (3.22) with $\alpha_0 = (\cosh \alpha)^2$, and bulk Hamiltonian given by

$$\begin{aligned} \tilde{H}_\Lambda &= - \sum_{-L \leq x \leq L-1} [S_x^1 S_{x+1}^1 + (\sinh 2\alpha) S_x^3 S_{x+1}^3] \\ &\quad + (\sinh \alpha)^2 \sum_{-L+1 \leq x \leq L-2} S_{[(x-1, \dots, x+2)]}^3. \end{aligned} \quad (3.30)$$

Next we analyse the case of arbitrary d . We have, for a bond $\langle x, y \rangle \in \Lambda$,

$$W_{x,y}(s_\Lambda) = 2 \sum_{b \in B_{x,y}^o} s_b, \quad s_b = s_z s_{z'}, \text{ if } \langle z, z' \rangle = b, \quad (3.31)$$

and hence

$$e^{-\frac{\alpha}{2} W_{x,y}(S_\Lambda^3)} = \prod_{\langle z, z' \rangle \in B_{x,y}^o} e^{-\alpha S_z^3 S_{z'}^3}. \quad (3.32)$$

where $B_{x,y}^o$ is the set of bonds stemming from the points x, y excluding the bond $\langle x, y \rangle$ itself. Another application of (3.25) yields

$$\begin{aligned} H_\Lambda &= - \sum_{\langle x, y \rangle \in \Lambda} S_x^1 S_y^1 + \\ &+ \sum_{\langle x, y \rangle \in \Lambda} \left\{ \left(\sum_{Z \subset N_x \setminus \{y\}} \gamma_x(|Z|) S_{[Z]_x}^3 \right) \left(\sum_{Z' \subset N_y \setminus \{x\}} \gamma_y(|Z'|) S_{[Z']_y}^3 \right) \right\} \end{aligned} \quad (3.33)$$

where $N_x = \{z \in \Lambda \mid |x - z| = 1\}$ and $N_y = \{z \in \Lambda \mid |y - z| = 1\}$, $[Z]_x = Z$ if $|Z|$ is even and $[Z]_x = Z \cup \{x\}$ if $|Z|$ is odd, and similarly for $[Z']_y$ and

$$\gamma_x(n) = (\cosh \alpha)^{n_x - n - 1} (\sinh \alpha)^n \quad (3.34)$$

and similarly for γ_y . This is clearly of the form (3.22) with $\alpha_0 = d(\cosh \alpha)^{2(2d-1)}$, and bulk Hamiltonian given by

$$\tilde{H}_\Lambda = - \sum_{\langle x, y \rangle \in \Lambda} [S_x^1 S_y^1 + \gamma S_x^3 S_y^3] + \sum_{\langle x, y \rangle \in \Lambda} \sum_{j=2}^{2(2d-1)} (-1)^j \gamma_j \sum_{\{b_1, \dots, b_j\} \subset B_{x,y}^o} S_{[\cup b_j]}^3, \quad (3.35)$$

where

$$\gamma = 2(2d-1)(\cosh \alpha)^{4d-3} (\sinh \alpha) \quad (3.36)$$

and

$$\gamma_j = (\cosh \alpha)^{4d-2-j} (\sinh \alpha)^j \quad (3.37)$$

and $\cup b_j$ includes x or y if they occur an odd number of times.

References

- [1] N. N. Bogoliubov (Jr.), J. G. Brankov, V. A. Zagrebnov, A. M. Kurbatov & N. C. Tonchev, *The Approximating Hamiltonian Method in Statistical Physics*. (Publ. Bulgarian Acad. Sciences, Sophia, 1981.)
- [2] N. N. Bogoliubov (Jr.) *On model dynamical systems in statistical mechanics*. *Physica* **32**, 933–944 (1966).
- [3] N. N. Bogoliubov (Jr.), J. G. Brankov, V. A. Zagrebnov, A. M. Kurbatov & N. C. Tonchev, *Some classes of exactly soluble models of problems in quantum statistical mechanics. The method of the approximating Hamiltonian*. *Russian Math. Surveys* **39** No. 6, 1–50 (1984).
- [4] W. Cegła, J. T. Lewis & G. A. Raggio, *The free energy of quantum spin systems and large deviations*. *Commun. Math. Phys.* **118**, 337–354 (1988).
- [5] N. Duffield & J. V. Pulé, *A New Method for the Thermodynamics of the BCS Model*. *Commun. Math. Phys.* **118**, 475–494 (1988).
- [6] N. G. Duffield & J. V. Pulé, *Thermodynamics and Phase Transitions in the Overhauser Model*. *J. Stat. Phys.* **54**, 449–475 (1989).
- [7] T. C. Dorlas, *Probabilistic derivation of a noncommutative version of Varadhan’s theorem*. Preprint DIAS-02-05.
- [8] M. Fannes, H. Spohn & A. Verbeure, *Equilibrium states for mean field models*. *J. Math. Phys.* **21**, 355–358 (1980).
- [9] D. Petz, G. A. Raggio & A. Verbeure, *Asymptotics of Varadhan type and the Gibbs variational principle*. *Commun. Math. Phys.* **121**, 271–282 (1989).
- [10] G. A. Raggio & R. Werner, *Quantum statistical mechanics of general mean-field systems*. *Helv. Phys. Acta* **62**, 980–1003 (1989).
- [11] V. A. Zagrebnov & J.-B. Bru, *The Bogoliubov model of weakly imperfect Bose gas*. *Phys. Rep.* **350**, 291–442 (2001).
- [12] J.-B. Bru & T. C. Dorlas, *Exact solution of the infinite-range-hopping Bose-Hubbard model*. *J. Stat. Phys.* **113**, 177–196 (2003).
- [13] F.R. Gantmacher, *Applications of the Theory of Matrices*, Interscience Publ. New York, 1959.

- [14] D. Serre, *Matrices. Theory and Applications*. Graduate Texts in Mathematics Vol. 216, Springer Verlag, New York etc., 2002.
- [15] J. R. Kirkwood, L. Thomas, *Expansions and phase transitions for the ground state of quantum lattice systems*. Commun. Math. Phys. **88**, 569–580 (1982).
- [16] N. Datta and T. Kennedy, *Expansions for one quasiparticle states in spin-1/2 systems*. J. Stat. Phys. **108**, 373–399 (2002).
- [17] D. A. Yarotsky, *Perturbations of ground states in weakly interacting quantum spin systems*. To appear in J. Math. Phys.
- [18] T. Matsui, *A link between quantum and classical Potts models*, J. Stat. Phys. **59**, Nos. 3/4, 781–798, (1990).
- [19] N. Macris & C.-A. Pignatelli, *Coexistence of long-range order for two observables at finite temperatures*, J. Stat. Phys. **105**, 909–935 (2001).
- [20] N. Macris, *Charge density wave and quantum fluctuations in a molecular crystal*, cond-mat/9906008.
- [21] Guang-Shan Tian, *A sufficient condition for two long-range orders coexisting in a lattice many-body system*, J. Phys. A **30**, 841–848 (1997).
- [22] R. A. Minlos, *Invariant subspaces of the stochastic Ising high temperature dynamics*, Markov Processes & Related Fields **2**, 263–284, (1996).
- [23] W. Skrypnik, *Long-range order in nonequilibrium systems of interacting Brownian linear oscillators*. J. Stat. Phys. **111**, 291–321 (2003).