

INTEGRABLE SOLUTIONS OF THE HIERARCHY
OF THE BBGKY-TYPE FOR BROWNIAN PARTICLES
IN THE MEAN-FIELD LIMIT

W.I. Skrypnik¹

Dublin Institute for Advanced Studies
10 Burlington Road, Dublin 4, Ireland

Introduction

Let us consider the gradient hierarchy of the BBGKY-type for correlation functions $\rho_t(X_m)$, $X_m = (x_1, \dots, x_m) \in \mathbb{R}^{dm}$, $x_j \in \mathbb{R}^d$, of a nonequilibrium system of diffusing particles, interacting via a pair, integrable smooth potential $\varepsilon \Phi(x)$. Let us assume also that the following asymptotic relation holds

$$\rho_t(X_m) = \varepsilon^{-m} \rho_t^\varepsilon(X_m), \quad \varepsilon \searrow 0$$

where the functions $\rho_t^\varepsilon(X_m)$ have a limit when $\varepsilon \searrow 0$. Then

¹On leave of absence from the Institute of mathematics of the Ukrainian Academy of Sciences, 252004, Kiev 4, Repin Street 3, USSR

the hierarchy is written as follows

(1)

$$\begin{aligned} \frac{\partial}{\partial t} \rho_t^\varepsilon(\mathbf{X}_m) = & \sum_{j=1}^m \frac{\partial}{\partial \mathbf{x}_j} \left\{ \beta^{-1} \frac{\partial}{\partial \mathbf{x}_j} \rho_t^\varepsilon(\mathbf{X}_m) + \varepsilon \rho_t^\varepsilon(\mathbf{X}_m) \frac{\partial}{\partial \mathbf{x}_j} U(\mathbf{X}_m) \right. \\ & \left. + \int_{\mathbb{R}^d} \left(\frac{\partial \phi}{\partial \mathbf{x}_j} \right) (\mathbf{x}_j - \mathbf{x}_{m+1}) \rho_t^\varepsilon(\mathbf{X}_{m+1}) d\mathbf{x}_{m+1} \right\} \end{aligned}$$

where β is the inverse temperature and $\varepsilon U(\mathbf{X}_m)$ is the potential energy

$$\begin{aligned} U(\mathbf{X}_m) = & \sum_{1 \leq i < j \leq m} \phi(\mathbf{x}_i - \mathbf{x}_j) , \\ \frac{\partial}{\partial \mathbf{x}_j} \left(f \frac{\partial h}{\partial \mathbf{x}_j} \right) = & \sum_{\nu=1}^d \left\{ \frac{\partial f}{\partial \mathbf{x}_j^\nu} \frac{\partial h}{\partial \mathbf{x}_j^\nu} + f \frac{\partial^2 h}{\partial (\mathbf{x}_j^\nu)^2} \right\} . \end{aligned}$$

In the mean-field limit ($\varepsilon \rightarrow 0$) the considered hierarchy is transformed into a hierarchy of the Vlasov type

(2)

$$\begin{aligned} \frac{\partial}{\partial t} \rho_t^0(\mathbf{X}_m) = & \sum_{j=1}^m \frac{\partial}{\partial \mathbf{x}_j} \left\{ \beta^{-1} \frac{\partial}{\partial \mathbf{x}_j} \rho_t^0(\mathbf{X}_m) + \right. \\ & \left. + \int_{\mathbb{R}^d} \left(\frac{\partial \phi}{\partial \mathbf{x}_j} \right) (\mathbf{x}_j - \mathbf{x}_{m+1}) \rho_t^0(\mathbf{X}_{m+1}) d\mathbf{x}_{m+1} \right\} \end{aligned}$$

In this paper we propose a justification of the mean-field limit in a class of integrable correlation

functions satisfying eq.(1) in a weak sense. We introduce a semigroup π_ε^t in a Banach space L_ξ^1 of sequences of symmetric itegrable functions and prove that the sequeense

$$\rho_\varepsilon^t(X_m) = (\pi_\varepsilon^t \rho^\varepsilon)(X_m), \quad \rho^\varepsilon \in L_\xi^1$$

satisfies eq.(1) in a weak sense. We show that, if the sequence $\exp(-\frac{t}{\varepsilon}U(X_m)) \rho_\varepsilon^t(X_m)$, $m \geq 1$, belongs to L_ξ^1 , then $\rho_\varepsilon^t(X_m)$ converges weakly to

$$\rho_t^0(X_m) = (\pi_0^t \rho^0)(X_m), \quad m \geq 1,$$

where π_0^t is defined as a map

$$L_\xi^1 \rightarrow L_{\xi(t)}^1, \quad \text{if } \xi(t) = \sqrt{2} \exp(-\hat{\nu}(t)) \xi < 1.$$

We also prove that the sequence ρ_t^0 is a weak solution of eq.(2).

The norm in the Banach space L_ξ^1 is defined as follows

$$\|\Psi\|_{L_\xi^1} = \max_{n \geq 1} \xi^{-n} \|\Psi_n\|_{L^1(\mathbb{R}^{dn})}.$$

The mean-field limit in a mechanical and a special random mechanical systems was studied earlier, respectively in [2,3] (see also [4,5]). The mean-field limit for eq.(1) in a class of bounded correlation functions is investigated in [6]. It is found in [7] that there is a space in which a solution of eq.(2) exists and is unique.

I. Main theorem.

Let $P_\varepsilon^t(X_n; Y_n)$ be a fundamental solution of the n-particle Smoluchowski equation (eq.(1) without the integral term in the r.s.). The operators $P_{\varepsilon, n}^t$, $t \geq 0$,

$$(P_{\varepsilon, n}^t \psi)(X_n) = \int_{\mathbb{R}^{dn}} P_\varepsilon^t(X_n; Y_n) \psi(Y_n) dY_n$$

defines a contraction strongly continuous semigroup in $L^1(\mathbb{R}^{dn})$. Let P_ε^t be a diagonal operator in \mathbb{L}_ξ^1 , given by

$$(P_\varepsilon^t \psi)(X_n) = (P_{\varepsilon, n}^t \psi)(X_n).$$

It is evident that P_ε^t is a contraction strongly continuous semigroup in \mathbb{L}_ξ^1 . Let us define an operator $\int d_x$ which is bounded in \mathbb{L}_ξ^1

$$(\int d_x \psi)(X_n) = \int_{\mathbb{R}^d} \psi(x, X_n) dx$$

Then π_ε^t

$$(1.1) \quad \pi_\varepsilon^t = \exp\{ \varepsilon^{-1} \int d_x \} P_\varepsilon^t \exp\{ -\varepsilon^{-1} \int d_x \}$$

is a strongly continuous semigroup in \mathbb{L}_ξ^1 . Its structure coincide with a structure of an evolution operator of the BBGKY-hierarchy [8].

Lemma 1.1

The sequence $\rho_t^\varepsilon = \pi_\varepsilon^t \rho^\varepsilon$, $\rho^\varepsilon \in \mathbb{L}_\xi^1$, is a weak solution of eq.(1), that is

$$\begin{aligned}
(1.2) \quad & \frac{\partial}{\partial t} \int_{\mathbb{R}^d} \rho_t^\varepsilon(X_m) h(X_m) dX_m = \\
& = \sum_{j=1}^m \int_{\mathbb{R}^d} \left\{ \beta^{-1} \rho_t^\varepsilon(X_m) \frac{\partial^2}{\partial x_j^2} h(X_m) - \right. \\
& \quad \left. - \left(\frac{\partial}{\partial x_j} h(X_m) \right) \left[\varepsilon \rho_t^\varepsilon(X_m) \frac{\partial}{\partial x_j} U(X_m) + \right. \right. \\
& \quad \left. \left. + \int_{\mathbb{R}^d} \left(\frac{\partial \phi}{\partial x_j} \right) (x_j - x_{m+1}) \rho_t^\varepsilon(X_{m+1}) dx_{m+1} \right] \right\} dX_m .
\end{aligned}$$

Lemma 1.2

Let $P_x(\tilde{dx})$ be the Wiener measure on $\Omega_d = (\mathbb{R}^d)^{\mathbb{R}^+}$ and $P_{X_m}(\tilde{dX}_m) = \prod_{j=1}^m P_{x_j}(\tilde{dx})$. Let $*$ be an operation of multiplication, defined on sequences

$$\{ \psi(X_n, \tilde{X}_n), X_n \in \mathbb{R}^{dn}, \tilde{X}_n \in \Omega_d^n \}_{n \geq 0}, \quad \psi(\emptyset, \emptyset) = 1$$

by

$$(\psi_1 * \psi_2)(X_n, \tilde{X}_n) = \sum_{\{s\} \in (1, \dots, n)} \psi_1(X_{\{s\}}, \tilde{X}_{\{s\}}) \psi_2(X_{\{n \setminus s\}}, \tilde{X}_{\{n \setminus s\}})$$

$$\{n \setminus s\} = (1, \dots, n) \setminus \{s\} .$$

Let $(\psi)^{-1}$ be the inverse element to ψ with respect to $*$ and

$$(D_{(X_m; \tilde{X}_m)} \psi)(X'_n; \tilde{X}'_n) = \psi(X_m, X'_n; \tilde{X}_m, \tilde{X}'_n)$$

There exists a measurable function $\hat{U}_t^\varepsilon(X_n, \tilde{X}_n)$ such that, if

$$\begin{aligned} & \Pi_\varepsilon(X_m, \tilde{X}_m | X_n', \tilde{X}_n') = \\ & = \varepsilon^{-n} (\exp\{ -\beta \hat{U}_t^\varepsilon \})^{-1} * D_{(X_m; \tilde{X}_m)} \exp\{ -\beta \hat{U}_t^\varepsilon \} (X_n'; \tilde{X}_n') , \end{aligned}$$

where the n-th component of the sequence $\exp\{ -\beta \hat{U}_t^\varepsilon \}$ equals $\exp\{ -\beta \hat{U}_t^\varepsilon(X_n; \tilde{X}_n) \}$, then the cluster expansion for π_ε^t holds

(1.3)

$$\begin{aligned} (\pi_\varepsilon^t \rho_t^\varepsilon)(X_m) &= \sum_{n \geq 0} \frac{1}{n!} \int_{\mathbb{R}^{dn}} dX_n' \int_{\Omega_d^{n+m}} P_{X_m, X_n'}(d\tilde{X}_m d\tilde{X}_n') \times \\ & \times \Pi_\varepsilon(X_m, \tilde{X}_m | X_n', \tilde{X}_n') \rho(\tilde{X}_m(t\beta^{-1}), \tilde{X}_n'(t\beta^{-1})) . \end{aligned}$$

Lemma 1.3

If the potential ϕ is a positive-definite function from $C^3(\mathbb{R}^d)$ and

$$|\phi(x)| \leq \phi^0, \quad |\nabla\phi(x)| \leq \phi^0, \quad |\Delta\phi(x)| \leq \phi^0, \quad \Delta = \nabla^2, \nabla = \frac{\partial}{\partial x} .$$

then the following uniform in ε bound holds

(1.4)

$$\text{ess sup}_{\text{all}(x, x', \tilde{x}, \tilde{x}')} \exp\{ -\frac{1}{2} \beta \hat{U}_t^\varepsilon(X_m, X_n'; \tilde{X}_m, \tilde{X}_n') \} |\Pi_\varepsilon(X_m, \tilde{X}_m | X_n', \tilde{X}_n')| \leq$$

$$\leq n! (\sqrt{2} \exp\{ \hat{\nu}(t) \})^{m+n} ,$$

$$\hat{\nu}(t) = (\beta \phi^0 x(t))^2 + \frac{1}{2} \phi(0) + \frac{1}{2} t (-\Delta\phi)(0) ,$$

and the functions $\Pi_\varepsilon(X_m, \tilde{X}_m | X_n', \tilde{X}_n')$ converge a.e. to functions

$\Pi_0(X_m, \tilde{X}_m | X'_n, \tilde{X}'_n)$, satisfying (1.4) for $\varepsilon = 0$.

Corollary .The operator π_0^t , defined by (1.3) for $\varepsilon = 0$ maps L^1_ξ into $L^1_{\xi(t)}$, if $\xi(t) < 1$.

Theorem 1.1

Let the conditions of the Lemma 1.3 be satisfied. If the following conditions are also satisfied

$$\exp \left\{ \frac{1}{2} \beta U \right\} \rho^\varepsilon \in L^1_\xi ,$$

$$\| \exp \left\{ \frac{1}{2} \beta U \right\} \rho^\varepsilon - \rho^0 \|_{L^1_\xi} \leq o(\varepsilon) ,$$

where $(\exp \left\{ \frac{1}{2} \beta U \right\})(X_n) = \exp \left\{ \frac{1}{2} \beta U(X_n) \right\}$,

then the functions $(\pi_\varepsilon^t \rho^\varepsilon)(X_m)$ converge weakly to the functions $(\pi_0^t \rho^0)(X_m)$, sequence of which satisfies eq.(2) in a weak sense

$$\begin{aligned} (1.5) \quad & \frac{\partial}{\partial t} \int_{\mathbb{R}^{dm}} \rho_t^0(X_m) h(x_m) dX_m = \\ & = \sum_{j=1}^m \int_{\mathbb{R}^{dm}} \left\{ \beta^{-1} \rho_t^0(X_m) \frac{\partial^2}{\partial (x_j)^2} h(X_m) - \right. \\ & \left. - \left(\frac{\partial}{\partial x_j} h(X_m) \right) \int_{\mathbb{R}^d} \left(\frac{\partial \phi}{\partial x_j} \right) (x_j - x_{m+1}) \rho_t(X_{m+1}) dx_{m+1} \right\} dX_m . \end{aligned}$$

2. Cluster expansion .

Let us prove the equality (1.3). We start from resumming in (1.1)

(2.1)

$$\pi_\varepsilon^t = \sum_{n \geq 0} \frac{\varepsilon^{-n}}{n!} \int_{\mathbb{R}^n} dX'_n \sum_{s=0}^n \frac{n!}{s!(n-s)!} (-1)^{n-s} D_{X'_s} P^t D_{X'_{\{n \setminus s\}}}$$

where $(D_{X'_m} \psi)(X'_n) = \psi(X'_m, X'_n)$.

Let $*$ be the operation of multiplication defined on sequences of kernels $\{K(X_n; Y_n)\}_{n \geq 0}$, $K(\emptyset, \emptyset) = 1$ (see Lemma 1.2) . $e_0 = (1, 0, 0, \dots)$ is the unit with the respect to $*$. Define

$$(K)^{-1} = \sum_{n \geq 0} (-1)^n (K - e_0) * \dots * (K - e_0) , \quad (K)^{-1} * K = e_0 ,$$

$$\langle K \rangle_\psi = \sum_{n \geq 0} \frac{1}{n!} \int_{\mathbb{R}^n} dX_n \int_{\mathbb{R}^n} dY_n K(X_n; Y_n) \psi(Y_n)$$

Then (1.2) is rewritten in the following fashion

(2.2)

$$(\pi_\varepsilon^t \rho^\varepsilon)(X_m) = \int_{\mathbb{R}^{dm}} \langle e_{-1} * D_{(X_m; Y_m)} P_\varepsilon^t \rangle_{\varepsilon^{-1}}^{-1} D_{Y_m} \rho^\varepsilon dY_m$$

where $(e_{\pm 1})(X_n; Y_n) = (\pm 1)^n \delta(X_n - Y_n)$, $(\hat{\varepsilon}^{-1})(\rho^\varepsilon)(X_n) = \varepsilon^{-n} \rho^\varepsilon(X_n)$

As a result we derive the cluster expansion

(2.3)

$$(\pi_\varepsilon^t \rho^\varepsilon)(X_m) = \int_{\mathbb{R}^{dm}} \langle (P_\varepsilon^t)^{-1} * D_{(X_m; Y_m)} P_\varepsilon^t \rangle_{\varepsilon^{-1}}^{-1} D_{Y_m} \rho^\varepsilon dY_m$$

(2.3) follows from the law of conservation of probability for the Smoluchowski equation and the equality $e_{-1} = (e_{+1})^{-1}$

in the following way

$$\begin{aligned}
 \langle (P_\epsilon)^{-1} * D_{(X_m; Y_m)} P_\epsilon \rangle_\psi &= \sum_{n \geq 0} \langle (P_\epsilon^t - e_0) * \dots * (P_\epsilon^t - e_0) * D_{(X_m; Y_m)} P_\epsilon^t \rangle_\psi = \\
 &= \sum_{n \geq 0} \langle (e_1 - e_0) * \dots * (e_1 - e_0) * D_{(X_m; Y_m)} P_\epsilon^t \rangle_\psi = \\
 &= \langle (e_{+1})^{-1} * D_{(X_m; Y_m)} P_\epsilon^t \rangle_\psi = \langle e_{-1} * D_{(X_m; Y_m)} P_\epsilon^t \rangle_\psi .
 \end{aligned}$$

It is well known that, if a function $\mu_t^\epsilon(X_n)$ satisfies the Smoluchowski equation, the function

$$\exp \left\{ -\frac{1}{2} \epsilon \beta U(X_n) \right\} \mu_t^\epsilon(X_n)$$

satisfies the heat equation with the potential $V_\epsilon(X_n)$

$$V_\epsilon(X_n) = \frac{1}{2} \beta \epsilon \sum_{j=1}^n \left\{ \frac{\partial^2}{\partial x_j^2} U(X_n) - \frac{1}{2} \beta \epsilon \left(\frac{\partial}{\partial x_j} U(X_n) \right)^2 \right\} .$$

Applying the Feynman-Kac formula we derive the following representation for the kernel P_ϵ^t

$$\begin{aligned}
 (2.4) \quad P_\epsilon^t(X_n; Y_n) &= \exp \left\{ -\frac{1}{2} \beta \epsilon (U(X_n) - U(Y_n)) \right\} \times \\
 &\times \int_{\Omega_d^n} P_{X_n} (d\tilde{X}_n) \exp \left\{ \int_0^{t\beta^{-1}} V_\epsilon(\tilde{X}_n(\tau)) d\tau \right\} \delta(\tilde{X}_n(t\beta^{-1}) - Y_n) .
 \end{aligned}$$

(2.2) and (2.4) yield (1.3) with

$$\tilde{U}_t^\epsilon(X_n; \tilde{X}_n) = \frac{1}{2} \epsilon (U(X_n) - U(\tilde{X}_n(t\beta^{-1}))) + \beta^{-1} \int_0^{t\beta^{-1}} V(\tilde{X}_n(\tau)) d\tau .$$

3. Proof of the main estimate .

In order to prove (1.4) we shall make use of the following identity

$$(3.1) \quad \exp \left\{ -\frac{1}{4} \beta^2 \varepsilon^2 \sum_{j=1}^n \int_0^{t\beta^{-1}} \left(\frac{\partial U}{\partial \mathbf{x}_j} \right)^2 (\tilde{\mathbf{X}}_n(\tau)) d\tau \right\} = \\ = \int_{\Omega_d^n} P(\mathbf{X}_n^*) \exp \left\{ -\frac{1}{2} \beta \varepsilon i \sum_{j=1}^n \int_0^{t\beta^{-1}} \left(\frac{\partial U}{\partial \mathbf{x}_j} \right) (\tilde{\mathbf{X}}_n(\tau)) , d\mathbf{x}_j^*(\tau) \right\} .$$

where $P(\mathbf{X}_n^*) = \prod_{j=1}^n P(d\mathbf{x}_j^*)$, $P(d\mathbf{x}^*) = P_0(d\mathbf{x}^*)$, $P_0(d\mathbf{x}^*)$ is the Wiener measure , $\int(\dots, d\mathbf{x}^*(\tau))$ is the stochastic integral, (\dots, \dots) is the scalar product of \mathbb{E}^d (we omit \sim over \mathbf{x}_j in derivatives).

Substituting (3.1) into (1.3) we obtain

(3.2)

$$\Pi_\varepsilon(\mathbf{X}_m, \tilde{\mathbf{X}}_m | \mathbf{X}'_n, \tilde{\mathbf{X}}'_n) = \int_{\Omega_d^{m+n}} P(d\mathbf{X}_m^*) P(d\mathbf{X}'_n^*) \Pi_\varepsilon(\hat{\mathbf{X}}_m | \hat{\mathbf{X}}'_n) ,$$

where $\hat{\mathbf{X}}_m = (\hat{\mathbf{x}}_1, \dots, \hat{\mathbf{x}}_m)$, $\hat{\mathbf{x}}_j = (\mathbf{x}_j, \tilde{\mathbf{x}}_j, \mathbf{x}_j^*) \in \mathbb{R}^d \times \Omega_d^2$,

$$\Pi_\varepsilon(\hat{\mathbf{X}}_m | \hat{\mathbf{X}}'_n) = \varepsilon^{-n} ((\exp \{-\varepsilon\beta \hat{U}_t\}) * D_{\hat{\mathbf{X}}_m} \exp \{-\varepsilon\beta \hat{U}_t\}) (\hat{\mathbf{X}}'_n)$$

$$\hat{U}_t(\hat{\mathbf{X}}_n) = \sum_{1 \leq k < j \leq n} \hat{\phi}_t(\hat{\mathbf{x}}_k | \hat{\mathbf{x}}_j) ,$$

$$\hat{\phi}_t(\hat{\mathbf{x}}_k | \hat{\mathbf{x}}_j) = \tilde{\phi}_t(\mathbf{x}_k - \mathbf{x}_j, \tilde{\mathbf{x}}_k - \tilde{\mathbf{x}}_j) + \phi_t^*(\tilde{\mathbf{x}}_k - \tilde{\mathbf{x}}_j | \mathbf{x}_k^*, \mathbf{x}_j^*) ,$$

$$\tilde{\phi}(\mathbf{x}, \tilde{\mathbf{x}}) = \frac{1}{2} [\phi(\mathbf{x}) - \phi(\tilde{\mathbf{x}}(t\beta^{-1}))] + \int_0^{t\beta^{-1}} (-\Delta\phi)(\tilde{\mathbf{x}}(\tau)) d\tau ,$$

$$\phi_t^*(\tilde{x}|x_k^*, x_j^*) = \frac{1}{2} i \{ \varphi_t(\tilde{x}|x_k^*) - \varphi_t(\tilde{x}|x_j^*) \} ,$$

$$\varphi_t(\tilde{x}|x^*) = \frac{1}{2} \int_0^{t\beta^{-1}} ((\nabla\phi)(\tilde{x}(\tau)), dx^*(\tau)) ,$$

The sequence $\Pi_\varepsilon(\hat{X}_m|\hat{X}_n')$ $m \geq 1, n \geq 1$ satisfies the standard relation [9]

$$(3.3) \quad \Pi_\varepsilon(\hat{X}_m|\hat{X}_n') = \exp \left\{ -\varepsilon\beta \sum_{\substack{l=1, \\ l \neq j}}^m \hat{\phi}_t(\hat{x}_l|\hat{x}_j) \right\} \times \\ \times \sum_{\{s\} \in (1, \dots, n)} K_\varepsilon(\hat{x}_j|\hat{X}'_{\{s\}}) \Pi_\varepsilon(\hat{X}_{m(j)}, \hat{X}'_{\{s\}}|\hat{X}'_{\{n \setminus s\}}) ,$$

$$\Pi_\varepsilon(\hat{X}_m|\emptyset) = \exp \left\{ -\varepsilon\beta \hat{U}_t(\hat{X}_m) \right\} , \quad \Pi_\varepsilon(\emptyset|\hat{X}_n) = 0 , \quad m(j) = (1, \dots, m) \setminus j$$

$$K_\varepsilon(\hat{x}|\hat{X}_n) = \prod_{j=1}^n \varepsilon^{-1} (\exp \{ -\varepsilon\beta \hat{\phi}_t(\hat{x}|\hat{x}_j) \} - 1) .$$

(3.3) has the limit for $\varepsilon \rightarrow 0$.

$$(3.4) \quad \Pi_0(\hat{X}_m|\hat{X}_n') =$$

$$= \sum_{\{s\} \in (1, \dots, n)} \prod_{l \in \{s\}} (-\beta) \hat{\phi}_t(\hat{x}_j|\hat{x}_l) \Pi_0(\hat{X}_{m(j)}, \hat{X}'_{\{s\}}|\hat{X}'_{\{n \setminus s\}}) .$$

Proposition 1.3 (The main bound)

If the conditions of Lemma 1.3 are satisfied then the following uniform in all variables, except t , bound

(3.5)

$$\left\{ \int_{\Omega_d^m} P(dX_m^*) \left[\int_{\Omega_d^n} P(dX_n'^*) |\Pi_\varepsilon(\hat{X}_m|\hat{X}_n')| \right]^2 \right\}^{\frac{1}{2}} \leq n! (\sqrt{2} \exp\{\hat{\nu}(t)\})^{m+n}$$

Proof.

To derive (3.5) we have to symmetrize (3.3), taking into consideration that the potential

$$\phi_{0t}(x, x) = \tilde{\phi}_t(x, \tilde{x}) + \frac{1}{2}\phi(\tilde{x}(t\beta^{-1}))$$

is stable (see also [6,9,10]). Now we prove (3.5) by induction for $\varepsilon = 0$.

Let us integrate (3.4) by $P(dX_n^{i*})$ and apply the Schwartz inequality

$$\begin{aligned} & \int_{\Omega_d^{<s>}} P(dX_{\{s\}}^{i*}) \left| \prod_{l \in \{s\}} \hat{\phi}_t(\hat{x}_j | \hat{x}_1) \right| \times \\ & \times \int_{\Omega_d^{<n-s>}} P(dX_{\{n \setminus s\}}^{i*}) \left| \Pi_0(\hat{X}_{m(j)}, \hat{X}'_{\{s\}} | \hat{X}'_{\{n \setminus s\}}) \right| \\ & \leq \left[\int_{\Omega_d^{<s>}} P(dX_{\{s\}}^{i*}) \left| \prod_{l \in \{s\}} \hat{\phi}_t(\hat{x}_j | \hat{x}_1) \right|^2 \right]^{\frac{1}{2}} \times \\ & \times \left[\int_{\Omega_d^{<s>}} P(dX_{\{s\}}^{i*}) \left[\int_{\Omega_d^{<n \setminus s>}} P(dX_{\{n \setminus s\}}^{i*}) \left| \Pi_0(\hat{X}_{m(j)}, \hat{X}'_{\{s\}} | \hat{X}'_{\{n \setminus s\}}) \right|^2 \right]^{\frac{1}{2}} \right]^{\frac{1}{2}} \end{aligned}$$

where $<s>$ is the number of elements in the sequence $\{s\}$.

Let us square the obtained inequality, split the sum over $\{s\}$ into two sums (over $\{s_1\}$ and $\{s_2\}$) and integrate by $P(dX_m^*)$ the resulting expression, utilizing the Schwartz inequality. Assume that (3.5) holds for all $<s> \leq m+n-1$. In this case

$$\begin{aligned} & \int_{\Omega_d^m} P(dX_m^*) \left[\int_{\Omega_d^n} \left| \Pi_0(\hat{X}_m | \hat{X}_n) \right| P(dX_n^{i*}) \right] \leq (n! \sum_{s \geq 0} \frac{1}{s!} K_{s,0})^2 \times \\ & \times (\sqrt{2} \exp \{ \hat{v}_0(t) \})^{2(m+n-1)} \end{aligned}$$

$$K_{s,0} = \beta^s \operatorname{ess\,sup}_{\text{all}(\mathbf{x}, \tilde{\mathbf{x}})} \int_{\Omega_d} P(d\mathbf{x}^*) \left[\int_{\Omega_d} P(d\mathbf{x}_s^*) \prod_{l=1}^s |\hat{\phi}_t(\tilde{\mathbf{x}}|\hat{\mathbf{x}}_l)|^2 \right]^{\frac{1}{2}}.$$

Making use of the Schwartz and generalized Helder inequalities we obtain the following bound for $K_{s,0}$

$$\begin{aligned} K_{s,0} &\leq \beta^s \operatorname{ess\,sup}_{\text{all}(\mathbf{x}, \tilde{\mathbf{x}})} \left[\int_{\Omega_d} P(d\mathbf{x}_s^*) \prod_{l=1}^s \int_{\Omega_d} P(d\mathbf{x}^*) |\hat{\phi}_t(\tilde{\mathbf{x}}|\hat{\mathbf{x}}_l)|^2 \right]^{\frac{1}{2}} \leq \\ &\leq \beta^s \operatorname{ess\,sup}_{\text{all}(\mathbf{x}, \tilde{\mathbf{x}})} \left[\prod_{l=1}^s \int_{\Omega_d} P(d\mathbf{x}_l^*) \int_{\Omega_d} P(d\mathbf{x}^*) |\hat{\phi}_t(\tilde{\mathbf{x}}|\hat{\mathbf{x}}_l)|^{2s} \right]^{\frac{1}{2s}}. \end{aligned}$$

It is well known that

$$\int_{\Omega} P(d\mathbf{x}^*) \left[\int_0^{t\beta^{-1}} ((\nabla\phi)(\tilde{\mathbf{x}}(\tau), d\mathbf{x}^*(\tau))) \right]^{2s} = \frac{(2s)!}{(s)!} \left[\int_0^{t\beta^{-1}} (\nabla\phi)^2(\tilde{\mathbf{x}}(\tau)) d\tau \right]^{\frac{1}{2}}$$

With the help of the inequalities

$$(a^s + b^s)^{\frac{1}{s}} \leq (a+b), \quad (a+b)^s \leq 2^s (a^s + b^s), \quad 3^{-n} n^n \leq n! \leq n^n$$

we have

$$K_{s,0} \leq (s!)^{\frac{1}{2}} \left\{ 72\beta \left(|\hat{\phi}_t|_0 + 2|\nabla\phi|_0 t\beta^{-1} \right) \right\}^s \leq (s!)^{\frac{1}{2}} (\phi^0 \beta x_0(t))^s$$

where

$$|\phi|_0 = \operatorname{ess\,sup}_{\mathbf{x} \in X} |\phi(\mathbf{x})|, \quad x_0(t) = 72(1 + \frac{1}{2} 5t\beta^{-1}), \quad \hat{\nu}_0 = (x_0(t)\beta\phi^0)^2.$$

Now it is not difficult to prove the proposition for $\varepsilon > 0$, using the symmetrized (3.3)[6] and the above arguments. As a result

$$x(t) = \exp \{ \beta\phi^0 (1 + \frac{1}{2} t\beta^{-1}) \} x_0(t).$$

Now we return to the Theorem 1.1. At first we consider the simplest case

$$\rho^\varepsilon = \exp \left\{ -\frac{1}{2} \varepsilon \beta U \right\} \rho^0 .$$

Since the algebraic structure of the functions $\Pi_\varepsilon(\hat{X}_m | \hat{X}'_n)$ are known, they converge to the functions $\Pi_0(\hat{X}_m | \hat{X}'_n)$ a.e. . From the Lebesgue theorem it follows that

$$\int_{\mathbb{R}^{dn}} (\pi_\varepsilon^t \rho^\varepsilon)(X_m) h(X_m) dX_m \xrightarrow{\varepsilon \rightarrow 0} \int_{\mathbb{R}^{dn}} (\pi_0^t \rho^0)(X_m) h(X_m) dX_m .$$

It means that the r.s. of (1.2) converges to the r.s. of (1.5). Since the r.s. of (1.5) is a continuous function of t (the series in (1.3) converges uniformly on a finite time interval) the l.s. of (1.2) converges to the left side of (1.5). It is clear that in a general case

$$\rho^\varepsilon = \exp \left\{ -\frac{1}{2} \varepsilon \beta U \right\} (\rho^0 + \rho'^\varepsilon) .$$

From Lemmas 1.1, 1.2 it follows that the remaining term $\pi_\varepsilon^t \exp \left\{ -\frac{1}{2} \varepsilon \beta U \right\} \rho'^\varepsilon$ converges to zero in L^1_ξ if

$$\xi(t) = \sqrt{2} \exp \left\{ \hat{\nu}(t) \right\} \xi < 1 .$$

The theorem is proved .

4. Proof of Lemma 1.1

Let us put

$$\rho_N^\varepsilon = \exp \left\{ \varepsilon \int d_x \right\} \mu_N^\varepsilon , \quad \mu_N^\varepsilon \in L^1_\xi , \quad \mu_N^\varepsilon(X_n) = 0 , \quad n > N .$$

Then

$$\begin{aligned} \rho_{t,N}^\varepsilon(X_m) &= (\pi_\varepsilon^t \rho_N^\varepsilon)(X_m) = (\exp\{\int d_x\} P_\varepsilon^t \mu_N^\varepsilon)(X_m) = \\ &= \sum_{n=0}^{N-m} \frac{1}{n!} \int_{\mathbb{R}^{dm}} dX_n' (P_\varepsilon^t \mu_N^\varepsilon)(X_m, X_n') . \end{aligned}$$

It can be shown that $\rho_{t,N}^\varepsilon$ satisfies (1) in a classical sense. Now let us differentiate it, taking into consideration that $(P_{\varepsilon,n}^t \mu_N^\varepsilon)(X_n) = \mu_t^\varepsilon(X_n)$ satisfies the Smoluchowski equation

$$\begin{aligned} \frac{\partial}{\partial t} \rho(X_m) &= \sum_{j=1}^m \frac{\partial}{\partial x_j} \left\{ \beta^{-1} \frac{\partial}{\partial x_j} \rho_{t,N}^\varepsilon(X_m) + \rho_{t,N}^\varepsilon(X_m) \frac{\partial}{\partial x_j} U(X_m) + \right. \\ &+ \varepsilon \sum_{n=0}^{N-m} \frac{n}{n!} \varepsilon^{-n} \int_{\mathbb{R}^{dn}} \left(\frac{\partial \phi}{\partial x_j} \right) (x_j - x_{m+1}) \mu_t^\varepsilon(X_m, X_n') dX_n \end{aligned}$$

The derivatives in the inner variables (X_n') disappeared since

$$\int_{\mathbb{R}} \left(\frac{\partial h}{\partial x} \right) (x) dx = 0, \quad f, \quad \frac{\partial h}{\partial x} \in L^1(\mathbb{R}).$$

and $\mu_t^\varepsilon(X_n) \in L^1 \cap C^2[10]$. As a result the last term is equal

$$\begin{aligned} \int_{\mathbb{R}^d} \left(\frac{\partial \phi}{\partial x_j} \right) (x_j - x_{m+1}) \left[\sum_{n=0}^{N-m-1} \frac{\varepsilon^{-n}}{n!} \int_{\mathbb{R}^{dn}} \mu_t^\varepsilon(X_m, X_n') dX_n' \right] dx_{m+1} = \\ = \int_{\mathbb{R}^d} \left(\frac{\partial \phi}{\partial x_j} \right) (x_j - x_{m+1}) \rho_{t,N}^\varepsilon(X_{m+1}) dx_{m+1}, \quad m \leq N. \end{aligned}$$

Hence the sequence $\{\rho_{t,N}^\varepsilon(X_m)\}_{m < N}$ satisfies (1) in a weak sense. Let $N \rightarrow \infty$. Then $\rho_{t,N}^\varepsilon$ converges to ρ_t^ε in the topology of L_ξ^1 . By the Lebesgue theorem the r.s. of (1.2) for $\rho_{t,N}^\varepsilon$ converges to the r.s. of (1.2). Since π_ε^t is a strongly continuous semigroup the same is true for the

corresponding left sides .The proof is complete .

REMARK . Our theorem does not allow the canonical correlation functions to converge in the mean-field limit since their limit satisfies the compatibility condition

$$\int_{\mathbb{R}^d} \rho_t(X_n) dx_n = \rho_t(X_{n-1}) \text{ .and } \rho_t \in \mathbb{L}_\xi^1 \text{ only if } \xi \geq 1 \text{ .}$$

But there is a possibility to improve our bounds in such a way that $\xi(t)$ goes to 0 when either t or ϕ^0 goes to 0 .

ACKNOWLEDGEMENT .The author expresses his sincere gratitude to professor John Lewis for invitation, kindness, and hospitality .He thanks also for help Margaret Matthews, Michel Vandyck and Guido Raggio .

REFERENCES

- 1.Streltsova E.A. Ukrainian Math.Journ.,1959,11, 1, 83-92.
- 2.Braun W., Hepp K. Comm.Math.Phys.,1977,56, 2, 101-113.
- 3.Skorohod A.V. Stochastic equations for complicated systems , Moscow, Nauka, 1983 (in Russian)
- 4.Dobrushin R.L., Sinai Ya.G., Suhov U.M. in : Itogi nauki i tehniki, series "Modern problems of mathematics", v.2, 233-284; Moscow, VINITI,1985 (in Russian)
- 5.Spohn H., Rev.Mod.Phys.,1980, 53, 3, 569-615.
- 6.Skrypnik W.I.,Teor.Mat.Fiz.,1986, 69,.1, 128-141;1988(to appear)
- 7.Chueshov I.D., Teor.Mat.Fiz.,1986, 67, 2, 304-308.
- 8.Petrina D.Ya.,Gerasimenko V.I., Malyshev P.V. Mathematical foundations of Classical Statistical Mechanics , Kiev, Naukova Dumka, 1985 (in Russian) .

9. Ruelle D., Statistical Mechanics. Rigorous results. Benjamin
1970.

10. Dynkin E.B. Markov processes, Moscow, Φ M, 1963 (in
Russian)

