

Effect of *Nepeta menthoides* on cognitive disorders in Alzheimer's disease: a clinical trial

A. Abdolahadi¹, A. Nazeri², S. Esmaeili³, Z. Amiri⁴, M. Movahhed^{1*}

¹ Department of Traditional Medicine, School of Traditional Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran; ² Psychoses Center, University of Social Welfare and Rehabilitation Science, Tehran, Iran; ³ Traditional Medicine and Materia Medica Research Center, Department of Traditional Pharmacy, School of Traditional Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran; ⁴ Department of Basic Sciences, National Nutrition and Food Technology Research Institute, Faculty of Nutrition Sciences and Food Technology, Shahid Beheshti University of Medical Sciences, Tehran, Iran

*E-mail: minamovahed1970@gmail.com

Received for publication: 08 January 2017.

Accepted for publication: 15 April 2017.

Abstract

Nowadays, given the outbreak of Alzheimer disease (AD) as a global phenomenon, it is absolutely essential to take efficacious measures against it. Heating up the brain is a recommended approach to decrease the symptoms of AD. Furthermore, using hot herb is an effective method to enhance the temperature of the brain. One of the hot herbs is *Nepeta menthoides* which is known as Ostokhodus and has neuroprotective effects. In the present paper, the effect of *Nepeta* on the treatment of Alzheimer was studied. A trial was carried out on two groups of AD patients. While the first group was prescribed the capsules of *Nepeta* extraction, the second group was given the placebo capsules. The results of the taken MMSE inventories from both of the tested groups as the comparison criteria revealed that *Nepeta* had positive influence on the treatment of AD.

Keywords: *Nepeta menthoides*, Alzheimer's disease, cognitive disorder

Introduction

As the age of the population augments all over the world, the need for extra resources to care for individuals afflicted with Alzheimer Disease (AD) is felt more than ever. In 2006, the global pervasiveness of AD was 26.6 million. It will break out approximately 4 times by 2050; 1 in 85 persons will be affected by Alzheimer's (Brookmeyer, Johnson, Ziegler-Graham, & Arrighi, 2007). Alzheimer's is a progressive neurodegenerative disorder which causes cognitive impairment, neuropsychiatric symptoms, disability, dependency, caregiver burden, substantial healthcare, expenditure and premature death (Brookmeyer et al., 2011). Objectively, AD attacks recent memories, the executive functions and verbal fluency. Progression in the disease culminates in changes in emotions, intensification of the psychotic symptoms, depression and changes in personality. Early damage in the brain structure happens in hippocampus and the interconnected cortical zones which perform a role in the memory function (Scheff et al., 2015). In AD, the major pathological changes that occur in the brain are (i) extracellular amyloid plaques, containing mainly amyloid- β (A β) peptide, (ii) intraneuronal neurofibrillary tangles (NFTs), made of hyperphosphorylated and misfolded tau; other pathological changes linked with this neurodegenerative illness are (iii) oxidative stress, (iv) gliosis, (v) inflammation, (vi) dystrophy of neurons, (vii) neuronal death, (viii) synapse death, and (ix) altered levels of neurotransmitters (Amtul, 2015).

Alzheimer as a neurological disorder is related to the cold intemperament of the brain. In other words, it means that the brain's texture has lower temperature than its normal, healthy form. One strategy or approach known as the Iranian Traditional Medicine (ITM) recommends increasing the temperature of the brain by either physical methods or medical "hot" herbs. One of these "hot" herbs is *Nepeta menthoides* Boiss and Buhse (Ostokhodus in Persian language)(Ahmadian-Attar, Ahmadiani, Kamalinejad, Dargahi, & Mosaddegh, 2014).

Genus *Nepeta* of the Lamiaceae family consists of about 300 herbaceous species. *Nepeta* has delightful flowers. The greatest diversity of this genus species is distributed in Himalayas and the Southwestern Asia particularly Iran, Turkey and Hindu Kush. Iran, especially, is one of the centers of the birthplace of the *Nepeta*. There are sixty-seven species of this genus in Iran about 53% of which are endemics(Asgarpanah, Sarabian, & Ziarati, 2014).

The *Nepeta menthoides* Boiss and Buhse, usually referred to as Ostokhodus-e Khorasani or in brief Ostokhodus, is used as a medicine to cure neurological diseases like epilepsy and melancholia(Naghibi, Mosaddegh, Mohammadi Motamed, & Ghorbani, 2010). Moreover, this species has been claimed to have neuroprotective influences on the axotomized spinal motoneurons(Azizzadeh Delshad & Farzan, 2013; Delshad, Naseri, Parvizi, Fattah, & Sharayeli, 2011). For their antiasmatic, antiseptic, antispasmodic, antitussive, diuretic and febrifuge effects, diverse species of *Nepeta* have widespread use in the traditional medicine(Miceli et al., 2005). The experts in the traditional medicine take pride in stating that Ostokhodus wards off the production of the infection inside the body and also reinforces the heart and the urinary system. Ostokhodus, moreover, is useful for the chest (lung) and kidney ailments and is suitable for the health condition of the elders(Avicenna., 2005; Biruni, 2004; Nafis., 2008). In ITM, the organs of the body comprise an amalgam of quadruple humors named akhlat, safra (with warm and dry qualities), sauda (with cold and dry qualities), balgham (with cold and wet qualities) and dam (with warm and wet qualities); these are the mixture of four basic elements viz. water, air, fire and soil(Abdolahadi Azam & Esmaeili Somayeh, 2016). This herb removes the black bile or sauda from the heart, head and balgham or phlegm of the body. The black bile and phlegm are two types of humor (in Persian named khelt). Sauda has cold and dry qualities and phlegm has cold and wet qualities(Aghili-Shirazi, 1773; SI, 1976).

The therapeutic impacts of the *Nepeta* genus are mainly due to the high contents of the flavonoids and essential oils in their leaves' textures(Jamzad et al., 2003). Because of their potentiality to function as antioxidant and scavenger of free radicals, these components are applicable in medicine(Soobrattee, Neergheen, Luximon-Ramma, Aruoma, & Bahorun, 2005). Also there is no need to resort to other remedies to compensate for the probable side effects of consuming Ostokhodus(El-Bitar, 1877). As of yet, however, the probable therapeutic effects of *Nepeta menthoides* Boiss and Buhse on the AD have not been studied. Hence, in line with this desideratum for research in this uncharted territory and due to the widespread applications of *Nepeta menthoides* in Iran, the present study was performed as an attempt to scrutinize the possible therapeutic capacities of this genus for the treatment of Alzheimer's disease.

Materials and Methods

The oral capsule (500 mg) of *Nepeta menthoides*, elicited from its hydro alcoholic extraction ,after the microbial and qualitative control of the *Nepeta* sample, was prepared. Also the placebo capsules were prepared by inserting 500 mg of starch into every capsule. A placebo is an artificial or clinically ineffectual treatment for an ailment or other medical condition with the purpose of deceiving the recipient (<https://en.oxforddictionaries.com/definition/placebo>, 2017).

Patients and design of the study

The study was conducted in 2015 at Shahid Beheshti University of Medical Sciences, Tehran, Iran. Both Nepeta and placebo capsules were 500 mg. The entrance criteria adopted by the researchers for selecting the AD patients as participants in the trial were as follows:

The age of the population being between 65 and 85, the scores being below 20 in the MMSE inventories at the onset of the trial, being literate at least up to fifth grade, taking consent from patient or the executor of the patient, underlying no other psychological disease such as depression, having no severe disease except psychiatric and neurological disorders based upon the assessment of an specialist, consuming no drugs except nicotine and caffeine, having no allergic history to the cholinesterase enzyme restrainers, and having no heart disease, hysteria, asthma, peptic trauma and urinary blockade history.

A total of 78 sick individuals as a statistical society (39 people in every group) were selected under the supervision of a psychiatrist. In case the patients felt the tendency to quit the trial or some side effects pertinent to the prescribed capsules were observed on the patients, the affected people were excluded from the trial and the additional or required medications were performed on them. After providing the needed explanations and having the informed consent forms signed by the patients, the history and the results of the MMSE inventories were determined before the trial. The two groups didn't have any meaningful differences in factors of age (p-value=0.496), sex (p-value=1.000), education (p-value=0.644), physical activity (p-value=1.000) and body mass index (p-value=0.792). All the patients were randomly assigned to two groups. The prepared capsules were distributed randomly amongst the patients based upon the codes assigned to them and, the researcher was kept unaware of the coding process. The patients in the first group were prescribed 3 capsules daily with 500 mg content of Nepeta L. The second group was given the same amount of placebos. At the first step, the duration of the treatment was one month. At the end of the due time the history and new scores of the MMSE inventories were surveyed again. At the second phase, the clinical trial was carried out for another three months and the new elicited results were gauged. Since Alzheimer's could be a serious and life-threatening disease, both group members were prescribed the conventional Alzheimer medications as the primary drugs. The analysis of the patients' data, including the scores of the MMSE inventories, was done utilizing T-test through the SPSS software.

Results

Among the 78 individuals who were registered in this research, 48.7% were male and 51.3% were female. 52.6% of the patients had the age range of 65 to 75. In addition, 47.4% of them were between 75 and 85 years old. 93.6 % of the patients were educated up to high school and 6.4 % had bachelor's degree. Body mass index distribution for 75.6 % of the patients was lower than 25 and for 24.4% was higher than 25. Two groups didn't have statistically meaningful differences based upon the physical activity. In both of the groups 74.4% of members had mild physical activity. The scores and standard deviations of the MMSE inventory before the treatment are displayed in Table 2. The juxtaposition of the results indicates that there is no significant difference between the two groups before launching the trial (p-value=0.488). The differences between the garnered scores of the MMSE inventory are depicted in diagram 1. It has been revealed that the differences between MMSE scores prior to the consumption (MMSE1) and after one month of consumption (MMSE2) are more significant for group 2 relative to group 1 (p-value=0.000). This shows that Nepeta is effective in improving MMSE scores. In addition, the diagram displays that the difference between MMSE2 and the MMSE3 score- the score of MMSE after three months and after the discontinuation of the consumption- is also more for the Nepeta consuming group (Group 2) in comparison with the placebo consuming group (Group 1)(p-value=0.000). This reveals that MMSE3 score in group 2 has

decreased more sharply compared to group 1. The difference between MMSE1 and MMSE3 scores in the two groups is not statistically meaningful (p-value=0.716). Based upon the data in table 2, three months after stopping the trial, patients' minimal scores dropped down and exhibited meaningful differences *visa a vise* the scores obtained in the the second step. These scores, nevertheless, did not reveal meaningful and considerable differences with the primary scores of the patients. Sex, age, education, body mass index and physical activity seemed to have no significant effects on the test results of the MMSE inventory

Table 1. Distribution of gender, age, education, BMI, physical activity and their p-values in both groups under study

	Sex		Age		Education		BMI		Physical-Activity	
	Male	Female	65-75	75-85	Illiterate	Literate	<25	>25	Mild	Moderate
Group 1	19	20	19	20	37	2	30	9	29	10
Group 2	19	20	22	17	36	3	29	10	29	10
P-Value	1.000		0.496		0.644		0.792		1.000	




Diagram 1 -The differences of MMSE scores: (a) MMSE 2 - MMSE1, (b) MMSE2 - MMSE3 and (c) MMSE1 – MMSE3 between the placebo consuming group (Group 1) and Nepeta consuming group (Group 2)

Table 2. Average Minimental Scores and their differences with p-value for both groups under study

	Mini1	Mini2	Mini3	Mini2 - Mini1	Mini2 - Mini3	Mini1 - Mini3
Group 1	13.33	13.44	12.87	0.1026	0.5641	0.4615
Group 2	13.79	14.64	13.38	0.8462	1.2564	0.4103
p- value	0.488	0.075	0.432	0.000	0.000	0.716

Discussion

The present research scrutinized the efficacy of one herbal product as a helper for the treatment of the Alzheimer disease. Today, using herbal medicines are prevalent as a way to treat the neurodegenerative diseases (Kahkeshani, Khanavi, & Hadjiakhoondi, 2014). The achieved data unveiled that *Nepeta menthoides* might be an effective remedy for the patients who have succumbed to Alzheimer. This study revealed further the efficacies of *Nepeta menthoides* accompanied by the conventional medications for diminishing the impairing impacts of AD. Diagram 1-(a) shows that consuming *Nepeta* can be highly effective but Diagram 1-(b) shows that discontinuation of the consumption of *Nepeta* can adversely affect the cognitive abilities of the patients to the point that there would be no meaningful difference between the two groups after 3 months (Diagram 1-(c)). *Nepeta menthoides* is referred to as *Ostokhodus Khorasani* in the Iranian traditional medicine (Amin, 1991). It is normally used as medicine to treat high blood pressure, rheumatic pains and nervous ailments (Amin, 1991; Ghahreman & Okhovvat, 2010; Hiroyukimoteki, Yamada, Hirotakakatsuzaki, & Komiya, 2002; Joharchi & Amiri, 2012). The presence of some of the components such as terpineol, terpinolene, 1,8-cineole and limonene in the essential oil of *Nepeta menthoides* have the antitumor activity (Hassan, Gali-Muhtasib, Göransson, & Larsson, 2010), reducing the protein expression of AKT1 and inhibiting cell proliferation (Okumura, Yoshida, Nishimura, Kitagishi, & Matsuda, 2012), antimicrobial activity (Sivropoulou et al., 1997), inhibition of acetylcholinesterase (Gould, 1997) and anticancer activity (Miyazawa & Yamafuji, 2005). Based on previous researches 1,8-cineol, α -pinene, limonene, α -terpineol and β -terpinene are shown to play preventive role of acetyl cholinesterase activity. Acetyl cholinesterase had the key role in Alzheimer involution (Aazza, Lyoussi, & Miguel, 2011). 1,8-cineol, which includes the major composition of *Nepeta menthoides*, plays a substantial role as the inhibitor of the enzyme (Aazza et al., 2011; Ciftci, Ozdemir, Tanyildizi, Yildiz, & Oguzturk, 2011; Dohi, Terasaki, & Makino, 2009; Perry, Houghton, Theobald, Jenner, & Perry, 2000). The existence of myrcene, 1,8-cineole, limonene, α -terpinene and α -terpinolene in the composition of *Nepeta menthoides* makes it a powerful antioxidant (Kim et al., 2004). The antioxidant activity can proffer protection cells from the oxidation and following growth cycle malfunction (which leads to cancer and neurodegenerative problems) (Kim et al., 2004; Roberto, Micucci, Sebastian, Graciela, & Anesini, 2010; Ruela de Sousa et al., 2007).

Flavonoids, as the most significant pigments for flower coloration in various *Nepeta* species, have anti-allergic, anti-inflammatory, antimicrobial, anti-cancer (Tereschuk, Riera, Castro, & Abdala, 1997), anti-infective (Al-Saleh, Gamal El-Din, Abbas, & Saeed, 1997; Aladesanmi, Sofowora, & Leary, 1986; Asadi Balsin Sharif Abadi, Nasri, Amin, & Bidaran, 2013; Bourrel, Perineau, Michel, & Bessiere, 1993; Mahmoud, Jawad, Hussain, Al-Omari, & Al-Naib, 1989; Quarengi, Tereschuk, Baigori, & Abdala, 2000; Rauha et al., 2000; Singh & Nath, 1999; Tarle &

Dvorzak, 1990; Torrenegra, Ricardo, Pedrozo, & Fuentes, 1989), antibacterial and anti-fungal effects(Daulatzai, 2010; Holtzman & Simon, 2000). One of the factors influential in the progression of AD is hypothermia. The molecular changes demonstrated that a 14-day Cold Water Hypothermia induced tau hyperphosphorylation, apoptosis and the formation of beta-amyloid plaques(A Whittington, Papon, Chouinard-Decorte, & Planel, 2010; Aid, Langenbach, & Bosetti, 2008; Aid et al., 2010; Planel et al., 2004).

Moreover, some neuroinflammatory features of the AD demonstrated that the COX-II has some influence on neuroinflammation(Blais, Turrin, & Rivest, 2005). Some research has reported that the inhibition or genetic removal of the COX-II would exacerbate the neuroinflammation(Choi, Aid, Choi, & Bosetti, 2010) . The amount of the COXII protein is considerable in the early stages of AD; it, however, decreases with the progression of the disease . Hypothermia increases COX-II. Surprisingly however, along with memory impairment, the expression of COX-II decreased after 14-day Cold Water Hypothermi(Ahmadian-Attari et al., 2015; Delshad & Parvizi, 2014; Tuo, Tuailon, Shen, & Chan, 2004). Nepeta, as a hot herb, has the therapeutic potential for the treatment of the AD occurred by hypothermia effects. One study signaled that the impaired memory due to hypothermia at rats coerced to swim in cold water could be neutralized by prescribing the Nepeta menthoides decoction. 100 mg/Kg of NM reversed memory impairment as well as tau hyperphosphorylation(Ahmadian-Attari et al., 2015). Other studies have surveyed the effects of the *N.menthoides* essential oil on mice regaining their memories and also its neuroprotective impacts on rats(Kiyani et al., 2012; Sarahroodi et al., 2012). Previously, different researches have been performed on the influences of Nepeta through the biological samples such as rats(Ahmadian-Attar et al., 2014; Ahmadian-Attari et al., 2015; Delshad & Parvizi, 2014) bacteria [58] and fungi(Ezzatzadeh, Sofla, Pourghasem, Rustaiyan, & Zarezadeh, 2014). In the present paper, the curing effect of Nepeta on AD with human samples was done for the first time.

Conclusion

To sum up, the present study revealed that using *Nepeta menthoides* as a herbal remedy accompanied by the conventional medications for the treatment of the neurodegenerative diseases such as Alzheimer's seems to be an efficacious, economical and safe therapeutic approach. Performing a trial on a group of AD patients that were prescribed oral capsules containing extraction of Nepeta along with normal AD medications unveiled that this herb could be effective in decreasing the AD symptoms. Further studies, nonetheless, are warranted to proffer a more vivid picture and to generalize the results of the present study.

Acknowledgement

This article is based on a thesis submitted to Shahid Beheshti University of medical sciences with SBMU.REC.1393.549.163 code.

References

- A Whittington, R., Papon, M.-A., Chouinard-Decorte, F., & Planel, E. (2010). Hypothermia and Alzheimer's disease neuropathogenic pathways. *Current Alzheimer Research*, 7(8), 717-725.
- Aazza, S., Lyoussi, B., & Miguel, M. G. (2011). Antioxidant and antiacetylcholinesterase activities of some commercial essential oils and their major compounds. *Molecules*, 16(9), 7672-7690.
- Abdolahadi Azam, M. M., & Esmaeili Somayeh, A. N. a. (2016). Nootropic plants in Iranian traditional medicine; an overview. [Review]. *International Journal of Advanced Biotechnology and Research (IJBR)*, Vol-7(Special Issue - Number 4), 456-471.

- Aghili-Shirazi, M. (1773). *Qarabadin-e Kabir (Great Pharmacopoeia)*. Tehran: The Institute for Medical History-Islamic and Complementary Medicine, Iran University of Medical Sciences, Tehran, Iran.
- Ahmadian-Attar, M. M., Ahmadiani, A., Kamalinejad, M., Dargahi, L., & Mosaddegh, M. (2014). Chronic cold-water-induced hypothermia impairs memory retrieval and *Nepeta menthoides* as a traditional “hot” herb reverses the impairment. *Iranian journal of pharmaceutical research: IJPR*, 13(Suppl), 185.
- Ahmadian-Attari, M. M., Dargahi, L., Mosaddegh, M., Kamalinejad, M., Khallaghi, B., Noorbala, F., & Ahmadiani, A. (2015). Impairment of Rat Spatial Learning and Memory in a New Model of Cold Water-Induced Chronic Hypothermia: Implication for Alzheimer’s Disease. *Neurotoxicity research*, 28(2), 95-107.
- Aid, S., Langenbach, R., & Bosetti, F. (2008). Neuroinflammatory response to lipopolysaccharide is exacerbated in mice genetically deficient in cyclooxygenase-2. *Journal of neuroinflammation*, 5(1), 17.
- Aid, S., Silva, A. C., Candelario-Jalil, E., Choi, S.-H., Rosenberg, G. A., & Bosetti, F. (2010). Cyclooxygenase-1 and-2 differentially modulate lipopolysaccharide-induced blood–brain barrier disruption through matrix metalloproteinase activity. *Journal of Cerebral Blood Flow & Metabolism*, 30(2), 370-380.
- Al-Saleh, G., Gamal El-Din, A., Abbas, J., & Saeed, N. (1997). Phytochemical and Biological Studies of Medicinal Plants in Bahrain: The Family Chenopodiaceae—Part 2. *International journal of pharmacognosy*, 35(1), 38-42.
- Aladesanmi, A., Sofowora, A., & Leary, J. (1986). Preliminary biological and phytochemical investigation of two Nigerian medicinal plants. *International Journal of Crude Drug Research*, 24(3), 147-153.
- Amin, G. R. (1991). *Popular medicinal plants of Iran (Vol. 1): Iranian Research Institute of Medicinal Plants Tehran*.
- Amtul, Z. (2015). Neural plasticity and memory: molecular mechanism. *Reviews in the Neurosciences*, 26(3), 253-268.
- Asadi Balsin Sharif Abadi, S., Nasri, S., Amin, G., & Bidaran, S. (2013). Anti-inflammatory and anti-nociceptive effects of hydroalcoholic extract of *Nepeta menthoides* on pain in aerial parts in male mice. *Journal of Jahrom University of Medical Sciences*, 11(3), 1-9.
- Asgarpanah, J., Sarabian, S., & Ziarati, P. (2014). Essential oil of *Nepeta* genus (Lamiaceae) from Iran: a review. *Journal of essential oil research*, 26(1), 1-12.
- Avicenna. (2005). *The Canon in Medicine*. Beirut: Institute of al-A'lami li al-Matbooa.
- Azizzadeh Delshad, A., & Farzan, A. (2013). The prophylactic capacity of *Nepeta Menthoides* (*Ostokhodus*) in prevention of spinal motoneuron injury. *Journal of Kerman University of Medical Sciences*, 20(1).
- Biruni, A. (2004). *Al Seydaneh fi al Teb*. Tehran, Iran: University Press (in Persian).
- Blais, V., Turrin, N. P., & Rivest, S. (2005). Cyclooxygenase 2 (COX-2) inhibition increases the inflammatory response in the brain during systemic immune stimuli. *Journal of neurochemistry*, 95(6), 1563-1574.
- Bourrel, C., Perineau, F., Michel, G., & Bessiere, J. (1993). Catnip (*Nepeta cataria* L.) essential oil: analysis of chemical constituents, bacteriostatic and fungistatic properties. *Journal of Essential Oil Research*, 5(2), 159-167.
- Brookmeyer, R., Evans, D. A., Hebert, L., Langa, K. M., Heeringa, S. G., Plassman, B. L., & Kukull, W. A. (2011). National estimates of the prevalence of Alzheimer’s disease in the United States. *Alzheimer's & Dementia*, 7(1), 61-73.

- Brookmeyer, R., Johnson, E., Ziegler-Graham, K., & Arrighi, H. M. (2007). Forecasting the global burden of Alzheimer's disease. *Alzheimer's & dementia*, 3(3), 186-191.
- Choi, S., Aid, S., Choi, U., & Bosetti, F. (2010). Cyclooxygenases-1 and-2 differentially modulate leukocyte recruitment into the inflamed brain. *The pharmacogenomics journal*, 10(5), 448-457.
- Ciftci, O., Ozdemir, I., Tanyildizi, S., Yildiz, S., & Oguzturk, H. (2011). Antioxidative effects of curcumin, β -myrcene and 1, 8-cineole against 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin-induced oxidative stress in rats liver. *Toxicology and Industrial Health*, 27(5), 447-453.
- Daulatzai, M. A. (2010). Conversion of Elderly to Alzheimer's Dementia: Role of Confluence of Hypothermia and Senescent Stigmata—the Plausible Pathway. *Journal of Alzheimer's Disease*, 21(4), 1039-1063.
- Delshad, A. A., Naseri, M., Parvizi, M., Fattah, N., & Sharayeli, M. (2011). The Iranian traditional herbal medicine ostokhodus can prevent axotomy-induced apoptosis in spinal motoneurons in neonate rats. *Journal of Medicinal Plants Research*, 5(18), 4446-4451.
- Delshad, A. A., & Parvizi, M. (2014). The Neuroprotective Effect of *Nepeta menthoides* on Axotomized Dorsal Root Ganglion Sensory Neurons in Neonate Rats. *Journal of Basic and Clinical Pathophysiology*, 2(2), 13-20.
- Dohi, S., Terasaki, M., & Makino, M. (2009). Acetylcholinesterase inhibitory activity and chemical composition of commercial essential oils. *Journal of agricultural and food chemistry*, 57(10), 4313-4318.
- El-Bitar, I. (1877). *Mofradat Al Adwiya*: Cairo: Egyptian Government Press.
- Ezzatzadeh, E., Sofla, S. F. I., Pourghasem, E., Rustaiyan, A., & Zarezadeh, A. (2014). Antimicrobial Activity and Chemical Constituents of the Essential Oils from Root, Leaf and Aerial Part of *Nepeta asterotricha* from Iran. *Journal of Essential Oil Bearing Plants*, 17(3), 415-421.
- Ghahreman, A., & Okhovvat, A. (2010). Matching the old descriptions of medicinal plants with the scientific ones: Tehran, Iran: tehran univer-sity press.
- Gould, M. N. (1997). Cancer chemoprevention and therapy by monoterpenes. *Environmental Health Perspectives*, 105(Suppl 4), 977.
- Hassan, S. B., Gali-Muhtasib, H., Göransson, H., & Larsson, R. (2010). Alpha terpineol: a potential anticancer agent which acts through suppressing NF- κ B signalling. *Anticancer Research*, 30(6), 1911-1919.
- Hiroyukimoteki, H. H., Yamada, Y., Hirotakakatsuzaki, K. I., & Komiya, T. (2002). Specific induction of apoptosis by 1, 8-cineole in two human leukemia cell lines, but not a in human stomach cancer cell line. *Oncology Reports*, 9, 757-760.
- Holtzman, A., & Simon, E. (2000). Body temperature as a risk factor for Alzheimer's disease. *Medical hypotheses*, 55(5), 440-444.
- <https://en.oxforddictionaries.com/definition/placebo> (Producer). (2017).
- Jamzad, Z., Grayer, R. J., Kite, G. C., Simmonds, M. S., Ingrouille, M., & Jalili, A. (2003). Leaf surface flavonoids in Iranian species of *Nepeta* (Lamiaceae) and some related genera. *Biochemical systematics and ecology*, 31(6), 587-600.
- Joharchi, M. R., & Amiri, M. S. (2012). Taxonomic evaluation of misidentification of crude herbal drugs marketed in Iran. *Avicenna journal of phytomedicine*, 2(2), 105.
- Kahkeshani, N., Khanavi, M., & Hadjiakhoondi, A. (2014). Quality Control of *Nepeta menthoides* Boiss & Buhse using Micromorphological Analysis and Phytochemical Screening. *Journal of Medicinal Plants*, 1(49), 34-38.

- Kim, H.-J., Chen, F., Wu, C., Wang, X., Chung, H. Y., & Jin, Z. (2004). Evaluation of antioxidant activity of Australian tea tree (*Melaleuca alternifolia*) oil and its components. *Journal of agricultural and food chemistry*, 52(10), 2849-2854.
- Kiyani, N., Kiyani, A., Khademizadeh, M., Pazuki, M. R., Mamghani, F. A., & Dehshiri, A. M. (2012). *Nepeta menthoides* as memory enhancer. *Planta Medica*, 78(11), PI449.
- Mahmoud, M. J., Jawad, A.-L. M., Hussain, A. M., Al-Omari, M., & Al-Naib, A. (1989). In vitro antimicrobial activity of *Salsola rosmarinus* and *Adiantum capillus-veneris*. *International Journal of Crude Drug Research*, 27(1), 14-16.
- Miceli, N., Taviano, M., Giuffrida, D., Trovato, A., Tzakou, O., & Galati, E. (2005). Anti-inflammatory activity of extract and fractions from *Nepeta sibthorpii* Benth. *Journal of ethnopharmacology*, 97(2), 261-266.
- Miyazawa, M., & Yamafuji, C. (2005). Inhibition of acetylcholinesterase activity by bicyclic monoterpenoids. *Journal of agricultural and food chemistry*, 53(5), 1765-1768.
- Nafis., I. a. (2008). *Al-Shamil fi al-sina'a al-tibbiyya*. Tehran: Iran University of Medical Sciences.
- Naghbi, F., Mosaddegh, M., Mohammadi Motamed, M., & Ghorbani, A. (2010). Labiatae family in folk medicine in Iran: from ethnobotany to pharmacology. *Iranian Journal of Pharmaceutical Research*, 63-79.
- Okumura, N., Yoshida, H., Nishimura, Y., Kitagishi, Y., & Matsuda, S. (2012). Terpinolene, a component of herbal sage, downregulates AKT1 expression in K562 cells. *Oncology letters*, 3(2), 321-324.
- Perry, N. S., Houghton, P. J., Theobald, A., Jenner, P., & Perry, E. K. (2000). In-vitro inhibition of human erythrocyte acetylcholinesterase by *Salvia lavandulaefolia* essential oil and constituent terpenes. *Journal of pharmacy and pharmacology*, 52(7), 895-902.
- Planel, E., Miyasaka, T., Launey, T., Chui, D.-H., Tanemura, K., Sato, S., Takashima, A. (2004). Alterations in glucose metabolism induce hypothermia leading to tau hyperphosphorylation through differential inhibition of kinase and phosphatase activities: implications for Alzheimer's disease. *Journal of Neuroscience*, 24(10), 2401-2411.
- Quarenghi, M., Tereschuk, M., Baigori, M., & Abdala, L. (2000). Antimicrobial activity of flowers from *Anthemis cotula*. *Fitoterapia*, 71(6), 710-712.
- Rauha, J.-P., Remes, S., Heinonen, M., Hopia, A., Kähkönen, M., Kujala, T., . . . Vuorela, P. (2000). Antimicrobial effects of Finnish plant extracts containing flavonoids and other phenolic compounds. *International journal of food microbiology*, 56(1), 3-12.
- Roberto, D., Micucci, P., Sebastian, T., Graciela, F., & Anesini, C. (2010). Antioxidant activity of limonene on normal murine lymphocytes: relation to H₂O₂ modulation and cell proliferation. *Basic & clinical pharmacology & toxicology*, 106(1), 38-44.
- Ruela de Sousa, R. R., Queiroz, K. C. S., Souza, A. C. S., Gurgueira, S. A., Augusto, A. C., Miranda, M. A., . . . Aoyama, H. (2007). Phosphoprotein levels, MAPK activities and NFκB expression are affected by fisetin. *Journal of enzyme inhibition and medicinal chemistry*, 22(4), 439-444.
- Sarahroodi, S., Jafari-Najafi, R., Nasri, S., Rohampour, K., Maleki-Jamshid, A., & Esmaeili, S. (2012). Effects of *Nepeta menthoides* aqueous extract on retention and retrieval of memory in mice. *Pakistan journal of biological sciences: PJBS*, 15(22), 1085-1089.
- Scheff, S. W., Price, D. A., Ansari, M. A., Roberts, K. N., Schmitt, F. A., Ikonovic, M. D., & Mufson, E. J. (2015). Synaptic change in the posterior cingulate gyrus in the progression of Alzheimer's disease. *Journal of Alzheimer's Disease*, 43(3), 1073-1090.
- SI, J. (1976). *Zakhireh Kharazmshahi (Treasure of Kharazmshah)*. The Iranian Culture Foundation.

- Singh, R., & Nath, G. (1999). Antimicrobial activity of *Elaeocarpus sphaericus*. *Phytotherapy research*, 13(5), 448-450.
- Sivropoulou, A., Nikolaou, C., Papanikolaou, E., Kokkini, S., Lanaras, T., & Arsenakis, M. (1997). Antimicrobial, cytotoxic, and antiviral activities of *Salvia fruticosa* essential oil. *Journal of Agricultural and food Chemistry*, 45(8), 3197-3201.
- Soobrattee, M. A., Neergheen, V. S., Luximon-Ramma, A., Aruoma, O. I., & Bahorun, T. (2005). Phenolics as potential antioxidant therapeutic agents: mechanism and actions. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*, 579(1), 200-213.
- Tarle, D., & Dvorzak, I. (1990). Antimicrobial activity of the plant *Cirsium oleraceum* (L.) Scop. *Acta Pharm Jugosl*, 40(4), 569-571.
- Tereschuk, M. a. L., Riera, M. V., Castro, G. R., & Abdala, L. R. (1997). Antimicrobial activity of flavonoids from leaves of *Tagetes minuta*. *Journal of Ethnopharmacology*, 56(3), 227-232.
- Torrenegra, R. D., Ricardo, A. A., Pedrozo, J. P., & Fuentes, O. C. (1989). Flavonoids from *Gnaphalium gracile* HBK. *International Journal of Crude Drug Research*, 27(1), 22-24.
- Tuo, J., Tuillon, N., Shen, D., & Chan, C.-C. (2004). Endotoxin-induced uveitis in cyclooxygenase-2-deficient mice. *Investigative ophthalmology & visual science*, 45(7), 2306-2313.