

## On Total Vertex Irregularity Strength of Cocktail Party Graph

Kristiana Wijaya<sup>1)</sup>, Slamir<sup>2)</sup>, Mirka Miller<sup>3)</sup>

<sup>1)</sup>Mathematic Department Faculty of Mathematics and Natural Sciences University of Jember

<sup>2)</sup>Informatic System Department University of Jember

<sup>3)</sup>School of Electrical and Computer Science The University of Newcastle Australia

### ABSTRACT

A vertex irregular total  $k$ -labeling of a graph  $G$  is a function  $\lambda$  from both the vertex and the edge sets to  $\{1, 2, 3, \dots, k\}$  such that for every pair of distinct vertices  $u$  and  $x$ ,  $\lambda(u) + \sum_{uv \in E} \lambda(uv) \neq \lambda(x) + \sum_{xy \in E} \lambda(xy)$ .

The integer  $k$  is called the total vertex irregularity strength, denoted by  $tvs(G)$ , is the minimum value of the largest label over all such irregular assignments. In this paper, we prove that the total vertex irregularity strength of the Cocktail Party graph  $H_{2,n}$ , that is  $tvs(H_{2,n}) = 3$  for  $n \geq 3$ .

Keywords : Total vertex irregularity strength, cocktail party graph

### INTRODUCTION

Throughout this paper all graphs are finite, simple, and undirected. A vertex irregular total  $k$ -labeling on a graph  $G$  is a function  $\lambda$  from both the vertex and the edge sets to  $\{1, 2, 3, \dots, k\}$  such that the weights calculated at vertices are distinct. The weight of a vertex  $v$  in  $G$  is defined as the sum of the label of  $v$  and the labels of all the edges incident with  $v$ , that is,  $wt(v) = \lambda(v) + \sum_{uv \in E} \lambda(uv)$ .

The notion of the vertex irregular total  $k$ -labeling was introduced by Bača, et al. (2007). The total vertex irregularity strength of  $G$ , denoted by  $tvs(G)$ , is the minimum value of the largest label over all such irregular assignments.

Many total vertex irregularity strength of  $G$  had already been identified for some classes of graph. Bača, et al. (2007) proved that if a tree  $T$  with  $n$  pendant vertices and no vertices of degree 2, then  $\left\lfloor \frac{n+1}{2} \right\rfloor \leq tvs(T) \leq n$ .

Additionally, they gave a lower and an upper bound on total vertex irregular strength for any graph  $G$  with  $v$  vertices and  $e$  edges, minimum degree  $\delta$  and maximum degree  $\Delta$ ,  $\left\lfloor \frac{|V| + \delta}{\Delta + 1} \right\rfloor \leq tvs(G) \leq |V| + \Delta - 2\delta - 1$ . In the same paper, they proved that  $tvs(C_n) = \left\lfloor \frac{n+2}{3} \right\rfloor$ ,  $tvs(K_n) = \left\lfloor \frac{n+1}{2} \right\rfloor$ , and

$tvs(K_n) = 2$ . Furthermore, the total vertex irregularity strength of complete bipartite graphs  $K_{m,n}$  for some  $m$  and  $n$  had been discovered by Wijaya et al. (2005), namely,

$$tvs(K_{2,n}) = \left\lfloor \frac{n+2}{3} \right\rfloor \text{ for } n > 3, \quad tvs(K_{n,n}) = 3$$

for  $n \geq 3$ ,  $tvs(K_{n,n+1}) = 3$  for  $n \geq 3$ ,  $tvs(K_{n,n+2}) = 3$  for  $n \geq 4$ , and

$$tvs(K_{n,an}) = \left\lfloor \frac{n(a+1)}{n+1} \right\rfloor \text{ for all } n \text{ and } a > 1.$$

Besides, they gave the lower bound on  $tvs(K_{m,n})$  for  $m < n$ , that is

$$tvs(K_{m,n}) \geq \max \left\{ \left\lfloor \frac{m+n}{m+1} \right\rfloor, \left\lfloor \frac{2m+n-1}{n} \right\rfloor \right\}.$$

Wijaya and Slamir (2008) found the values of total vertex irregularity strength of wheels  $W_n$ , fans  $F_n$ , suns  $S_n$  and friendship graphs  $f_n$  by

$$\text{showing that } tvs(W_n) = \left\lfloor \frac{n+3}{4} \right\rfloor,$$

$$tvs(F_n) = \left\lfloor \frac{n+2}{4} \right\rfloor, \quad tvs(S_n) = \left\lfloor \frac{n+1}{2} \right\rfloor,$$

$$tvs(f_n) = \left\lfloor \frac{2n+2}{3} \right\rfloor. \text{ Ahmad et al. (in press)}$$

had determined total vertex irregularity strength of Halin graph. Whereas the total vertex irregularity strength of several types of trees determined by Nurdin et al. (to appears)

and the total vertex irregularity strength of disjoint union of  $t$  copies of a path had been determined by Nurdin *et al.* (2009). Ahmad & Bača (Gallian 2009) proved that

$$tvs(J_{n,2}) = \left\lceil \frac{n+1}{2} \right\rceil \text{ for } n \geq 4 \text{ and conjectured that for } n \geq 3 \text{ and } m \geq 3,$$

$$tvs(J_{m,n}) \geq \max \left\{ \left\lceil \frac{n(m-1)+2}{3} \right\rceil, \left\lceil \frac{nm+2}{4} \right\rceil \right\}. \text{ They also proved that for the circulant graph,}$$

$$tvs(C_n(1,2)) = \left\lceil \frac{n+4}{5} \right\rceil, \text{ and conjectured that for the circulant graph } C_n(a_1, a_2, \dots, a_m) \text{ with degree at least } 5, 1 \leq a_i \leq \left\lfloor \frac{n}{2} \right\rfloor,$$

$$tvs(C_n(a_1, a_2, \dots, a_m)) = \left\lceil \frac{n+r}{1+r} \right\rceil.$$

In this paper, we determine the total vertex irregularity strength of the cocktail party graph  $H_{2,n}$  for  $n \geq 3$ , as described in the following section.

**RESULTS AND DISCUSSION**

The cocktail party graph  $H_{2,n}$ ,  $n \geq 3$  is a graph with a vertex set  $V = \{v_1, v_2, \dots, v_{2n}\}$  partitioned into  $n$  independent sets  $V = \{I_1, I_2, \dots, I_n\}$  each of size 2 such that  $v_i v_j \in E$  for all  $i, j \in \{1, 2, \dots, 2n\}$  where  $i \in I_p, j \in I_q, p \neq q$ . It is the complement of a disjoint union of  $n$  copies of  $K_2$ . We present the total vertex irregularity strength of the cocktail party graph  $H_{2,n}$  for  $n \geq 3$ , as follows.

Theorem 1.  $tvs(H_{2,n}) = 3$  for  $n \geq 3$ .

Proof. Let  $H_{2,n}$  be the cocktail party graph with  $n \geq 3$ . Then  $H_{2,n}$  has  $2n$  vertices of degree  $2n-2$ . The smallest weight of vertices of  $H_{2,n}$  must be  $2n-1$  and the largest weight of vertices of  $H_{2,n}$  is at least  $4n-2$ . Since  $4n-2$  is obtained from the sum of  $2n-1$  numbers, then at least all edges and one vertex must have label  $\frac{4n-2}{2n-1} = 2$ . This is impossible

because there must be at least one edge with label 1 to obtain lower weight than  $4n-2$ .

Thus  $tvs(H_{2,n}) \geq 3$ .

To show that  $tvs(H_{2,n}) \leq 3$ , we label the vertices and edges of  $H_{2,n}$  as a vertex irregular total 3-labeling. Suppose the cocktail party graph has the set of vertices,

$$V(H_{2,n}) = \{u_1, u_2, \dots, u_n, v_1, v_2, \dots, v_n\} \text{ for } i = 1, 2, \dots, n \text{ and the set of edges,}$$

$$E(H_{2,n}) = \{u_i u_j\}_{i \neq j} \cup \{u_i v_j\}_{i \neq j} \cup \{v_i v_j\}_{i \neq j}$$

The labels of the vertices of  $H_{2,n}$  are described in the following formulas:

$$\lambda(u_i) = 1, \text{ for } i = 1, 2, \dots, n. \\ \lambda(v_i) = \begin{cases} 1, & \text{for } i = 1, 2, \dots, \left\lfloor \frac{n}{2} \right\rfloor \\ 2, & \text{for } i = \left\lfloor \frac{n}{2} \right\rfloor, \text{ if } n \text{ is odd} \\ 3, & \text{for } i = \left\lfloor \frac{n}{2} \right\rfloor + 1, \left\lfloor \frac{n}{2} \right\rfloor + 2, \dots, n. \end{cases}$$

The labels of edges of  $H_{2,n}$  are as follows.

For  $i \neq j = j$ , where  $i, j = 1, 2, \dots, n$ .

$$\lambda(u_i u_j) = 1, \\ \lambda(v_i v_j) = 2, \\ \lambda(u_i v_j) = \begin{cases} 1, & \text{if } i < j \text{ and } i + j \leq n + 1, \\ 1, & \text{if } i > j \text{ and } i + j \leq n, \\ 2, & \text{if else.} \end{cases}$$

The weights of vertices  $u_i$  and  $v_i$  are:

$$wt(u_i) = 2n - 2 + i \text{ for } i = 1, 2, \dots, n; \\ wt(v_i) = 3n - 2 + i \text{ for } i = 1, 2, \dots, n.$$

It is easy to see that the weights calculated at vertices are distinct. So, the labelling is vertex irregular total. Therefore  $tvs(H_{2,n}) = 3$  for  $n \geq 3$ . □

As an illustration, we show the vertex irregular total 3-labeling of  $H_{2,6}$  (in Table 1) and  $H_{2,7}$  (in Table 2). We note that label 0 means that there is no edge connecting two vertices.

Table 1. The vertex irregular total 3-labeling of  $H_{2,6}$ .

$H_{2,6}$	$u_1$	$u_2$	$u_3$	$u_4$	$u_5$	$u_6$	$v_1$	$v_2$	$v_3$	$v_4$	$v_5$	$v_6$	$wt$
	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>3</b>	<b>3</b>	<b>3</b>	
$u_1$	<b>1</b>	0	1	1	1	1	1	0	1	1	1	1	11
$u_2$	<b>1</b>	1	0	1	1	1	1	1	0	1	1	1	2
$u_3$	<b>1</b>	1	1	0	1	1	1	1	1	0	1	2	2
$u_4$	<b>1</b>	1	1	1	0	1	1	1	1	2	0	2	2
$u_5$	<b>1</b>	1	1	1	1	0	1	1	2	2	2	0	2
$u_6$	<b>1</b>	1	1	1	1	1	0	2	2	2	2	2	0
$v_1$	<b>1</b>	0	1	1	1	1	2	0	2	2	2	2	2
$v_2$	<b>1</b>	1	0	1	1	2	2	2	0	2	2	2	2
$v_3$	<b>1</b>	1	1	0	2	2	2	2	2	0	2	2	2
$v_4$	<b>3</b>	1	1	1	0	2	2	2	2	2	0	2	2
$v_5$	<b>3</b>	1	1	2	2	0	2	2	2	2	2	0	2
$v_6$	<b>3</b>	1	2	2	2	2	0	2	2	2	2	2	0
$wt$		11	12	13	14	15	16	17	18	19	20	21	22

Table 2. The vertex irregular total 3-labeling of  $H_{2,7}$ .

$H_{2,7}$	$u_1$	$u_2$	$u_3$	$u_4$	$u_5$	$u_6$	$u_7$	$v_1$	$v_2$	$v_3$	$v_4$	$v_5$	$v_6$	$v_7$	$wt$
	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>3</b>	<b>3</b>	
$u_1$	<b>1</b>	0	1	1	1	1	1	1	0	1	1	1	1	1	13
$u_2$	<b>1</b>	1	0	1	1	1	1	1	1	0	1	1	1	1	2
$u_3$	<b>1</b>	1	1	0	1	1	1	1	1	1	0	1	1	2	2
$u_4$	<b>1</b>	1	1	1	0	1	1	1	1	1	1	0	2	2	2
$u_5$	<b>1</b>	1	1	1	1	0	1	1	1	1	2	2	0	2	2
$u_6$	<b>1</b>	1	1	1	1	1	0	1	1	2	2	2	2	0	2
$u_7$	<b>1</b>	1	1	1	1	1	1	0	2	2	2	2	2	2	0
$v_1$	<b>1</b>	0	1	1	1	1	1	2	0	2	2	2	2	2	2
$v_2$	<b>1</b>	1	0	1	1	1	2	2	2	0	2	2	2	2	2
$v_3$	<b>1</b>	1	1	0	1	2	2	2	2	2	0	2	2	2	2
$v_4$	<b>2</b>	1	1	1	0	2	2	2	2	2	2	0	2	2	2
$v_5$	<b>3</b>	1	1	1	2	0	2	2	2	2	2	2	0	2	2
$v_6$	<b>3</b>	1	1	2	2	2	0	2	2	2	2	2	2	0	2
$v_7$	<b>3</b>	1	2	2	2	2	2	0	2	2	2	2	2	2	0
$wt$		13	14	15	16	17	18	19	20	21	22	23	24	25	26

### CONCLUSION

We conclude this paper with a result that the total vertex irregularity strength of the Cocktail Party graph  $H_{2,n}$ , that is  $tvs(H_{2,n}) = 3$  for  $n \geq 3$ . We also give an open problem, that is, determining the total vertex irregularity strength of generalization of cocktail party graph  $H_{m,n}$  for  $m > 2$  and  $n \geq 3$

### REFERENCES

- Ahmad A, Nurdin & Baskoro ET. On the irregular total labeling of generalized Halin graph. (In Press) *Ars Combinatoria*.
- Bača M, Jendrol' S, Miller M & Ryan J. 2007. On irregular total labellings. *Discrete Math.* **307**: 11-12, 1378–1388.
- Gallian JA. 2009. A Dynamic Survey of Graph Labeling. *The Electronic Journal of Combinatorics.* **16**.
- Nurdin, Baskoro ET, Salman ANM & Gaos NN, *On total vertex-irregular labellings for several types of trees, to appears in Utilitas Mathematica.*
- Nurdin, Baskoro ET, Salman ANM & Gaos NN. 2009. On the total vertex-irregular strength of a disjoint union of  $t$  copies of a path. *J. Combin. Math. Combin. Comput.* **71**: 227- 233.
- Wijaya K, Slamini, Surahmat, & Jendrol' S. 2005. Total vertex irregular labeling of complete bipartite graphs. *J. Combin. Math. Combin. Comput.* **55**: 129–136.
- Wijaya K & Slamini. 2008. Total vertex irregular labeling of wheels, fans, suns and friendship graphs. *J. Combin. Math. Combin. Comput.* **65**:103-112.