

OSP-IMMUNOFLUORESCENT REMYELINATING OLIGODENDROCYTES IN THE BRAINSTEM OF TOXICALLY-DEMYELINATED WISTAR RATS

Eliza Simone Viégas Sallis¹, Cinthia Melazzo Mazzanti², Alexandre Mazzanti¹, Luis Antonio Violin Pereira³, Kélen Fabíola Arroiteia³, Rafael Fustigatto⁴, Charles Pelizzari¹, Aline Rodrigues¹, Dominguita Lühers Graça¹

ABSTRACT - Central nervous system (CNS) remyelination following toxically-induced demyelination is a well known process. Oligodendrocytes constitute the bulk of the myelinating cells in the brain whereas Schwann cells overwhelm oligodendrocytes numbers in spinal cord remyelination. Despite the common knowledge of these facts, we still do not know completely the origin of both remyelinating cells. The present study investigated the participation of mature oligodendrocytes in remyelination after ethidium-bromide (EB) induced demyelination in the brainstem of normal and cyclosporin A-immunosuppressed Wistar rats. Thirty adult female rats were divided into three experimental groups. In group 1 the rats received a single intracisternal injection of 10 μ L of 0.1% ethidium bromide (EB) in 0.9% saline (n=10); in group 2 the rats received the EB injection while immunosuppressed with cyclosporin A (n=10); in group 3 the rats received a single 10 μ L injection of 0.9% saline while treated with cyclosporin A. The rats were killed at 15, 21 and 31 days after injection. Within the EB lesions, from 15 days onward many cells within the periphery of the lesions stained positive for OSP (oligodendrocyte specific protein) a marker for mature oligodendrocytes and myelin. This cell marking signals that, at least, part of the process of repairing the myelin sheaths is carried out by mature cells of the oligodendrocyte lineage.

KEY WORDS: toxic demyelination, remyelination, oligodendrocytes, oligodendrocyte specific protein (OSP), ethidium bromide.

Oligodendrócitos remielinizantes positivos para OSP - proteína específica do oligodendrócito no tronco encefálico de ratos Wistar desmielinizados toxicamente

RESUMO - A remielinização do sistema nervoso central após desmielinização tóxica é um processo bem conhecido. No encéfalo, os oligodendrócitos remielinizam uma área maior do que na medula espinhal, onde as células de Schwann são preponderantes. Embora esses fatos sejam bem conhecidos, ainda não se conhece com certeza a origem das células remielinizantes. Esta investigação foi desenhada para esclarecer a participação de oligodendrócitos maduros na reconstrução das bainhas perdidas após a desmielinização induzida por brometo de etídio (BE) no tronco encefálico de ratos Wistar normais e imunossuprimidos com ciclosporina A. Trinta ratos fêmeas adultas foram divididos em três grupos experimentais. No grupo 1, os ratos receberam uma injeção de 10 μ L de BE em 0,9% salina (n=10) na cisterna basal; no grupo 2, os ratos receberam a injeção de BE e foram tratados com ciclosporina A (n=10); no grupo 3 os ratos receberam uma injeção de 10 μ L de 0,9% salina e foram tratados com ciclosporina A. Os ratos foram sacrificados aos 15, 21 e 31 dias após a injeção. A partir dos 15 dias muitas células da periferia das lesões tiveram marcação positiva para OSP (proteína específica do oligodendrócito), marcador de oligodendrócitos maduros e mielina. Assim, foi possível comprovar que células maduras da linhagem oligodendroglial participam do processo de remielinização neste modelo gliotóxico.

PALAVRAS-CHAVE: desmielinização tóxica, remielinização, oligodendrócitos, proteína específica do oligodendrócito (OSP), brometo de etídio.

¹PPPGMV da Universidade Federal de Santa Maria RS, Brazil; ²PPGBQ da Universidade Federal do Rio Grande do Sul, Brazil; ³Departamento de Histologia e Embriologia da Universidade Estadual de Campinas (UNICAMP) SP, Brazil; ⁴Curso de Graduação em Medicina Veterinária da Universidade Federal de Santa Maria RS, Brazil.

Received 18 August 2005, received in final form 28 November 2005. Accepted 23 January 2006.

Dra. Dominguita L. Graça - Departamento de Patologia UFSM - 97105-900 Santa Maria RS - Brasil. E-mail: dlgraca@smail.ufsm.br

Oligodendrocytes are the myelin-producing cells of the central nervous system (CNS). They occur in both the gray matter- as satellites for neurons- and the white matter- as interfascicular cells. Yet, satellite cells have the potentiality to form myelin¹. Demyelination (loss of myelin of intact axons) due to local injection of gliotoxic agents proved to be useful to study the pathogenesis of remyelination². Among those agents, the fluorescent dye ethidium bormide (EB) has been used to induce demyelination in Wistar rats within the spinal cord^{3,4}, within the brain^{5,6} and, within the sciatic nerve⁷. In the CNS remyelination was carried out by oligodendrocytes and Schwann cells³.

Due to the presence of lymphocytes within the lesions, Bondan et al.^{6,8} investigated the response of immunosuppressed Wistar rats to a local injection of EB in the pons. The rats treated with either cyclophosphamide or cyclosporin A showed partial remyelination by oligodendrocytes and occasional Schwann cells intermingled with cystic cavities. At the time of the investigation, the origin of the remyelinating oligodendrocytes was not known.

It was the aim of this study to find out if the remyelinating cells were mature or immature oligodendrocytes by marking the cells with OSP⁹. OSP (oligodendrocyte-specific protein) is a transmembrane protein, the third most abundant CNS myelin protein contributing to 7% of the total myelin protein¹⁰, and expressed in mature reactive cells and myelin.

METHOD

Thirty female Wistar rats (250-300 g) were used. The rats were divided into three groups and all had an intracisternal injection of either EB or 0.9% saline as follows. In group 1 the rats received 10 µL of 0.1% EB in 0.9% saline in the basal cisterna; in group 2, the rats received the EB injection and were immunosuppressed with cyclosporin-A; in group 3, the rats received 10 µL of 0.9% saline while immunosuppressed with cyclosporin A.

The rats were anesthetized with ketamin and xylazine (5:1; 0.1 ml/100 g) and a burr-hole was made on the right side of the skull, 0.8 cm rostral to the fronto-parietal suture. The injections were made according to Pereira et al.⁵, with a hand-held Hamilton syringe through the burr hole in a vertical position and the contents freed when the needle reached the base of the skull, into the basal surface of the pons. The skin was sutured and the rats allowed to recover. The rats from group 2 were injected cyclosporin A (Sandimun[®]) intraperitoneally, 10 mg/Kg on a daily basis for seven days and subsequently three times a week with a 48 h interval. The first cyclosporin A injection was made soon after surgery. The rats were maintained in collective cages (3-5 rats) and fed ration and water ad libitum.

For each group the rats were perfused under deep anes-

thesia with 10% buff e red formaline via the left ventricle at 15 (3) 21 (4) and 31 days (3) after injection (a.i). Brainstem coronal slices with the lesion were separated into two matching portions: one was immersed in TissueTek[®] and frozen at -80°C, the other half was embedded in paraffin for routine processing.

For the immunofluorescence studies, 8 to 12 µm frozen sections were obtained in cryostat and allowed to dry for 1h at room temperature (RT) before the immunoreaction. Selected sections were post-fixed with methanol for 2 minutes at -20°C and allowed to dry. To blocking non-specific sites, the sections were incubated with 0.05M tris-buffer-saline (TBS) pH 7.4 added with 1% bovine serum albumin (BSA) for 30 minutes at room temperature and washed with sequential incubations with 0.05M TBS. To make membranes permeable, the sections were incubated with 0.05M TBS added with 1% BSA and 0.1% of triton-X for 30 minutes at room temperature and washed with 0.05M TBS. The sections were then incubated with anti-OSP primary antibody diluted 1:100 in 0.05 M TBS with 1% BSA and 0.01% triton-X, overnight at 4°C. After washing with 0.05M TBS, the sections were incubated with fluorescein conjugated (FITC) anti-rabbit secondary antibody diluted 1:100 in 0.05M TBS for 60 minutes at room temperature and protected from light. The sections were washed with 0.05M TBS and mounted in fluorescent medium (Vectashild). Glass coverslips were walled with enamel. Fluorescence of the slides was observed with a Nikon fluorescence photomicroscope and the images were processed in the Image software Adobe Photoshop.

Paraffin embedded tissues were trimmed at 6 µm, dehydrated with increased concentrations of ethanol, stained with hematoxylin and eosin (H&E) and Weill staining, mounted in Entellan[®] with glass coverslips and studied and photographed under an Olympus BX41 light microscope.

RESULTS

EB induces a demyelinating lesion following destruction of glial cells as previously demonstrated by our group³⁻⁶. Naked axons undergo remyelination by oligodendrocytes and occasional Schwann cells. Within the brain, most of remyelination is carried out by oligodendrocytes that are firstly recognized in the lesions around day 13 after EB injection⁶.

In the H&E-stained lesions of this investigation, many round nuclei cells, interpreted as glial cells, slender cell processes and abundant gitter cells were found at 15 days a.i. (Fig 1A). Weill staining emphasized new myelin being produced as the lesions developed (Fig 1B).

In normal Wistar rats as well as cyclosporine A-immunosuppressed Wistar rats, remyelination was well advanced at 21 days after EB injection in groups 1 and 2. Many cells related to axons and thin new sheaths could be seen. From 15 days onward OSP-immunofluorescent oligodendrocytes were conspicuous within the lesions. A strong immunofluorescence mar-




Fig 1. 21 days lesion from a group 2 rat within the pons. A. Many round-nuclei glial cells are observed in this rather compact demyelinated lesion. H&E, 100X: B. New thin myelin sheaths (arrow) stain positive round the periphery of the lesion. Weil, 200X.

ked broadly branched cells (Fig 2) that extended processes in an expanded extracellular space where repair of myelin sheaths was under way. Immunofluorescent protein localized primarily within the outer membranes and appeared as concentrated both on the cell body as on the processes.

At all times in sections from groups 1 and 2, some round cells with eosinophilic dense cytoplasm within the tissue, interpreted as immature oligodendrocytes in EM studies⁸ did not mark positive for OSP. Both, mature and immature cells in normal as in immunosuppressed rats, lied in areas where astrocytic processes were conspicuous.

The lesions induced by the saline injection in group 3 consisted on a mild traumatic lesion along the needle track that induced astrocytic isomorphic gliosis.

DISCUSSION

CNS demyelination induced by EB is followed by remyelination by oligodendrocytes and Schwann cells. The area of Schwann cell remyelinated sheaths is larger in the spinal cord than in the brain due to regional differences in numbers of astrocytes within the CNS¹¹. The early disappearance of astrocytes and the glial limiting membrane after EB injection allows Schwann cells




Fig 2. Large broadly branched OSP-immunofluorescent oligodendrocytes from the periphery of the lesion in Fig 1, 400X.

to invade the CNS in order to remake the lost sheaths after myelinating cells commit to the naked axons^{12,13}. Once oligodendrocytes become the main myelin repairing cells within the brain, it was relevant to find out if they are either mature or immature considering the postmitotic nature of these cells.

In EB-induced lesions in the brainstem of normal Wistar rats oligodendrocytes remyelinate over the narrow area lining the normal tissue⁵, suggesting that mature cells with a functional reserve rebuild the sheaths. After EB injection in cyclosporin A-immunosuppressed Wistar rats, many round oligodendrocytes with large amounts of rough endoplasmic reticulum approach the naked axons. These cells are very suggestive of immature oligodendrocytes⁸, which could be derived from a common progenitor that gives rise to astrocytes and oligodendrocytes¹⁴. Confirmation of their condition of newly differentiated cells of the lineage will be provided by NG2 labelling¹⁵.

The fact that some oligodendrocytes label positive for fluorescent OSP signals to the formation of tight junctions between adjacent myelin lamellae made by quiescent nonproliferating and nonmigratory mature reactive cells from the area¹⁶, which constitute a source to remyelinate the demyelinated axons as previously suggested¹⁷. It is proposed that these mature cells initiate remyelination while newly differentiated cells reach the status to proceed the repair of the lost sheaths.

Both cells, mature and immature, and in normal and immunosuppressed animals, lie in areas where

astrocytic processes are conspicuous, confirming the need of astrocytes as the third element of the CNS for a stable relationship between axons and oligodendrocytes¹³.

In chronically demyelinated lesions as those of multiple sclerosis, the lack of remyelination may be ascribed more to the unchecked chronic immune reaction and less to the glial scar that does not completely hinder myelinating cells migration toward the myelin-demanding naked axons¹⁸. The discovery of progenitor oligodendrocytes within the adult brain even within multiple sclerosis lesions¹⁹, brought in some hope although the complex molecular environment in which remyelination takes place remains largely unknown. Molecular biology techniques may be of great help to find out which factors concerning remyelination are dysregulated in either chronically or recurrently demyelinated lesions²⁰.

Acknowledgements - The authors appreciate the technical assistance of Maria Andreia Inkelmann, DVM and João Francisco Nunes.

REFERENCES

1. Ludwin SK. The function of the perineuronal satellite oligodendrocyte: an immunohistochemical study. *Neuropath Appl Neurobiol* 1984;10: 143-149.
2. Woodruff RH, Franklin RJM. Demyelination and remyelination of the caudal cerebellar peduncle of adult rats following stereotaxic injections of lysolecithin, ethidium bromide, and complement/anti-galactocerebroside: a comparative study. *Glia* 1999;25:216-228.
3. Graça DL, Blakemore WF. Delayed remyelination of rat spinal cord following ethidium bromide injection. *Neuropath Appl Neurobiol* 1986;12: 593-608.
4. Fernandes CG, Pereira LAV, Graça DL. Inflammatory response of the spinal cord to multiple episodes of blood-brain barrier disruption and toxic demyelination in Wistar rats. *Braz J Med Biol Res* 1998; 31: 933-936.

5. Pereira LAV, Dertkigil MSJ, Graça DL, Cruz-Höfling MA. Dynamics of remyelination in adult rat brain after exposure to ethidium bromide. *J Submicr Cytol Pathol* 1998;30:297-301.
6. Bondan EF, Lallo MA, Sinhorini IL, Pereira LAV, Graça DL. The effect of cyclophosphamide on brainstem remyelination following local ethidium bromide injection in Wistar rats. *J Submicr Cytol Pathol* 2000;32:431-438.
7. Riet-Correa G, Fernandes CG, Pereira LAV, Graça DL. Ethidium bromide-induced demyelination of the sciatic nerve of adult Wistar rats. *Braz J Med Biol Res* 2002;35:99-104.
8. Bondan EF, Lallo MA, Graça DL. Efeitos do brometo de etídio no tronco encefálico de ratos Wistar imunossuprimidos com ciclosporina. *Cad Est Pesq UNIP* 1998;IV:1-46.
9. Bronstein JM, Popper P, Micevych PE, Farber DB. Isolation and characterization of a novel oligodendrocyte-specific protein. *Neurology* 1996;47:772-778.
10. Bronstein JM, Micevych PE, Cehn K. Oligodendrocyte-specific protein (OSP) is a major component of CNS myelin. *J Neurosci Res* 1997;50:713-720.
11. Sims TJ, Gilmore SA, Waxman SG, Klinge E. Dorsal-ventral differences in the glia limitans of the spinal cord: an ultrastructural study in developing and irradiated rats. *J Neuropathol Exp Neurol* 1985;44:415-429.
12. Franklin RJ, Blakemore WF. Requirements for Schwann cells migration within CNS environments: a viewpoint. *Int J Dev Neurosci* 1993;11:641-649.
13. Graça DL, Bondan EF, Pereira LAV, Fernandes CG, Maiorka PC. Behaviour of oligodendrocytes and Schwann cells in an experimental model of toxic demyelination of the central nervous system. *Arq Neuropsiquiatr* 2001;59:358-361.
14. Shigeko F, Shirabe T. The reaction of glial progenitor cells in remyelination following ethidium bromide-induced demyelination in the mouse spinal cord. *Neuropathology* 2002;22:233-242.
15. Levine JM, Reynolds, R. Activation and proliferation of endogenous oligodendrocyte precursor cells during ethidium bromide-induced demyelination. *Exp Neurol* 1999;160:333-347.
16. Tiwari-Woodruff SK, Buznikov AG, Vu TQ, et al. OSP/Claudin-11 forms a complex with a novel member of the tetraspanin super family and $\beta 1$ integrin and regulates proliferation and migration of oligodendrocytes. *J Cell Biol* 2005;153:295-305.
17. Pereira LAV, Crua-Höfling MA, Dertkigil MSJ, Graça DL. Biology of the repair of central nervous system demyelinated lesions. *Arq Neuropsiquiatr* 1996;54:331-334.
18. Noble M, Atalio P, Barnett SC, et al. Development, regeneration and neoplasia of glial cells in the central nervous system. *An NY Acad Sci* 1991;633:35-47.
19. Chang A, Nishiyama A, Peterson J, Prineas J, Trapp BD. NG2 positive oligodendrocyte progenitor cells in adult human brain and multiple sclerosis lesions. *J Neurosci* 2000;20:6404-6412.
20. Franklin RJM. Why does remyelination fail in multiple sclerosis? *Nature Rev Neurosci* 2002;3:705-771.