

Isolation and Characterization of Selenate Resistant Mutants of *Acremonium chrysogenum*

Airton Vialta^{1*}, Cleide F. Catani², Renato B. Junior¹ and João L. Azevedo³

¹Instituto de Tecnologia de Alimentos. Avenida Brasil, 2880. CEP.13073-001.Campinas-SP, Brasil; ²Universidade Estadual de Campinas. Departamento de Microbiologia e Imunologia. CEP 13083-970. Campinas- SP, Brasil;

³ESALQ - USP. Departamento de Genética. CEP 13418-900. Piracicaba - SP, Brasil.

ABSTRACT

Mutants unable to convert exogenous sulfate to sulfite were isolated using the toxic analogue selenate. Three of twenty-eight isolated mutants were chromate sensitive. They showed a possible lesion in the gene that codes the ATP sulfurylase. The others were chromate resistant, and probably had a lesion in one or both of the genes that code the sulfate permease. Methionine increased the resistance levels to selenate. In addition, the frequency of spontaneous mutants obtained in a medium containing methionine was higher (between 2.4×10^{-6} and 18.0×10^{-6}) than that obtained using a medium without any intentional source of sulfur (between 0.7×10^{-6} and 5.0×10^{-6}). The original strain, as well as the mutants, were able to grow in a sulfur-free liquid medium even after 4 consecutive inoculation procedures. These results indicated the existence of sulfur traces in the medium and/or an efficient intracellular storage system. There was no significant difference between cephalosporin C production in mutants and the original strain.

Key words: Selenate resistant mutant, *Acremonium chrysogenum*, cephalosporin C

INTRODUCTION

Microbiological and biochemical analyses of *Acremonium chrysogenum* mutants for sulfur metabolism showed that wild strains of this species probably have three regulation mechanisms, two of which are related to the conversion of sulfate to cysteine, where the control is carried out by methionine. The third, *O*⁻ acetyl serine, induces the enzymes that convert sulfate to cysteine (Queener *et al.* 1984). Methionine also plays a fundamental role in the biosynthetic pathway of cephalosporin C. It is the major donor of sulfur atoms, in addition to the stimulation of antibiotic synthesis (Nüesch *et al.* 1973; Drew & Demain, 1975). Using *Aspergillus nidulans*, it is possible to isolate mutants that cannot use toxic analogues, such as selenate and chromate, to convert exogenous sulfate to sulfite (Arst, 1968). The most common selenate resistant mutations in *A. nidulans* result in either loss of sulfate permease, produced by the *sB* gene, or of ATP sulfurylase, produced by the *sC* gene. These two mutations can be easily distinguished. *sB*⁻ mutants are both chromate

and selenate resistant, whereas the *sC*⁻ mutants are selenate resistant and chromate sensitive (Arst, 1968). *sC*⁻ mutants of *A. nidulans*, *A. niger* and *A. oryzae* were transformed by complementation, using the *sC* gene isolated from *A. nidulans* (Buxton *et al.*, 1989; Yamada *et al.*, 1997). When the transformants were plated on minimal medium containing sulfate as the sole source of sulfur, they showed normal growth.

This system has been studied for yeasts in the last few years. Smith *et al.* (1995) isolated a mutant of *Saccharomyces cerevisiae*, deficient in its capacity to transport sulfate into the cells, obtained a clone which complemented this mutation and identified a cDNA fragment that encoded a high affinity sulfate transporter, responsible for the transfer of sulfate across the plasma membrane from the external medium. Other genes related to sulfate transport were isolated (Cherest *et al.*, 1997) and a structural relationship between ATP sulfurylase and sulfate membrane transporters was proposed (Logan *et al.*, 1996). Treichler *et al.* (1978)

* Author for correspondence

carried out studies using strain 8650-S⁻ of *A. chrysogenum*, which was not able to absorb inorganic sulfate, but showed growth in other sulfur sources such as sulfite, thiosulfate and sulfide. Despite its incapacity to metabolize sulfate, strain 8650-S⁻ exhibited a residual growth in media without sulfate, while wild strains showed a vigorous residual growth. The existence of residual growth in selective medium may be a problem during the characterization of mutants or during the isolation of transformant colonies. This growth was thought to be caused by sulfur compound traces that cannot be eliminated from chemically defined culture media (Treichler *et al.* 1978). These studies were usually carried out in liquid media, because agar contains significant quantities of covalently bound organic sulfur in a form that may support the growth of the strains (Queener *et al.* 1984).

The objectives of this study were to isolate selenate resistant mutants of the C-10 strain of *A. chrysogenum*, to check their level of resistance to selenate as well as their sensitivity to chromate, and to transform sC⁻ mutants by complementation, using a plasmid with the sC gene and genes that could cause an improvement in cephalosporin C production. Growth and titer levels of cephalosporin C of resistant mutants were determined and compared with those of the original strain.

MATERIALS AND METHODS

Microorganisms and maintenance: Strain C-10 of *Acremonium chrysogenum* (ATCC 48272) was used. Stock cultures were maintained on CM slants at 5°C.

Media used: Minimum medium (MM) contained (g.L⁻¹): sucrose, 20; NaNO₃, 3.0; K₂HPO₄, 0.5; KH₂PO₄, 0.5; KCl, 0.5; MgSO₄·7H₂O, 0.5; FeSO₄·7H₂O, 0.01, pH 7.2. Solid MM was prepared with 15 g.L⁻¹ of agar (Difco). Complete medium (CM) contained MM plus (g.L⁻¹): yeast extract, 4.0 and bactopectone (Difco), 4.0. Minimum medium without sulfur source (MMS⁻) was prepared same as MM, but replacing MgSO₄·7H₂O and FeSO₄·7H₂O with MgCl₂·6H₂O and FeCl₃·6H₂O. Seed medium (SM): SM contained (g.L⁻¹):

soluble starch, 40; corn steep liquor, 30; soybean meal, 10; soybean oil, 20; CaCO₃, 3.0; (NH₄)₂SO₄, 1.0, pH 7.0. Fermentation medium (FM) contained (g.L⁻¹): sucrose, 36; glucose, 27; yeast extract (Difco), 20; soybean oil, 24; CaSO₄, 7.5; CaCO₃, 10; (NH₄)₂SO₄, 8.0; D,L-methionine, 3.0, pH 6.4.

All the media were autoclaved at 121°C for 15 min. Incubations were carried out at 25°C (250 rpm) for 2 and 6 days for seed and fermentation (including mycelial growth in MM and MMS⁻), respectively.

In order to isolate and characterize resistant mutants, mycelia of 7 day old cultures of *A. chrysogenum* C-10 in MM were suspended in 6 ml of saline. Suspensions were transferred to test tubes containing glass beads and homogenized for 1 min. in a vortex. These suspensions were mixed with solid MMS⁻ at 45°C containing 10 mg.L⁻¹ of D,L-methionine and sodium selenate (0 to 102.4 mM) and transferred to petri dishes. After incubation, colonies with vigorous growth were isolated, and the mutation frequency estimated. Mutation frequencies were also obtained on MMS⁻ without D,L-methionine. In this case, selenate concentrations varied between 0 and 0.8 mM. The resistant mutants were characterized on MMS⁻ containing 2 mg.L⁻¹ of D,L-methionine and selenate at the following concentrations: 0, 0.1, 0.2, 0.4, 0.8 and 1.6 mM. In order to determine resistance or sensitivity to chromate, the suspensions were mixed with MMS⁻ containing 0.1 mM of sodium chromate.

Chromate resistant mutants were characterized pouring suspensions prepared as described above into MMS⁻ containing 10-mg.L⁻¹ of D, L-methionine and chromate at the following concentrations: 0.005, 0.01, 0.02, 0.04, 0.08 and 0.16 mM.

Suspensions prepared as described above were centrifuged, washed 3 times with saline, diluted and inoculated in to 250 ml erlenmeyers containing 25 ml of liquid MM in order to achieve a final concentration of 10⁶ c.f.u./ml. After incubation, the cultures were filtered and the dry weight was determined. One ml of

cultures grown in MMS⁻ was used to inoculate the erlenmeyers containing MM and MMS⁻. After incubation, 1 ml of cultures growing in both media was used to inoculate a new set of MM and MMS⁻. This procedure was repeated 4 times. This experiment was performed in duplicate.

Cephalosporin C production was determined by transferring 2.5 ml of the above mentioned suspension to a 500 ml erlenmeyer that contained 50 ml of seed medium. After incubation, 1.6 ml were used to inoculate the fermentation flasks. Fermentation was carried out in 500-ml erlenmeyers containing 40 ml of FM. Cephalosporin C was determined by HPLC after filtration through a 0.20 µm filter (Millipore). Analysis were carried out in a Varian (9010 and 9050) chromatograph with a Lichrospher RP-18 column (Merck) using a 254 nm detector. The mobile phase was 0.03% KH₂PO₄ (w/v). The Mann-Withney test (Roscoe, 1975) was used for the statistical analysis.

RESULTS

It was observed that concentrations higher than 12.8 mM completely inhibited the growth of sensitive colonies (Table 1). However, when MMS⁻ without D,L-methionine was used, a selenate concentration of only 0.05 mM (256 times lower) was sufficient to cause total inhibition (Table 2). These results showed clearly that the presence of D,L-methionine in MMS⁻ increased the resistance of the C-10 strain to selenate.

Colonies showing vigorous growth were observed in all plates containing selenate. These

colonies were classified as possible mutants and their frequencies estimated between 2.4×10^{-6} and 18.0×10^{-6} for experiments carried out with MMS⁻ containing D,L-methionine, and between 0.7×10^{-6} and 5.0×10^{-6} for MMS.

Table 1: Growth and frequency of possible spontaneous mutants in MMS⁻ containing different concentrations of selenate and 10 mg.L⁻¹ of D,L-methionine (1.33×10^6 c.f.u./plate)

Selenate (mM)	Growth ^a	Number of mutants per plate	Frequency of mutants (10^{-6})
0	+++	NE	
0.4	++	4.6	3.5
0.8	+	17.5	13.2
1.6	+	17.0	12.8
3.2	+	15.5	11.6
6.4	+	24.0	18.0
12.8	-	18.0	13.5
25.6	-	16.5	12.4
51.2	-	14.0	10.5
102.4	-	3.0	2.4

NE = not estimated, (a) +++ = vigorous growth, ++ = moderate growth, + = poor growth, - = only mutants grew

Table 2: Growth and frequency of spontaneous mutants on MMS⁻ containing different concentrations of selenate (1.4×10^6 c.f.u./plate).

Selenate (mM)	Growth ^a	Number of mutants per plate	Frequency of mutants ($\times 10^{-6}$)
0	+++	NE	
0.05	-	NE	
0.10	-	7	5.0
0.15	-	5	3.6
0.20	-	5	3.6
0.30	-	1	0.7
0.40	-	3	2.1
0.80	-	3	2.1

NE = not estimated, (a) +++ = vigorous growth, - = only mutants grew

Table 3: Dry weight (in mg) of the C-10 strain and mutants growing in liquid media (MM and MMS⁻) with agitation, in four serial replications.

Step	Mutants							
	SC1		SC2		SB1		C-10	
	MM	MMS ⁻	MM	MMS ⁻	MM	MMS ⁻	MM	MMS ⁻
1	118*±6	99±13	102±15	98±11	113±8	100±16	67±17	113±9
2	153±14	11±3	88±16	9±1	46±8	12±2	135±14	12±2
3	145±9	9±3	115±5	14±4	127±12	15±5	140±18	17±5
4	165±23	7±4	126±10	6±3	125±7	2±1	141±11	7±2

* average of 2 measures.

Thirty two possible mutants were isolated and purified from MMS⁻ and MMS⁻ with D,L⁻ methionine, both containing selenate. All isolates had resistance to selenate confirmed in MMS⁻ containing 2 mg.L⁻¹ of D,L⁻ methionine. A lower concentration of this aminoacid was used in order to utilize lower concentrations of selenate.

As previously commented, there is an inverse relationship between methionine concentration and sensitivity to selenate. In MMS⁻ supplemented with 2 mg.L⁻¹ of D,L-methionine, strain C-10 was completely inhibited by 0.1 mM of selenate. All mutants tested were able to grow in the highest concentration (1.6 mM).

Strain C-10 and all selenate resistant mutants were poured into MMS⁻ containing 10 mg.L⁻¹ of D,L-methionine and different concentrations of chromate, in order to detect their sensitivity to this ion. Only 3 selenate resistant mutants were chromate sensitive, inhibited by 0.04 mM of chromate (data not shown).

Strain C-10 and mutants SC1, SC2 and SB1, were tested in liquid MM and MMS⁻. They were able to grow without any intentional source of sulfur, even after four serial inoculations (Table 3). Although mutants had their sulfur metabolism pathways blocked, they showed growth similar to that of the strain C-10. The cephalosporin C titer of the 28 mutants tested were determined (Figure 1). A statistical analysis of the data showed that there was no significant difference in cephalosporin C

production between the mutants and the strain C-10 (Table 4).


Table 4: Statistical analysis of cephalosporin C production showed by C-10 strain and mutants using Mann – Withney nonparametrical test

C-10 strain	Mutants	Average	Standard Deviation	Z
n = 28	n = 28	μU = 392	δ = 60,99	0,631*
R = 836,5	= R = 759,5			
U = 353,5	= U = 430,5			

*Samples can be considered as coming from the same population (P>0,95).

DISCUSSION

It was observed that concentrations higher than 12.8 mM of selenate were necessary to inhibit the growth of sensitive colonies in media containing 10 mg.L⁻¹ of D,L-methionine. However, concentrations of 0.05mM, 256 times lower, were sufficient to inhibit growth in media without D,L-methionine. These results showed clearly that D,L-methionine increased the resistance of strain C-10 to selenate. Similar results were observed by Queener *et al.* (1984), who obtained a complete selenate toxicity reversion of M-8650 by adding 100mg.L⁻¹ of D,L-methionine to the medium. The level of tolerance to selenate shown by strain C-10 in MMS⁻ containing D,L-methionine, was higher than that found with *Aspergillus* sp. (Arst, 1968; Buxton *et al.*, 1989), whose inhibition levels were about 0.05 mM.



Cephalosporin C concentration (g.L⁻¹)

Figure 1: Cephalosporin C production shown by strain C-10 and mutants.

The frequencies observed of possible mutants were lower in media without D,L-methionine. As MMS⁻ did not contain any intentional source of sulfur, the cells were required to make use of their own reserves or of traces of this element that could be present in the medium. This could also explain the small number of mutants obtained with MMS⁻. An inverse relationship was observed between

methionine concentration and sensitivity to selenate. Similar results were obtained by Queener *et al.* (1984) working with strain M-8650 of *A. chrysogenum*. Strain C-10 showed levels of tolerance to selenate higher than those found in *Aspergillus* sp. (Arst, 1968; Buxton *et al.*, 1989).

Only three selenate resistant mutants were chromate sensitive, which therefore showed lesions in the *sC* gene, which encodes sulfate adenylyltransferase. The chromate resistant mutants had lost their sulfate permease and sulfate adenylyltransferase activities. This could be due to lesions in the *sB* genes, or in a combination of *sB* and *sC*, as observed with *A. nidulans* (Arst, 1968) and *A. niger* (Buxton *et al.*, 1989).

As previously mentioned, strain C-10 and mutants SC1, SC2 and SB1 were able to grow

without any intentional source of sulfur. These results pointed out the existence of sulfur traces in the medium and/or an efficient intracellular storage system for compounds containing this element. The mutants carrying the *sB* gene were unable to take up extracellular sulfate, and mutants carrying the *sC* gene did not metabolize it. Therefore, the cells probably used a reduced sulfur compound, like sulfite or thiosulfate, or an organic molecule that contained sulfur. Due to this characteristic, it was impossible to carry out the transformation using the *sC* system.

There was no significant difference in the cephalosporin C production between the mutants and strain C-10. These results could be explained by the presence of D,L-methionine in the fermentation medium. As discussed above, this aminoacid can be easily metabolized by mutants unable to absorb sulfate ions. Seemingly, this mutation does not affect either the growth or cephalosporin C production by the microorganism. Another possibility could be the utilization of intracellular stores.

Thus, the present study showed the successful isolation of the mutants resistant to selenate using strain C-10 of *A. chrysogenum*, and, despite their incapacity of metabolize sulfate,

they exhibited a residual growth in media without sulfate.

ACKNOWLEDGMENTS

The authors would like to thank Dr. Robert T. Rowlands from the Fermentation Services Division - PANLABS for providing strain C-10 and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for the RHAE fellowships.

RESUMO

Mutantes incapazes de converter o sulfato extracelular em sulfato foram isolados utilizando o análogo tóxico selenato. De 28 mutantes isolados, apenas 3 foram sensíveis ao cromato, provavelmente apresentando lesão no gene que codifica a ATP sulfúrilase. Os demais foram resistentes ao cromato e devem conter lesão no gene *sB* ou também no gene *sC*. A metionina elevou os níveis de resistência ao selenato e a frequência de mutantes espontâneos obtida em meio contendo este aminoácido foi maior (entre $2,42 \times 10^{-6}$ e $18,04 \times 10^{-6}$) do que a obtida no meio sem a adição de qualquer fonte intencional de enxofre (entre $0,71 \times 10^{-6}$ e $5,0 \times 10^{-6}$). A linhagem original e os mutantes foram capazes de crescer, mesmo depois de quatro etapas de inóculo, fato que pode ser explicado pela existência de traços do referido elemento no meio e/ou a presença de um sistema eficiente de estocagem intracelular. A produção de cefalosporina C foi estudada e a análise dos dados revelou que não houve diferença significativa entre os níveis produzidos pelos mutantes e os produzidos pela linhagem original.

REFERENCES

- Arst, H.N. (1968), Genetic analysis of the first steps of sulphate metabolism in *Aspergillus nidulans*. *Nature* (London), **219**, 268-270.
- Buxton, F.P.; Gwynne, D.I & Davies, R.W. (1989), Cloning of a new bidirectionally selectable marker for *Aspergillus* strains. *Gene*, **84**, 329-334.
- Cherest, H.; Daidian, J.C.; Thomas, D.; Benes, V.; Ansorge, W. & Surdin-Kerjan, Y. (1997), Molecular characterization of two high affinity sulfate transporters in *Saccharomyces cerevisiae*. *Genetics*, **145**, 627-635.
- Drew, S.W. & Demain, A.L. (1975), Production of cephalosporin C by single and double sulphur auxotrophic mutants of *Cephalosporium acremonium*. *Antimicrob. Agents Chemoth.*, **8**, 5-10.
- Logan, H.M.; Cathala, N.; Grignon, C. & Davidian, J. C. (1996), Cloning of a cDNA encoded by a member of the *Arabidopsis thaliana* ATP sulfurylase multigene family: Expression studies in yeast and in relation to plant sulfur nutrition. *J. Biol. Chem.*, **271**, 12227-12233.
- Nüesch, J.; Treichler, H.J. & Liersch, M. (1973), The biosynthesis of cephalosporin C. In - *Genetic of Industrial Microorganisms*, eds. Z. Vanek, Z. Hostalek & J. Cudlin, Elsevier Publishing Co., Amsterdam, pp. 309-334.
- Queener, S.W.; Wilkerson, S.; Tunin, D.R.; Mcdermott, J.R.; Chapman, J.L.; Nash, C.; Platt, C & Westpheling, J. (1984), Cephalosporin C fermentation: biochemical and regulatory aspects of sulphur metabolism. In - *Biotechnology of Industrial Antibiotics*, eds. E.J.Vandamme, vol. 22, Marcel Dekker Inc., N.Y., pp.141-170.
- Roscoe, J.T. (1975), *Fundamental Research Statistics for Behavioral Sciences*, 2nd ed., Holt, Rinehart and Winston Inc., New York.
- Smith, F.W.; Hawkesford, M.J.; Prosser, I.M. & Clarkson, D. T. (1995), Isolation of cDNA from *Saccharomyces cerevisiae* that encodes a high affinity sulphate transporter at the plasma membrane. *Mol. Gen. Genetics*, **247**, 709-715.
- Treichler, H.J.; Liersch, M. & Nüesch, J. (1978), Genetic and biochemistry of cephalosporin biosynthesis. In - *Antibiotics and Other Secondary Metabolites*, eds. Hütter, R. et al., FEMS Symposium 5., Academic Press, London, pp.177-199.
- Yamada, O.; Lee, B.R. & Gomi, K. (1997), Transformation system for *Aspergillus oryzae* with double auxotrophic mutations, *nia D* and *sC*. *Bioscience Biotechnol. Biochem.*, **61**, 1367-1369.

Received: March 16, 1999;
Revised: April 01, 1999;
Accepted: June 22, 1999.