



---

## One New Species and One New Subspecies of *Cissus* (Vitaceae) from Colombia

Author(s): Julio Antonio Lombardi

Source: *Novon*, Vol. 6, No. 2 (Summer, 1996), pp. 191-194

Published by: [Missouri Botanical Garden Press](#)

Stable URL: <http://www.jstor.org/stable/3391918>

Accessed: 22-06-2015 19:05 UTC

### REFERENCES

Linked references are available on JSTOR for this article:

[http://www.jstor.org/stable/3391918?seq=1&cid=pdf-reference#references\\_tab\\_contents](http://www.jstor.org/stable/3391918?seq=1&cid=pdf-reference#references_tab_contents)

You may need to log in to JSTOR to access the linked references.

---

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



Missouri Botanical Garden Press is collaborating with JSTOR to digitize, preserve and extend access to *Novon*.

<http://www.jstor.org>

---

# One New Species and One New Subspecies of *Cissus* (Vitaceae) from Colombia

Julio Antonio Lombardi

Departamento de Taxonomia e Morfologia Vegetais, Instituto de Biologia, Universidade Estadual de Campinas, Campinas, SP, Brazil. Current address: Departamento de Botânica, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, Av. Antônio Carlos 6627, 31270-110, Belo Horizonte, MG, Brazil

---

**ABSTRACT.** *Cissus colombiensis* and *C. verticillata* subsp. *colombiana*, new taxa of Vitaceae, are described from Colombia. In addition to descriptions and illustrations, comments on their probable relationships and distributions are presented.

*Cissus* L. has approximately 55 species in South America and is the largest genus in the Vitaceae. Many of the species described by Baker (1871; placed in the genus *Vitis*), and Planchon (1887) were considered synonyms in a recent taxonomic survey by the author (Lombardi, 1995), while one new species and one new subspecies were found in Colombia: *Cissus colombiensis* and *C. verticillata* subsp. *colombiana*.

***Cissus colombiensis* Lombardi, sp. nov. TYPE:**  
Colombia. Antioquia: Frontino, Corregimiento Nutibara, Región Murí, camino hacia La Blanquita, 1440 m, 10 July 1986, Acevedo et al. 1204 (holotype, NY). Figure 1.

Frutex scandens, foliis ternatis, hispidis, *C. ulmifoliae* (Baker) Planchon similis, sed inflorescentia haud umbelliformi, calyce urceolato, fructibus 1.5 cm longis, 1 cm latis, seminibus raphe rugosa, differt.

Liana, stem hispid, terete. Trichomes chiefly multicellular and not secretory, mixed with sparse non secretory and unicellular trichomes, and secretory trichomes. Tendrils with several bifurcate branches, hispid, adhesive discs not seen; scales ca. 0.15 cm long, triangular, puberulent, ciliate. Stipules ca. 0.4 cm long, 0.3 cm wide, subfalcate, fleshy, glabrous, the base gibbous, caducous. Petioles 1.4–8.5 cm long, terete, hispid to villous. Leaf blades ternately compound, much reduced in reproductive shoots; central petiolules 0.35–2.7 cm long, lateral petiolules 0–1.8 cm long, hispid to villous; blades of central leaflets subelliptic, elliptic to obovate, 4.3–18.8 cm long, 1.3–9.7 cm wide, blades of lateral leaflets subovate, subelliptic to elliptic, 3.6–14.35 cm long, 1.15–7.3 cm wide, central and lateral leaflets herbaceous, sparsely hispid

to hispid chiefly along the prominent veins on the abaxial side, sparsely puberulent along the veins on the adaxial side, the base subreniform, oblique, cuneate to rounded, the apex acuminate to acute, the margin denticulate. Compound cymes elongated, the peduncle 1.8–3.5 cm long, puberulent; the bracts triangular, 0.1–0.2 cm long, the base gibbous, puberulent, ciliate. Flowers greenish; pedicels 0.25–0.3 cm long, puberulent; calyx truncate, 0.1–0.15 cm long, 0.2–0.3 cm diam., urceolate, fleshy, glabrous; corolla tetramerous, 0.15–0.25 cm long, 0.15–0.3 cm diam., petals coherent, the margin elevated in the junction, caducous, papillose; stamens 4, connective deltate, granular, drying slightly dark brown, anther dehiscence extrorse; disc apex concave with central depression; the style conic, the stigma slightly clavate. Fruit purple, sub-spherical, with sparse lenticels, 1.5 cm long, 1 cm wide. Seed 1, subobconic, 1.1 cm long, 0.9 cm wide, the hilum acute, the raphe prominent and crisplate, the sides rugose.

This species is named after Colombia, the country with the most complete collections. It is distinctive among the South American trifoliate species by its large leaves and flowers, the hispid leaves prominently nerved, the elongate inflorescence, the urceolate calyx, and the peculiar seeds.

This species is similar to *Cissus ulmifolia* (Baker) Planchon, but differs in the urceolate calyx, the elongate inflorescence (umbelliform in *Cissus ulmifolia*), and in the seeds with a crisplate raphe and rugose sides (raphe not crisplate and sides deeply sulcate in *Cissus ulmifolia*).

The apparent disjunct distribution of *Cissus colombiensis* probably can be explained by a lack of collections of this species, a large liana in the rainforests of the Amazon basin and adjacent areas.

**Paratypes.** COLOMBIA. Boyacá: upper Chapon, 100 mi. NW de Bogotá, 6500 ft., high forest fronts, 8 Aug. 1932, Lawrence 397 (A—2 sheets, F, S). BRAZIL Amazonas: Novo Japurá, rio Japurá, margem direita, mata da várzea, 9 Nov. 1982, Amaral et al. 379 (NY); São Paulo

NOVON 6: 191–194. 1996.




Figure 1. *Cissus colombiensis* Lombardi. —A. Reproductive branch. —B. Stamens, ventral, lateral, and dorsal views. —C. Vegetative leaf. —D. Flower bud. —E. Stipule. —F. Nectariferous disc and pistil. —G. Seed. A–F from Acevedo et al. 1204; G from Lawrence 397. Scales A, C = 1 cm, B, D–G = 1 mm. Drawn by the author.




Figure 2. *Cissus verticillata* subsp. *colombiana* Lombardi. —A. Reproductive branch. —B. Stipule. A, B from Haught 4182. Scales A = 1 cm; B = 1 mm. Drawn by the author.

de Olivença, várzea, 21 May 1945, Fróes 20945 (F). BOLIVIA. La Paz: Nor Yungas, Serranía de Bella Vista, 17.6 km N of the bridge to Carrasco, 1600 m, 11 June 1985, Solomon 13980 (M, U), Solomon 13983 (U).

***Cissus verticillata* subsp. *colombiana* Lombardi, subsp. nov.** TYPE: Colombia. Magdalena: dry hills 12 km W of Codazzi, ca. 100 m, 26 May 1944, Haught 4182 (holotype, F; isotype, S). Figure 2.

*C. verticillatae* (L.) Nicolson & Jarvis subsp. *verticillatae* similis, sed folia in ramis vegetativis suborbicularia, in ramis fertilibus depresso-ovata, transverso-elliptica vel deltoidea, basi cuneata vel reniformia, stipularum margine flectere ad basem et apicem, et in inflorescentia 1.4–2.0 cm longo, 1.45–2.7 cm lato longo differt.

Stem glabrous. Tendrils bifurcate. Stipules ovate to spatulate, glabrous, the base rounded with reflected margin, the apex reflected. Leaves alternate; petioles 0.8–3 cm long, glabrous; blade simple and entire, in the vegetative shoots subcircular, in the reproductive shoots depressed ovate, transversely elliptic to deltate, 2–6.2 cm long, 1.9–7.1 cm wide. Inflorescence glabrous, pauciflorous, 1.4–2.0 cm long, 1.45–2.7 cm wide; the peduncle 0.55–1.85 cm long. Flowers greenish to sometimes reddish; pedicels 0.1–0.3 cm long; calyx 0.1 cm long, 0.1–0.15 cm diam.; corolla 0.1–0.2 cm long, 0.1–0.15 cm diam., slightly granulose. Fruit spherical, 0.4 cm diam.

*Cissus verticillata* is a very variable species, with a wide range of variation in indumentum, leaf shape, and rarely in flower color (pink to red in some collections); these phenotypic forms have been given several specific or infraspecific epithets (for a South American account see Lombardi, 1995).

Subspecies *colombiana* has leaves that are subcircular on vegetative shoots, and depressed ovate, transversely elliptic to deltate on reproductive shoots (in subspecies *verticillata* they are ovate, triangular, lanceolate, elliptic, obovate, wide-ovate to subpannulariform on vegetative shoots, and ovate, oblong to triangular on reproductive shoots); stipules with margin and apex reflected (only apex reflected in subspecies *verticillata*); and smaller and pauciflorous inflorescences (3.5–7.4 cm long, 3.0–5.9 cm wide, and multiflorous in subspecies *verticillata*).

The peculiar leaf shape, and the restricted geographic occurrence of this variation, not encountered in the rest of the distribution area of *Cissus verticillata* (south Florida to North Argentina), justify the recognition of subspecies *colombiana* as a distinct entity.

**Paratypes.** COLOMBIA. Atlántico: Magdalena, Santa Marta, 100 ft., Aug. 1898–1901, H. H. Smith 570 (BM); Santa Marta, near seacoast, Sep. 1898–1901, H. H. Smith 1319 (BR); Santa Marta, 500 ft., July 1898–1901, H. H. Smith 2724 (BM, BR, E, F, L, S, U). VENEZUELA. Aragua: Maracay, Facultad de Agronomía, cerro con bosque caducifolio, detrás del Instituto de Botánica Agrícola, 29 Aug. 1974, Guevara 1753 (F); Maracay, 1934, Vogl 1294 (M). Carabobo: environs de El Palito, 50 m, 24 Sep. 1920, Pittier 9084 (Z). Distrito Federal: bosque seco en las faldas costáreas frente el mar, entre Punta Picure y Oricao, 100–200 m, 1 July 1966, Steyermark & Aristeguieta 115 (F). Falcón: Dto. Silva, bosque tropófilo por encima de la línea de peñascos calcáreos, al S de la Punta Faustino, al SE de Chichiriviche, 15–25 m, 29 Aug. 1974, Steyermark & Manara 110404 (U). Nueva Esparta: Isla Margarita, 0–850 m, Aug. 1955, Bernardi 2391 (NY); Island of Margarita, 1 Aug. 1901, Miller 244 (ARIZ, F).

**Acknowledgments.** I thank the curators of the cited herbaria for the loan of specimens, G. W. Fernandes for the English revision, and G. M. Barroso for the Latin diagnoses. Work was partially sup-

ported by Comissão de Aperfeiçoamento do Pessoal do Ensino Superior (CAPES), and Universidade Federal de Minas Gerais (Pró-Reitoria de Pesquisa grant number: 23072.050311/94-99).

Literature Cited

- Baker, J. G. 1871. Ampelideae. Pp. 197–220 in C. F. P. Martius (editor), *Flora Brasiliensis*, Vol. 14. Frid. Fleischer, Leipzig.
- Lombardi, J. A. 1995. Typification of names of South American *Cissus* (Vitaceae). *Taxon* 44: 193–206.
- Planchon, J. E. 1887. Monographie des Ampélidées vraies. Pp. 305–654 in A. & C. DeCandolle (editors), *Monographiae Phanerogamarum*, Vol. 5. G. Masson, Paris.