

# Fumonisin B<sub>1</sub> and ochratoxin A in beers made in Brazil

## Fumonisina B<sub>1</sub> e ocratoxina A em cervejas fabricadas no Brasil

Luciane Mie KAWASHIMA<sup>1</sup>, Ana Paula VIEIRA<sup>1</sup>, Lucia Maria VALENTE SOARES<sup>1\*</sup>

### Abstract

Samples of beer made in Brazil were analyzed for the presence of fumonisin B<sub>1</sub> (FB<sub>1</sub>) and ochratoxin A (OTA). FB<sub>1</sub> was searched for in 58 beer samples from 30 plants located in nine states. The samples were concentrated and cleaned up with strong ion exchange column, derivatized with OPA and analyzed by HPLC with fluorescence detection. The limit of detection was 0.26 ng.mL<sup>-1</sup> and the average recovery was 98%. Twenty-five samples contained FB<sub>1</sub>, ranging from 1 to 40 ng.mL<sup>-1</sup>. Beer (123 samples) from 36 plants located in 5 states were analyzed for OTA by means of immunoaffinity column cleanup followed by liquid chromatography associated with fluorescence. The detection limit was 0.1 ng.mL<sup>-1</sup> and the average recovery was 92%. Five samples contained OTA in concentrations from 1 to 18 ng.mL<sup>-1</sup>. The results indicate that FB<sub>1</sub> and OTA contamination in Brazilian beer is not geographically limited and that beer does not contribute significantly to FB<sub>1</sub> intake by consumers. In the case of regular high ingestion, beer could contribute sizably to OTA, intake although still below the maximum considered tolerable for the toxin.

**Keywords:** mycotoxins; fumonisin B<sub>1</sub>; ochratoxin A; beer.

### Resumo

A presença de fumonisina B<sub>1</sub> (FB<sub>1</sub>) e de ocratoxina A (OTA) foi investigada em amostras de cerveja fabricada no Brasil. FB<sub>1</sub> foi pesquisada em 58 amostras de cerveja provenientes de 30 fábricas localizadas em nove Estados. As amostras foram concentradas, e a toxina isolada através de coluna de troca iônica forte, derivação com OPA e análise por CLAE com fluorescência associada. O limite de detecção foi 0,26 ng.mL<sup>-1</sup> e a recuperação média foi de 98%. Vinte e cinco amostras continham FB<sub>1</sub> em concentrações de 1 a 40 ng.mL<sup>-1</sup>. Cerveja (123 amostras) proveniente de 36 fábricas localizadas em 5 Estados foi analisada para OTA através de coluna de imunoafinidade seguida de CLAE com detector de fluorescência. O limite de detecção foi 0,1 ng.mL<sup>-1</sup> e a recuperação média foi de 92%. Cinco amostras continham OTA em concentrações de 1 a 18 ng.mL<sup>-1</sup>. Os resultados indicam que a contaminação da cerveja brasileira por FB<sub>1</sub> e por OTA não é geograficamente limitada e que não contribui significativamente para a ingestão de FB<sub>1</sub> por consumidores. Por outro lado, no caso de consumo alto e regular, esta pode contribuir substancialmente na ingestão de OTA, porém ainda abaixo do máximo considerado tolerável para a toxina.

**Palavras-chave:** micotoxinas; fumonisina B<sub>1</sub>; ocratoxina A; cerveja.

## 1 Introduction

Fumonisin consists of a group of mycotoxins produced mainly by strains of *Fusarium verticilloides* and *Fusarium proliferatum*<sup>5</sup>. Within this group of toxins, Fumonisin B<sub>1</sub> (FB<sub>1</sub>), B<sub>2</sub>, and B<sub>3</sub> are the most commonly found in food and feed. FB<sub>1</sub> represents up to 70% of the fumonisins produced in laboratory conditions or found in naturally contaminated food or feed. It is also the most toxic of the group<sup>9</sup>. In rats, FB<sub>1</sub> causes cancer in liver, is toxic to kidneys and liver and weakens the immune system<sup>22</sup>. FB<sub>1</sub> brings about leukoencephalomalacia in horses<sup>22,49</sup> and pulmonary edema in swine<sup>11</sup>. There is no direct evidence of FB<sub>1</sub> carcinogenic effect in humans, but an increase of oesophageal cancer is observed in areas where maize, either contaminated by *Fusarium* or containing high levels of FB<sub>1</sub>, is used as the main staple<sup>5,50</sup>. Fumonisin, mainly FB<sub>1</sub>, have been detected in a large array of maize products in all areas of the world where analyses of such commodities have been conducted<sup>4,6,7,16,17,25,40,45</sup>.

Fumonisin may be introduced when making beer when maize is used as an adjunct to fermentation<sup>18</sup>. Corn starch and corn syrup are among the adjuncts alternatively used for beer production, but no contamination by fumonisins has

been found in these materials<sup>4,28</sup>. FB<sub>1</sub> is stable to heat<sup>1,19,20</sup> and to conditions found during fermentation of corn to produce ethanol<sup>37</sup>.

The presence of fumonisins in beer was described for the first time by SCOTT and LAWRENCE<sup>36</sup> in beverages commercialized in Canada. The levels found were low and only 4 of the 41 samples analyzed had concentrations higher than 2 ng.mL<sup>-1</sup>. In a subsequent study, SCOTT et al.<sup>38</sup> found the toxins in 20 of 46 samples of beer. The concentrations of FB<sub>1</sub> + FB<sub>2</sub> in the positive samples ranged from 0.2 to 64 ng.mL<sup>-1</sup> (average = 5.6 ± 13.6 ng.mL<sup>-1</sup>). Both studies included foreign and Canadian beers. TORRES et al.<sup>51</sup> analyzed 32 samples of Spanish beers and found that 14 samples contained FB<sub>1</sub> in concentrations ranging from 4.76 to 85.53 ng.mL<sup>-1</sup> (average = 28 ± 22 ng.mL<sup>-1</sup>). HLYWKA and BULLERMAN<sup>18</sup> examined 29 samples of beers commercialized in the U.S.A. The authors detected FB<sub>1</sub>, FB<sub>2</sub>, and FB<sub>3</sub> in 25 of the samples. The contaminated samples contained FB<sub>1</sub> + FB<sub>2</sub> concentrations from 0.3 to 13.5 ng.mL<sup>-1</sup>.

Ochratoxin A (OTA) is nephrotoxic to monogastric animals, especially swine, and a potent renal carcinogen in rats. It behaves as immunotoxic and teratogenic agent. It shows genotoxic properties in some tests, but not in others such as the Ames test<sup>23,30</sup>. Evidence gathered so far, although not conclusive, points to OTA as the possible causative agent of the endemic chronic renal disorder observed in humans in the Balkan countries<sup>44,46</sup>. The occurrence of OTA in food has been shown to be worldwide, especially in cereals<sup>24,31,43</sup>. *Penicillium ver-*

Recebido para publicação em 21/6/2006

Aceito para publicação em 23/4/2007 (001768)

<sup>1</sup> Departamento de Ciência de Alimentos, Faculdade de Engenharia de Alimentos, Universidade Estadual de Campinas – UNICAMP, CP 6121, CEP 13083-682, Campinas - SP, Brasil, E-mail: valente@fea.unicamp.br

\*A quem a correspondência deve ser enviada

*rucosum*, *Aspergillus ochraceus*, and *A. carbonarius* are OTA major producers, but their individual temperature and other growth preferences cause them to occur in different areas of the globe, as well as infecting diverse food<sup>29,48</sup>. OTA residues may occur in beer, if the toxin is present in barley and malt used in its making<sup>2</sup>. The fermentation of wort to which OTA had previously been added showed that the toxin decreased from 2 to 13%<sup>37</sup>. A remainder of 13 to 32% of the toxin was found in the final product when OTA contaminated malt was obtained from barley inoculated with *Penicillium verrucosum* and subsequently used for beer making<sup>2</sup>.

Beer has been surveyed for OTA in various countries. MEDINA et al.<sup>26</sup> examined 88 samples of beer (domestic and imported) marketed in Spain and OTA was detected in 82.9% of them. The range for positive samples was 0.007-0.204 ng.mL<sup>-1</sup>. In Hungary, VARGA et al.<sup>52</sup> analyzed 25 samples of beer and all but one of the samples were found to be contaminated with small amounts of OTA with a mean concentration of 0.127 ng.mL<sup>-1</sup> and a range from 0.030 to 0.250 ng.mL<sup>-1</sup>. DASKO et al.<sup>12</sup> analyzed OTA 20 samples of local and 4 samples of foreign beer commercialized in Slovenia. No positive results for OTA presence were detected in any of the beer samples. German beers (35 samples) were analyzed by DEGELMANN et al.<sup>13</sup> and OTA levels in the range of 0.1-0.2 ng.mL<sup>-1</sup> were found in 9 beer samples, 21 samples contained trace amounts of OTA (<0.1 ng.mL<sup>-1</sup>) and in 5 samples no OTA was detectable. SCOTT and KANHERE<sup>35</sup> found traces of OTA in 26 out of 46 samples of Canadian and imported beers. In Denmark, JORGENSEN<sup>21</sup> found OTA in 21 samples of beer analyzed in concentrations ranging from 0.0001 to 0.160 ng.mL<sup>-1</sup>, with a mean of 0.049 ng.mL<sup>-1</sup>. In Japan, 94 imported and 22 Japanese made beers were tested for OTA by NAKAJIMA et al.<sup>27</sup> and a mean level of 0.010 ng.mL<sup>-1</sup> was determined in 86 out of the imported beers and a mean level of 0.0125 ng.mL<sup>-1</sup> in 21 of the Japanese beers. VISCONTI et al.<sup>53</sup> analyzed Italian and imported beers (61 samples) and found that half of the samples contained OTA in concentrations from <0.01 to 0.135 ng.mL<sup>-1</sup>. No correlation between the incidences of OTA and country or beverage type was found by SOLEAS et al.<sup>42</sup> in beer commercialized in Canada (107 samples). In Belgium, OTA was found by TANGNI et al.<sup>47</sup> in all samples of 62 Belgian and 20 imported beers analyzed and the highest concentration was 0.185 ng.mL<sup>-1</sup>. PRADO et al.<sup>31</sup> investigated the presence of OTA in 26 samples of beer produced in Brazil and found traces of the toxin in 6 samples.

Brazil is the fifth world largest producer of beer after China, US, Germany, and Russia with a yearly production of 8.5 billion liters according to data collected from 2002 to 2003. The annual per capita beer intake of 47.6 liters in 2004, on the other hand, ranks ninth after countries such as the Czech Republic, Germany, the UK, Australia, the US, Spain, Japan, and Mexico<sup>41</sup>. So far Brazilian beers have not been examined for FB<sub>1</sub> and a limited number of samples have been analyzed for OTA. The present work investigated beers produced in various locations within the country for the possible presence of FB<sub>1</sub> and OTA.

## 2 Material and methods

### 2.1 Samples

Beers made in Brazil (44 Pilsner and other light colored types and 14 stout and other dark colored types) were acquired at food markets and liquor outlets during 2000 and 2001 to be analyzed for FB<sub>1</sub>. One hundred and twenty-three samples of beer (94 light and 29 dark types) were acquired at food markets during 2003 and 2004 to be analyzed for OTA. The beers were either canned or glass bottled. A sample from each type of beer formulation was acquired whenever more than one type of beer was produced by the plant in order to cover its whole range of products.

### 2.2 Analytical reagents

Reagent grade solvents and salts were used for sample extraction, cleanup and derivatization. Ultra-pure water (Milli-Q Plus, Millipore, Milford, USA) and chromatographic grade acetonitrile and methanol were used for the mobile phase. Fumonisin B<sub>1</sub> and ochratoxin A were obtained from Sigma (St Louis, USA).

### 2.3 Sample extraction and cleanup for fumonisin B<sub>1</sub> determination

The AOAC Method 995.15<sup>34</sup> originally developed for corn and its products and modified by CAMARGOS et al.<sup>8</sup> was used as follows. In short, the pH of beer samples was brought to the 5.8-6.5 range with 1N NaOH and the samples were filtered through qualitative filter paper. For sample cleanup and concentration, an aliquot of 50 mL beer was applied to a strong anion exchange SPE column (10 cm<sup>3</sup>.500 mg<sup>-1</sup>, SAX, Bond Elut LRC, Varian, Walnut Creek, USA) previously conditioned with 10 mL methanol followed by 10 mL de methanol/water (3:1). The sample was followed by 10 mL methanol/water (3:1) and 6 mL methanol. FB<sub>1</sub> was eluted with 20 mL methanol/acetic acid (95:5). The elution was dried under nitrogen stream in a 60 °C water bath.

### 2.4 Fumonisin B<sub>1</sub> determination by liquid chromatography

The dried extract was suspended in 500 µL acetonitrile/water (1:1), an aliquot of 100 µL was transferred to a reaction vessel and 200 µL OPA reagent (40 mg o-ftaldialdehyde in 1 mL ethanol diluted with 0.1 M borate buffer and 50 µL 2-mercaptoethanol) was added. The flask was kept in an ultrasonic bath at 5-15 °C for 30 seconds. After 60 a second reaction time the derivatized sample was injected into a Model 1050 liquid chromatograph (Hewlett Packard, Palo Alto, USA), Rheodyne manual injector with a 20 µL loop, HP 1046 fluorescence detector (335 nm and 440 nm), HP 3393A integrator, Spherisorb ODS-2, 5 µm, 250 x 4.6 mm column mm (Supelco, Bellefonte, USA), guard column Varian (Walnut Creek, USA), 2 cm, filled with C18 ODS, 32 µm (Alltech, Deerfield, USA) and mobile phase water/acetonitrile/acetic acid (54:46:1) at a flow rate of 1.0 mL/minute. Identification was based on a comparison of

standard and sample retention times, and external standards were used for quantification.

### 2.5 Sample extraction and cleanup for ochratoxin A determination

Ochratoxin A was extracted from beer samples according to the method from R-Biopharm Rhone<sup>33</sup> modified as follows. The beer samples were degassed by mixing with a magnetic stirrer at 100 rpm for 60 minutes. The pH was adjusted to 7.2 with 2 M NaOH. An aliquot of 30 mL was applied to the immunoaffinity column (Ochraprep, R-Biopharm Rhone, Glasgow, Scotland). The column was subsequently washed with 20 mL ultrapure water at the flow rate of 5 mL/minute (Mili-Q Plus, Millipore, Milford, USA). The column was allowed to dry and air was flushed through it. The toxin was eluted into a vial with 3.0 mL chromatographic grade methanol and the elution was dried under nitrogen.

### 2.6 Ochratoxin A determination by liquid chromatography

The dried extract was suspended in a 1.0 mL mobile phase and 100  $\mu$ L of it was injected into an HPLC system consisting of a Rheodyne manual injector, pump system model 1050 (Hewlett Packard, Palo Alto, USA), C18 Chromolith 100 x 4.6 mm analytical column and 5 x 4.6 mm guard column (Merck, Darmstadt, Germany), fluorescence detector model 1046A (Hewlett Packard), integrator model 3393A (Hewlett Packard) under the following analytical conditions: mobile phase acetic acid/methanol (35:65) at 1 mL/minute, excitation and emission wavelengths 333 nm and 470 nm, respectively. Calibration curves were prepared in the range of 0.027 to 0.136  $\mu$ g OTA mL<sup>-1</sup>.

### 2.7 Analytical quality control


The recovery tests for FB<sub>1</sub> were conducted with samples spiked with standard to achieve concentrations of 5, 10, 20 or 50 ng.mL<sup>-1</sup>. A reagent blank and a spiked sample with 8 ng.mL<sup>-1</sup> FB<sub>1</sub> were added to each series of 8-12 samples analyzed. A sample spiked with 3 ng OTA mL<sup>-1</sup> was analyzed in triplicate and accompanied each series of 8 – 12 samples during the OTA survey. The results of these spiked samples were used to calculate recovery. All positive samples were analyzed twice, each duplicate analyzed on a different day. The detection limit for FB<sub>1</sub> was based on 3 times the average SD response for 3 injections from 8 different contaminated samples containing the toxin standard at concentrations close to 5 ng.mL<sup>-1</sup>. The detection limit for OTA was taken as 3 times the SD derived from the area of 7 injections of a spiked beer sample (3 ng.mL<sup>-1</sup>). The quantification limit was taken as 5 times the detection limit in both cases.

## 3 Results and discussion

The average recovery was 98  $\pm$  17% (n = 12) for FB<sub>1</sub> concentrations in beer between 5 and 50 ng.mL<sup>-1</sup> (Table 1). No interfering compound was observed in the chromatograms (Figure 1). The detection limit for FB<sub>1</sub> in beer samples was 0.26 ng.mL<sup>-1</sup>.

**Table 1.** Recovery of fumonisin B<sub>1</sub> (FB<sub>1</sub>) added to beer.

FB <sub>1</sub> added (ng.mL <sup>-1</sup> )	Recovery (%)
5	118.3
5	121.2
5	78.0
10	74.7
10	105.9
10	71.6
20	99.0
20	90.1
20	100.6
50	87.5
50	104.8
50	120.4



**Figure 1.** a) Chromatogram of fumonisin B<sub>1</sub> standard (4 ng.mL<sup>-1</sup>); b) chromatogram of uncontaminated beer sample; and c) chromatogram of naturally contaminated beer sample (14.3 ng FB<sub>1</sub> mL<sup>-1</sup>).

The beers used in the FB<sub>1</sub> survey were produced at 30 plants placed at 23 locations mostly located in the south-east (states of São Paulo, Minas Gerais, and Rio de Janeiro, 45 samples), and a few in the south (states of Paraná and Rio Grande do Sul, 6 samples), the north-east (states of Sergipe, Pernambuco, and Paraíba, 6 samples), and the north (state of Pará, 1 sample). The south-eastern region of the country besides being the most industrialized also concentrates 43% of the population.

FB<sub>1</sub> was detected in 25 (21 light and 4 dark) of the 58 samples of beer analyzed (43.1%) in concentrations ranging from 1 to 40 ng.mL<sup>-1</sup> (Table 2). The average concentration FB<sub>1</sub> for contaminated samples was  $9.6 \pm 10.2$  ng.mL<sup>-1</sup>. The contaminated samples were produced in plants at 16 locations, of which 11 were located in the southeast, 3 in the northeast, and 2 in the south. The distribution of the contaminated samples shows that the contamination is not confined to a single area of the country and it is probably connected with the use of maize as an adjunct. The levels found are within the range and the magnitude found in other countries where toxin has been searched for in beer<sup>18,38,51</sup>.

**Table 2.** Concentration of fumonisin B<sub>1</sub> in positive samples of commercial beer produced in Brazil during 2000 and 2001<sup>a</sup>.

Sample no.	FB <sub>1</sub> conc. <sup>b</sup> (ng.mL <sup>-1</sup> ) $\pm$ SD
1	4.5 $\pm$ 0.3
2	15 $\pm$ 1
3	14 $\pm$ 1
6	7.3 $\pm$ 0.5
8	14.7 $\pm$ 0.2
14	2.9 $\pm$ 0.3
15	39 $\pm$ 2
25	8 $\pm$ 2
27	17 $\pm$ 1
28	11 $\pm$ 2
29	3.0 $\pm$ 0.6
30	3.0 $\pm$ 0.7
34	10.7 $\pm$ 0.1
36	1.2 $\pm$ 0.2
37	12 $\pm$ 1
38	40 $\pm$ 4
39	2.3 $\pm$ 0.3
40	2.2 $\pm$ 0.4
42	10 $\pm$ 1
43	1.4 $\pm$ 0.2
45	4 $\pm$ 1
46	3.0 $\pm$ 0.3
51	4.3 $\pm$ 0.6
54	4.3 $\pm$ 0.3
56	4.0 $\pm$ 0.2

<sup>a</sup>Total number of samples analysed = 58; and <sup>b</sup>Mean of duplicate results.

The average per capita intake of beer in Brazil is 130 mL.day<sup>-1</sup><sup>40</sup>. The Provisional Maximum Tolerable Daily Intake (PMTDI) by the Joint FAO/WHO Expert Committee on Food Additives<sup>5</sup> is 2000 ng.kg<sup>-1</sup> bw for fumonisins B<sub>1</sub>, B<sub>2</sub>, and B<sub>3</sub>, alone or in group. In the present survey, the average

concentration of FB<sub>1</sub> in beer was 4.1 ng.mL<sup>-1</sup>. The intake of FB<sub>1</sub> due to beer of a consumer who ingested 1300 mL of the beverage daily (10 times the national average) and that had a body weight of 70 kg would be 3.8% of the PMTDI.

The average recovery of OTA for a concentration of 3 ng.mL<sup>-1</sup> of the toxin was 93% (n = 16) for beers of light color such as Pilsner and 87% (n = 6) for dark beers such as stout and Malzbier (Table 3). The detection and quantification limits were 0.1 and 0.5 ng.mL<sup>-1</sup>, respectively, for light and dark beers. The RSD was 1.5% for the recovery tests (n = 18) for light beers and 1.2% (n = 6) for dark beers. No interfering compound was observed in the chromatograms (Figure 2).

**Table 3.** Recovery of ochratoxin A added to beer<sup>a</sup>.


Beer type	Recovery (%) $\pm$ SD <sup>b</sup>
Pilsner	79.7 $\pm$ 5.7
	107.3 $\pm$ 5.5
	82.3 $\pm$ 5.1
	101.3 $\pm$ 1.1
	79.3 $\pm$ 7.1
	93.0 $\pm$ 2.6
	95.3 $\pm$ 6.8
	93.3 $\pm$ 5.9
	108.7 $\pm$ 3.0
	101.7 $\pm$ 8.6
	89.3 $\pm$ 2.5
	92.0 $\pm$ 1.7
	89.7 $\pm$ 2.1
	98.7 $\pm$ 1.5
95.3 $\pm$ 6.4	
95.0 $\pm$ 4.0	
83.0 $\pm$ 14.0	
89.0 $\pm$ 8.5	
Dark	85.0 $\pm$ 2.6
	65.3 $\pm$ 4.1
	114.0 $\pm$ 2.6
	62.0 $\pm$ 3.6
	88.3 $\pm$ 0.6
	109.0 $\pm$ 8.2

<sup>a</sup>Uncontaminated beer samples spiked with OTA (3 ng.mL<sup>-1</sup>); and <sup>b</sup>Mean of triplicate results.

Beers used in the survey for OTA were produced at 36 plants corresponding to 15 locations in the states of Paraná (southern region, 115 samples), São Paulo, Rio de Janeiro (south-eastern region, 3 samples), Paraíba (north-eastern region, 2 samples), and Pará (northern region, 3 samples). OTA was found in 5 samples from the 123 beers analyzed. All were light colored beers, with OTA concentrations varying from 1 to 18 ng.mL<sup>-1</sup> (Table 4). Four of the samples were from 3 plants belonging to the same beer company. One of the samples was from a plant located in the south of the country, 3 from the south-east, and one from the north. Although the number of contaminated samples is small, it shows the contamination is not restricted in terms of region.

Incidences of 83 to 100% of OTA in beer have been reported in all countries where the toxin has been searched for, except for Slovenia where no contamination has been found by OTA





**Figure 2.** a) Chromatogram of ochratoxin A standard ( $0.091 \text{ ng mL}^{-1}$ ); b) chromatogram of uncontaminated beer sample; and c) chromatogram of naturally contaminated beer sample ( $7 \text{ ng OTA mL}^{-1}$ ).

**Table 4.** Concentration of ochratoxin A in positive samples of commercial beer produced in Brazil during 2003 and 2004<sup>a</sup>.

Sample n <sup>o</sup>	OTA conc. <sup>b</sup> ( $\text{ng mL}^{-1}$ ) $\pm$ SD
48	$18.0 \pm 0.08$
85	$7.56 \pm 0.01$
113	$11.66 \pm 0.004$
115	$1.03 \pm 0.03$
116	$3.52 \pm 0.20$

<sup>a</sup>Total number of samples analysed = 123; and <sup>b</sup>Mean of duplicate results.

in beer. The incidence found in Brazilian beer in the present survey (4.8%) is therefore low. The levels found, on the other hand are higher than the levels reported by other countries by various orders of magnitude<sup>12,13,21,26,27,35,47,52,53</sup>.

The Provisional Tolerable Weekly Intake (PTWI) by JECFA<sup>3</sup> for OTA is  $100 \text{ ng kg}^{-1} \text{ bw}$ . Considering that the average contamination of beer by OTA found in the present work was  $0.3 \text{ ng mL}^{-1}$ , a consumer of 1,300 mL beer per day (10 times the national average) would be ingesting 2,730 ng OTA weekly and that would amount to 39 % of the PTWI. The incidence of OTA in beer was low but its high toxicity turns it into a significant contributor at an OTA intake in case of regular high ingestion of the beverage.

## 4 Conclusions

The survey conducted in Brazilian beer found  $\text{FB}_1$  and OTA to be present in 43% and 4.8% of the samples, respectively. The toxicity of  $\text{FB}_1$  combined with the low levels present in the samples examined, indicate that the presence of this toxin in Brazilian beer is more of a quality issue for the industry than a reason for public health concern. On the other hand, the high levels of OTA found and its high toxicity are cause for public health concern even with the low incidence observed. The elimination of both toxins from the product should be contemplated by the industry either for quality or health reasons.

## Acknowledgements

The present work was supported by a research grant from FAPESP (Fundação de Amparo à Pesquisa do Estado de São Paulo). Graduate scholarships were granted by CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico) and are gratefully acknowledged by the first two authors.

## References

- ALBERTS, J. F. et al. Effects of temperature and incubation period on production of fumonisin B<sub>1</sub> by *Fusarium moniliforme*. **Appl. Environ. Microbiol.**, v. 56, n. 6, p. 1729-1733, 1990.
- BAXTER, D.; SLAIDING, I. R.; KELLY, B. Behavior of ochratoxin A in brewing. **J. Am. Soc. Brewing Chem.**, v. 59, n. 3, p. 98-100, 2001.
- BENFORD, D. et al. Ochratoxin A. In: WHO FOOD ADDITIVES SERIES N. 47, **Safety evaluation of certain mycotoxins in food**. Geneva: World Health Organization, 2001. p. 281-415.
- BENNETT, G. A. et al. Distribution of fumonisins in food and feed products prepared from contaminated corn. In: JACKSON, L. S.; DE VRIES, J. W.; BULLERMAN, L. B.; (eds.) **Fumonisin in food**. New York: Plenum, 1996. p. 317-322.
- BOLGER, M. et al. Fumonisin. In: WHO FOOD ADDITIVES SERIES N. 47, **Safety evaluation of certain mycotoxins in food**. Geneva: World Health Organization, 2001. p. 103-279.
- BULLERMAN, L. B. Occurrence of *Fusarium* and fumonisins on food grains and in foods. In: JACKSON, L. S.; DE VRIES, J. W.; BULLERMAN, L. B. (eds.) **Fumonisin in food**. New York: Plenum, 1996. p. 27-38.
- BULLERMAN, L. B.; TSAI, W. J. Incidence and levels of *Fusarium moniliforme*, *Fusarium proliferatum* and fumonisins in corn

- and corn-based foods and feeds. **J. Food Protection**, v. 57, n. 6, p. 541-546, 1994.
8. CAMARGOS, S. M.; MACHINSKI JR, M.; VALENTE SOARES, L. M. Avaliação de métodos para determinação de fumonisinas B<sub>1</sub> e B<sub>2</sub> em milho. **Rev. Inst. Adolfo Lutz**, v. 58, n. 2, p. 71-79, 1999.
  9. CAWOOD, M. E. et al. Isolation of the fumonisin mycotoxins – a quantitative approach. **J. Agric. Food Chem.**, v. 39, n. 11, p. 1958-1962, 1991.
  10. CHELKOWSKI, J.; LEW, H. *Fusarium* species of Liseola section occurrence in cereals and ability to produce fumonisins. **Microbiol. Alim. Nutr.**, v. 10, n. 1/2, p. 49-53, 1992.
  11. COLVIN, B. M.; HARRISON, L. R. Fumonisin-induced pulmonary edema and hydrothorax in swine. **Mycopathologia**, v. 117, n. 1/2, p. 79-82, 1992.
  12. DASKO, L. et al. Determination of ochratoxin A in beer. **Czech J. Food Sci.**, v. 23, n. 2, p. 69-73, 2005.
  13. DEGELMANN, P. et al. Determination of ochratoxin A in beer by high-performance liquid chromatography. **Chromatographia**, v. 49, n. 9/10, p. 543-546, 1999.
  14. DOKO, M. B.; VISCONTI, A. Occurrence of fumonisins B<sub>1</sub> and B<sub>2</sub> in corn and corn-based human foodstuffs in Italy. **Food Add. Contam.**, v. 11, n. 4, p. 433-439, 1994.
  15. GELDERBLOM, W. C. A. et al. Toxicity and carcinogenicity of *Fusarium moniliforme* metabolite, fumonisin B<sub>1</sub>, in rats. **Carcinogenesis**, v. 12, n. 7, p. 1247-1251, 1991.
  16. HENNINGEN, M. R. et al. Fumonisin levels in commercial corn products in Buenos Aires, Argentina. **Food Add. Contam.**, v. 17, n. 1, p. 55-58, 2000a.
  17. HENNINGEN, M. R. et al. Fumonisin in corn hybrids grown in Argentina for two consecutive seasons. In: DE KOE, W. J.; SAMSON, R. A.; VAN EGMOND, H. P.; GILBERT, J.; SABINO, M. (eds.) **Mycotoxins and phycotoxins in perspective at the turn of the millennium**. Wageningen: Posen and Looten, 2000b. p. 331-339.
  18. HLYWKA, J. J.; BULLERMAN, L. B. Occurrence of fumonisin B<sub>1</sub> and B<sub>2</sub> in beer. **Food Add. Contam.**, v. 16, n. 8, p. 319-324, 1999.
  19. JACKSON, L. S. et al. Effects of time, temperature and pH on the stability of fumonisins B<sub>1</sub> in an aqueous model system. **J. Agric. Food Chem.**, v. 44, n. 4, p. 906-912, 1996a.
  20. JACKSON, L. S. et al. Effects of thermal processing on the stability of fumonisins B<sub>2</sub> in an aqueous model system. **J. Agric. Food Chem.**, v. 44, n. 10, p. 1984-1987, 1996b.
  21. JORGENSEN, K. Survey of pork, poultry, coffee, beer and pulses for ochratoxin A. **Food Add. Contam.**, v. 15, n. 4, p. 550-554, 1998.
  22. KELLERMAN, T. S. et al. Leukoencephalomalacia in two horses induced by oral dosing of fumonisin B<sub>1</sub>. **J. Vet. Res.**, v. 57, n. 4, p. 269-275, 1990.
  23. KUIPER-GOODMAN, T. Risk assessment of ochratoxin A: an update. **Food Add. Contam.**, v. 13 (Supplement), p. 53-57, 1996.
  24. KUIPER-GOODMAN, T.; SCOTT, P.M. Risk assessment of the mycotoxin Ochratoxin A. **Biomed. Environ. Sci.**, v. 2, n. 1, p. 179-248, 1989.
  25. MACHINSKI JR., M.; VALENTE SOARES, L.M. Fumonisin B<sub>1</sub> and B<sub>2</sub> in Brazilian corn-based food products. **Food Add. Contam.**, v. 17, n. 10, 875-879, 2000.
  26. MEDINA, A. et al. Determination of ochratoxin A in beer marketed in Spain by liquid chromatography with fluorescence detection using lead hydroxyacetate as a cleanup agent. **J. Chromatogr. A.**, v. 1083, n. 1/2, p. 7-13, 2005.
  27. NAKAJIMA, M.; TSOUBOUCHI, H.; MIYASABE, M. A survey of ochratoxin A and aflatoxins in domestic and imported beers in Japan by immunoaffinity and liquid chromatography. **J. AOAC Int.**, v. 82, n. 6, p. 897-902, 1999.
  28. PATEL, S. et al. Surveillance of fumonisins in UK maize based foods and other cereals. **Food Add. Contam.**, v. 40, n. 11, p. 1352-1354, 1997.
  29. PITT, J. I. *Penicillium viridicatum*, *Penicillium verrucosum*, and production of ochratoxin A. **Appl. Environ. Microbiol.**, v. 53, n. 2, p. 266-269, 1987.
  30. PLESTINA, R. Nephrotoxicity of ochratoxin A. **Food Add. Contam.**, v. 13, (Supplement), p. 49-50, 1996.
  31. POHLAND, A. E.; NESHEIM, S.; FRIEDMAN, L. Ochratoxin A: a review. **Pure Appl. Chem.**, v. 64, n. 9, p. 1029-1046, 1992.
  32. PRADO, G. et al. Ochratoxin A determination in beer by immunoaffinity column clean-up and high-performance liquid chromatography. **Ciênc. Tecnol. Alim.**, v. 23, suplemento, p. 58-61, 2003.
  33. R-BIOPHARM RHÔNE LTD. **Quantitative detection of ochratoxin A**. Glasgow, Scotland, 2002, 5 p.
  34. SCOTT, P.M. Natural toxins. In: CUNNIF, P. (ed.). **Official Methods of Analysis of the AOAC International**. Gaithersburg: AOAC International, 1997.
  35. SCOTT, P.M.; KANHERE, S.R. Determination of ochratoxin A in beer. **Food Add. Contam.**, v. 12, n. 4, p. 591-598, 1995.
  36. SCOTT, P. M.; LAWRENCE, G. A. Analysis of beer for fumonisins. **J. Food Protec.**, v. 58, n. 12, p. 1379-1382, 1995.
  37. SCOTT, P. M. et al. Fermentation of wort containing added ochratoxin A and fumonisins B<sub>1</sub> and B<sub>2</sub>. **Food Add. Contam.**, v. 12, n. 1, p. 31-40, 1995.
  38. SCOTT, P. M. et al. Evaluation of enzyme-linked immunosorbent assay for analysis of beer for fumonisins. **Food Add. Contam.**, v. 14, n. 5, p. 445-450, 1997.
  39. SHEPHARD, G. S. et al. Quantitative determination of fumonisin B<sub>1</sub> and B<sub>2</sub> by high performance liquid chromatography with fluorescence detection. **J. Liquid Chrom.**, v. 13, n. 10, p. 2077-2087, 1990.
  40. SHEPHARD, G. S. et al. Worldwide survey of fumonisin contamination of corn and corn-based products. **J. AOAC Int.**, v. 79, n. 3, p. 671-687, 1996.
  41. SINDICATO NACIONAL DE INDÚSTRIAS DA CERVEJA. Dados de consumo. São Paulo, 2006. Disponível em: <<http://www.sindicerv.com.br>>. Acesso em: 15 maio 2006.
  42. SOLEAS, G. J.; YAN, J.; GOLDBERG, D. M. Assay of ochratoxin A in wine and beer by high-pressure liquid chromatography photodiode array and gas chromatography mass selective detection. **J. Agric. Food Chem.**, v. 49, n. 11, p. 2733-2740, 2001.
  43. SPEIJERS, J. A.; VAN EGMOND, H. P. Worldwide ochratoxin A levels in food and feeds. In: CREPPY, E. E.; CASTEGNARO, M.; DIRHEIMER, G. (eds.) **Human ochratoxicosis and its pathologies**. Paris: Libbey Eurotext, 1993. pp. 85-100.
  44. STOEV, S.D. The role of ochratoxin A as a possible cause of Balkan endemic nephropathy and its risk evaluation. **Vet. Human Toxicol.**, v. 40, n. 4, p. 352-360, 1998.

45. SYDENHAM, E. W. et al. Fumonisin contamination of commercial corn-based human foodstuffs. **J. Agric. Food Chem.**, v. 39, n. 9, p. 2014-2018, 1991.
46. TATU, C. A. et al. The etiology of Balkan endemic nephropathy: still more questions than answers. **Environ. Health Perspec.**, v. 106, n. 7, p. 689-700, 1998.
47. TANGNI, E. K. et al. Ochratoxin A in domestic and imported beers in Belgium: occurrence and exposure assessment. **Food Add. Contam.**, v. 19, n. 12, p. 1169-1179, 2002.
48. TEREM, J. et al. Immunochemical detection of ochratoxin A in black *Aspergillus* strains. **Mycopathol.**, v. 134, n. 1, p. 171-176, 1996.
49. THIEL, P. G. et al. The implications of naturally occurring levels of fumonisins in corn for human and animal health. **Mycopathol.**, v. 117, n. 1, p. 3-9, 1992.
50. THIEL, P. G. et al. Levels of fumonisins B<sub>1</sub> and B<sub>2</sub> in feeds associated with confirmed cases of equine leukoencephalomalacia. **J. Agric. Food Chem.**, v. 39, n. 1, p. 109-111, 1991.
51. TORRES, M. R.; SANCHIS, V.; RAMOS, A. J. Occurrence of fumonisins in Spanish beers analyzed by an enzyme-linked immunosorbent assay method. **Int. J. Food Microbiol.**, v. 39, n. 1/2, p. 139-143, 1998.
52. VARGA, J. et al. Detection of ochratoxin A in Hungarian wines and beers. **Acta Aliment.**, v. 34, n. 4, p. 381-392, 2005.
53. VISCONTI, A.; DOKO, M. B. Survey of fumonisin production by *Fusarium* isolated from cereals in Europe. **J. AOAC Int.**, v. 77, n. 6, p. 546-550, 1994.