

## Original

**Anthropometry of height, weight, arm, wrist, abdominal circumference and body mass index, for Bolivian Adolescents 12 to 18 years - Bolivian adolescent percentile values from the MESA study**A. Baya Botti<sup>1,5</sup>, F. J. A. Pérez-Cueto<sup>1,2</sup>, P. A. Vasquez Monllor<sup>3,5</sup> and P. W. Kolsteren<sup>1,4</sup><sup>1</sup>Nutrition and Child Health Unit. Department of Public Health. Prince Leopold Institute of Tropical Medicine. Antwerp. Belgium.<sup>2</sup>Department of Agricultural Economics. Faculty of Bioscience Engineering. Ghent University. Ghent. Belgium.<sup>3</sup>Faculdade de Engenharia dos Alimentos. Universidade de Campinas. Campinas. SP. Brasil.<sup>4</sup>Department of Food Safety and Food Quality. Faculty of Bioscience Engineering. Ghent. Belgium.<sup>5</sup>Unidad de Nutrición. Dirección Nacional de Investigación. Universidad del Valle. Cochabamba. Bolivia.**Abstract**

Anthropometry is important as clinical tool for individual follow-up as well as for planning and health policy-making at population level. Recent references of Bolivian Adolescents are not available. The aim of this cross sectional study was to provide age and sex specific centile values and charts of Body Mass Index, height, weight, arm, wrist and abdominal circumference from Bolivian Adolescents. Data from the METabolic Syndrome in Adolescents (MESA) study was used. Thirty-two Bolivian clusters from urban and rural areas were selected randomly considering population proportions, 3445 school going adolescents, 12 to 18 y, 45% males; 55% females underwent anthropometric evaluation by trained personnel using standardized protocols for all interviews and examinations. Weight, height, wrist, arm and abdominal circumference data were collected. Body Mass Index was calculated. Smoothed age- and gender specific 3<sup>rd</sup>, 5<sup>th</sup>, 10<sup>th</sup>, 25<sup>th</sup>, 50<sup>th</sup>, 75<sup>th</sup>, 85<sup>th</sup>, 90<sup>th</sup>, 95<sup>th</sup> and 97<sup>th</sup> Bolivian adolescent percentiles (BAP) and Charts (BAC) were derived using LMS regression. Percentile-based reference data for the anthropometrics of for Bolivian Adolescents are presented for the first time.

(Nutr Hosp. 2009;24:304-311)

Key words: *Anthropometry. Body Mass Index (BMI). Growth percentiles. Waist circumference. Abdominal circumference. Height. Weight. Adolescents. Bolivia.*

**REFERENCIAS ANTROPOMÉTRICAS DE LOS ADOLESCENTES BOLIVIANOS DE 12 A 18 AÑOS: ESTATURA, PESO, CIRCUNFERENCIA DEL BRAZO, MUÑECA, Y ABDOMINAL, ÍNDICE DE MASA CORPORAL. PERCENTILES DE ADOLESCENTES BOLIVIANOS (PAB) DEL ESTUDIO MESA**

**Resumen**

La antropometría es una herramienta clínica importante para el seguimiento individual de los pacientes así como para la planificación de políticas públicas. En Bolivia no existen referencias antropométricas nacionales para adolescentes. El objetivo de este estudio transversal fue de desarrollar percentiles y diagramas de crecimiento para peso, talla, índice de masa corporal, presión arterial sistólica y diastólica, circunferencia de muñeca, brazo y abdominal de adolescentes bolivianos. Los datos antropométricos en el estudio MESA (Síndrome metabólico en adolescentes bolivianos) fueron obtenidos a partir de 32 unidades muestrales, considerando proporcionalidad muestral con reposición. Fueron evaluados 3445 adolescentes de 12 a 18, 45% hombres; 55% mujeres, de colegios de áreas urbanas y rurales. La evaluación fue efectuada por personal entrenado siguiendo procedimientos estandarizados. Se tomaron medidas del peso, talla circunferencias de muñeca, brazo y abdominal. El índice de masa corporal fue calculado. Se obtuvieron los valores de los percentiles 3<sup>o</sup>, 5<sup>o</sup>, 10<sup>o</sup>, 25<sup>o</sup>, 50<sup>o</sup>, 75<sup>o</sup>, 85<sup>o</sup>, 90<sup>o</sup>, 95<sup>o</sup> y 97<sup>o</sup> utilizando regresión por el método LMS. Las referencias antropométricas para los adolescentes bolivianos son presentadas por vez primera a la comunidad médica. (En la versión electrónica de Nutrición Hospitalaria se puede consultar el texto íntegro en castellano de este artículo).

(Nutr Hosp. 2009;24:304-311)

Palabras clave: *Antropometría. Índice de Masa Corporal (IMC). Percentiles de crecimiento. Circunferencia de la cintura. Circunferencia abdominal. Talla. Peso. Adolescentes. Bolivia.*

**Correspondence:** Ana Baya Botti.

Pasaje La Sevillana 1353.

P. O. Box 6557 Cochabamba - Bolivia.

E-mail: [baya.ana@gmail.com](mailto:baya.ana@gmail.com)

Recibido: 5-VI-2008.

Aceptado: 1-VII-2008.

## Introduction

Anthropometric parameters and their derived indices are frequently used by physicians and health workers as a valuable instrument to determine health and disease, to define nutritional status, to assess growth and development, to determine differences in body proportion between populations as well as to optimize diagnosis and treatment.<sup>1-3</sup>

Decisions for policy making and planning in public health nutrition must be based on anthropometric accurate information about the population for which it is intended to be used. Since little is known regarding anthropometry of Bolivian adolescents,<sup>4</sup> and no national reference percentiles or charts have been developed, international references<sup>5-7</sup> have been used systematically for growth monitoring and nutritional classification of individuals.

Previous studies in La Paz, and from other regions of Bolivia<sup>4,8-10</sup> confirmed the need for updated information to address the nutritional status of adolescents in Bolivia. The country faces nutritional transition and adolescents are among the most vulnerable group to its impact. Increased numbers of overweight and obese adolescents has been described recently.<sup>4</sup> For the Bolivian health system, having access to local growth references and clinical evaluation parameters for adolescents is urgent and crucial to measure trends in nutritional status and to develop concurrent policies.

Local anthropometric references are indispensable to perform high-quality clinical practices. Health care providers base their diagnosis on percentile values to decide extent of a problem and level of treatment. This is particularly true for predicting and assessing risk for cardiovascular diseases and metabolic syndrome, which uses among other factors, percentile values of Body Mass Index (BMI), abdominal circumference and blood pressure. For the assessment of high blood pressure percentiles of height is required for adolescents. The use of heights derived from other populations could induce to diagnosis error.

For nutritional intervention programs local population percentile values could help to portrait future risk associated to nutritional transition outcomes, and initiate activities to reduce morbidity and mortality rates associated with risk factors for chronic diseases, such as hyperlipidaemia, hyperinsulinaemia, hypertension, and early atherosclerosis in adulthood.<sup>6,11-14</sup> Treatment of diet related diseases depletes the Bolivian limited health budget resources. Therefore interventions based in early detection and correct targeting of populations at risk is likely to reduce future expenditures.

In 2005 a national study called the METabolic Syndrome in Adolescents (MESA) was carried out to assess the cardiovascular and metabolic syndrome of Bolivian adolescents in relation to obesity, diabetes, income, food intake and physical activity in Bolivia. The first component of the MESA study was to document references of anthropometric parameters needed

to measure risk. This document provide age and sex specific percentile values and charts of BMI, height, weight, arm, and wrist and waist circumference from Bolivian adolescents.

## Methods

Sample size was estimated using Epi info v 3 Software assuming a prevalence of 2.5% obesity, at 95% confidence level. A total of 32 clusters proportional to population size, with replacement from the national list of Counties were selected randomly. A cluster was defined as a school. Schools were chosen from the corresponding Education District's list of the County. Individuals were chosen from the school register. The sample size calculated for each school was 120 subjects, about 20 per grade from 7<sup>th</sup> to 12<sup>th</sup> grade. Data was collected from September 2005 to June 2007.

The random selection ensured that the Bolivian population from the Andean highlands, valleys and tropics were appropriately represented. Rural, semi-urban and urban settings were also represented in the sample. The study protocol was approved by the ethics committee of the Universidad del Valle and the Bolivian Ministry of Education. Ethical procedures comply with the Helsinki declaration of 1975 reviewed in 2000.<sup>15</sup> Informed consent was obtained from all participants, and a parent or legal guardian.

Adolescents completed a self administered questionnaire on sociodemographic, nutritional intake and physical activity aspects. A date for a school visit was scheduled for data collection.

Weight, height, wrist, arm and abdominal circumference were recorded twice for each individual by trained personnel following WHO's recommendations.<sup>1</sup> Average values were used for the analysis. Weight was recorded in light, indoor clothing with a Beurer's digital scale to the nearest 0.1 kg, height was measured without shoes to the nearest 0.1 cm using a portable metal stadiometer. Abdominal circumference was measured to the nearest 0.1 cm at the high point of the iliac crest at minimal respiration when the participant was in a standing position, using a steel measuring tape following WHO recommendations.<sup>1</sup> Pregnant adolescents were excluded.

For each participant of all locations, the same equipment was used for the anthropometric measurements. All evaluations were carried out from eight to eleven in the morning. Data quality was assured by previous extensive training of the medical assistants.

BMI was calculated applying the standard formula: Weight in kilograms divided by the square of height.

For comparability with other studies, the 3<sup>rd</sup>, 5<sup>th</sup>, 10<sup>th</sup>, 25<sup>th</sup>, 50<sup>th</sup>, 75<sup>th</sup>, 85<sup>th</sup>, 90<sup>th</sup>, 95<sup>th</sup> and 97<sup>th</sup> percentiles were chosen as reference values. Smoothed age- and gender specific values and charts for each percentile value and for each anthropometric index were derived using the least median squares (LMS) regression. The Cole's

**Table I**  
Anthropometric characteristics of the sample by age: Mean (SD)

Age	n = 3,445	Boys				Girls			
		n	Weight (kg)	Height (cm)	BMI (kg/m <sup>2</sup> )	n	Weight (kg)	Height (cm)	BMI (kg/m <sup>2</sup> )
12	342	172	43.8 (9.93)	147.6 (8.7)	19.9 (3.4)	170	44.4 (9.4)	149.1 (7.4)	19.8 (3.2)
13	435	221	47.4 (9.98)	153.7 (8.2)	19.9 (3.2)	214	47.6 (7.8)	152.1 (7.9)	20.6 (3.1)
14	486	221	52.6 (10.0)	160.9 (8.0)	20.3 (2.9)	265	51 (8.4)	153.3 (5.7)	21.7 (3.3)
15	621	267	55.0 (10.0)	163.0 (6.7)	20.7 (3.0)	354	52.7 (7.3)	154.7 (5.8)	22.0 (2.9)
16	717	306	57.5 (7.9)	165.2 (6.7)	21 (2.4)	411	54.0 (8.2)	154.3 (6.2)	22.6 (3.0)
17	561	220	58.6 (2.8)	165.5 (6.8)	21.4 (2.8)	341	54.5 (8.7)	154.9 (6.1)	22.7 (3.4)
18	283	144	60.3 (8.5)	166.6 (6.0)	21.7 (2.7)	139	56.8 (9.4)	155.4 (6.5)	23.4 (3.4)

LMS method<sup>16</sup> also called maximum penalized likelihood approach was used because it has proven to be a powerful and compact technique for deriving and presenting reference charts. It calculates the Box-Cox power needed to transform the data to normality at each age, and displaying the results as a smooth curve of power plotted against age allowing the original centiles to be reconstructed to high accuracy. The LMS Pro software used for data management was obtained from the institute of Child Health, London. Descriptive statistics were computed using SPSS v 12 and graphs and charts from LMSChartMaker 2006.

## Results

Data was collected on 3,445 adolescents, 1,551 boys and 1,894 girls, from rural (34.8%) and urban areas (65.2%), and from public (76.4%) and private (23.6%) schools. Although 4,013 adolescents were selected ini-

tially to participate on the study, 3,445 finally participated in the study. Adolescents dropped out of the study due to refusal of parental consent, failure to attend the day of data collection, failure to fill the birth date or name, failure to return the questionnaire or failure to have their anthropometric data taken or completed. Characteristics of the population sample by age are presented in table I.

Table II to VII shows smoothed percentile values respectively for BMI, height, weight, abdominal, arm and wrist circumference by age- and gender.

Figures 1 to 12 show the smoothed charts for each anthropometric parameter in order to be available for practical clinical application.

## Conclusion

Smoothed age- and gender specific 3<sup>rd</sup>, 5<sup>th</sup>, 10<sup>th</sup>, 25<sup>th</sup>, 50<sup>th</sup>, 75<sup>th</sup>, 85<sup>th</sup>, 90<sup>th</sup>, 95<sup>th</sup> and 97<sup>th</sup> Bolivian adolescent per-

**Table II**  
Body Mass Index (BMI) percentile values for Bolivian adolescents 12<sup>th</sup> to 18<sup>th</sup> years, by age and gender

Percentile	3 <sup>th</sup>	5 <sup>th</sup>	10 <sup>th</sup>	25 <sup>th</sup>	50 <sup>th</sup>	75 <sup>th</sup>	85 <sup>th</sup>	90 <sup>th</sup>	95 <sup>th</sup>	97 <sup>th</sup>
<i>Age</i>										
<i>Girls</i>										
12	14.9	15.3	16.1	17.5	19.3	21.5	22.8	23.8	25.4	26.5
13	15.9	16.4	17.1	18.5	20.3	22.4	23.8	24.7	26.3	27.5
14	16.9	17.3	18.0	19.4	21.1	23.2	24.5	25.5	27.1	28.3
15	17.6	18.0	18.7	20.0	21.7	23.8	25.1	26.1	27.7	28.9
16	18.0	18.4	19.1	20.4	22.1	24.2	25.5	26.5	28.2	29.4
17	18.3	18.7	19.4	20.6	22.4	24.5	25.9	27.0	28.8	30.1
18	18.5	18.9	19.6	20.9	22.7	24.9	26.4	27.5	29.4	30.9
<i>Boys</i>										
12	15.1	15.4	16.1	17.3	19.1	21.3	22.7	23.9	25.9	27.5
13	15.6	16.0	16.6	17.8	19.5	21.6	22.9	24.0	25.8	27.1
14	16.1	16.5	17.1	18.3	19.9	21.8	23.1	24.0	25.7	26.9
15	16.7	17.0	17.6	18.8	20.3	22.1	23.3	24.2	25.6	26.7
16	17.1	17.5	18.1	19.2	20.7	22.5	23.6	24.4	25.8	26.8
17	17.5	17.8	18.4	19.6	21.0	22.8	23.9	24.7	26.1	27.0
18	17.7	18.1	18.7	19.8	21.3	23.1	24.2	25.1	26.4	27.4

**Table III**  
*Height percentile values (m) for Bolivian adolescents 12<sup>th</sup> to 18<sup>th</sup> years by age and gender*

Percentile	3 <sup>th</sup>	5 <sup>th</sup>	10 <sup>th</sup>	25 <sup>th</sup>	50 <sup>th</sup>	75 <sup>th</sup>	85 <sup>th</sup>	90 <sup>th</sup>	95 <sup>th</sup>	97 <sup>th</sup>
<i>Age</i>										
<i>Girls</i>										
12	1.35	1.37	1.40	1.44	1.49	1.54	1.56	1.58	1.61	1.63
13	1.40	1.41	1.43	1.47	1.52	1.56	1.59	1.60	1.63	1.65
14	1.42	1.44	1.46	1.49	1.53	1.58	1.60	1.61	1.64	1.65
15	1.43	1.45	1.47	1.50	1.54	1.58	1.61	1.62	1.64	1.66
16	1.44	1.45	1.47	1.51	1.55	1.59	1.61	1.62	1.65	1.66
17	1.44	1.45	1.47	1.51	1.55	1.59	1.61	1.63	1.65	1.67
18	1.44	1.45	1.47	1.51	1.55	1.60	1.62	1.63	1.66	1.67
<i>Boys</i>										
12	1.33	1.35	1.38	1.43	1.48	1.54	1.57	1.59	1.59	1.65
13	1.40	1.41	1.44	1.49	1.55	1.60	1.63	1.65	1.65	1.70
14	1.45	1.47	1.50	1.55	1.60	1.65	1.68	1.70	1.70	1.75
15	1.50	1.51	1.54	1.58	1.63	1.68	1.71	1.72	1.72	1.77
16	1.52	1.54	1.56	1.60	1.65	1.69	1.72	1.74	1.74	1.78
17	1.54	1.55	1.58	1.62	1.66	1.70	1.73	1.74	1.74	1.78
18	1.55	1.57	1.59	1.63	1.67	1.71	1.73	1.74	1.74	1.78

**Table IV**  
*Weight percentile values (kg) for Bolivian adolescents 12<sup>th</sup> to 18<sup>th</sup> years by age and gender*

Percentile	3 <sup>th</sup>	5 <sup>th</sup>	10 <sup>th</sup>	25 <sup>th</sup>	50 <sup>th</sup>	75 <sup>th</sup>	85 <sup>th</sup>	90 <sup>th</sup>	95 <sup>th</sup>	97 <sup>th</sup>
<i>Age</i>										
<i>Girls</i>										
12	30.8	32.0	34.0	37.8	42.7	48.6	52.2	54.9	59.3	62.4
13	35.2	36.4	38.3	42.0	46.8	52.5	56.1	58.7	63.1	66.1
14	38.6	39.7	41.6	45.1	49.7	55.3	58.8	61.4	65.6	68.6
15	40.9	42.0	43.8	47.2	51.7	57.1	60.5	63.1	67.2	70.3
16	42.1	43.2	45.0	48.3	52.8	58.2	61.7	64.3	68.6	71.7
17	42.8	43.9	45.7	49.0	53.5	59.1	62.7	65.5	70.1	73.5
18	43.4	44.4	46.2	49.6	54.2	60.0	63.9	66.8	71.8	75.6
<i>Boys</i>										
12	28.7	30.0	32.3	36.6	42.2	49.1	53.3	56.5	61.6	65.3
13	33.2	34.6	36.8	41.1	46.7	53.6	57.9	61.2	66.5	70.3
14	37.6	38.9	41.1	45.3	50.8	57.5	61.7	64.9	70.2	74.0
15	41.4	42.6	44.7	48.6	53.8	60.2	64.2	67.2	72.1	75.8
16	44.4	45.6	47.6	51.3	56.1	62.0	65.7	68.5	73.1	76.4
17	46.6	47.7	49.6	53.1	57.7	63.3	66.8	69.4	73.7	76.9
18	48.4	49.5	51.3	54.6	59.0	64.3	67.6	70.1	74.2	77.3

centiles(BAP) and Charts(BAC) for the anthropometric parameters of height, weight, arm, wrist, and abdominal circumference and body mass index for Bolivian Adolescents 12 to 18y were developed.

The data of the sample distribution resembles the population distribution of the country. As stated in the 2007 projection of the 2001 National Census<sup>17,18</sup> the urban and rural population is estimated respectively at 65.4% and 34.6%. The study was carried out with school attending adolescents missing the ones that do not attend it, thus no claim can be made for complete representativeness. However there is no indication that

anthropometric values of this group could vary due to genetic factors from the adolescents that have participated in the study. This factor may on the other hand have reduced data collection from a more vulnerable adolescent population that may have suffered more from infection or disease two elements that can negatively affect malnutrition and growth.

Anthropometry provides the single most convenient, universally applicable, inexpensive and non-invasive technique for assessing size, proportions and composition of the human body. It reflects both health and nutritional status and predicts performance, health and survival.<sup>1,4</sup>

**Table V**  
Abdominal circumference percentile values (cm) for Bolivian adolescents 12<sup>th</sup> to 18<sup>th</sup> years by age and gender

Percentile	3 <sup>th</sup>	5 <sup>th</sup>	10 <sup>th</sup>	25 <sup>th</sup>	50 <sup>th</sup>	75 <sup>th</sup>	85 <sup>th</sup>	90 <sup>th</sup>	95 <sup>th</sup>	97 <sup>th</sup>
<i>Age</i>										
<i>Girls</i>										
12	57.7	58.8	60.6	64.0	68.3	73.5	76.7	79.1	83.0	85.9
13	60.0	61.1	62.9	66.3	70.7	75.8	79.1	81.4	85.3	88.1
14	61.7	62.9	64.7	68.1	72.5	77.8	81.0	83.4	87.3	90.1
15	63.0	64.1	66.0	69.4	73.9	79.2	82.4	84.9	88.8	91.7
16	63.7	64.9	66.7	70.3	74.8	80.2	83.5	86.0	90.1	93.0
17	64.0	65.2	67.1	70.7	75.4	81.0	84.5	87.0	91.3	94.4
18	64.3	65.5	67.5	71.2	76.0	81.8	85.4	88.1	92.6	95.9
<i>Boys</i>										
12	57.7	58.8	60.6	64.0	68.7	74.7	78.6	81.7	87.0	91.1
13	60.5	61.5	63.2	66.4	70.7	76.2	79.9	82.7	87.7	91.5
14	63.1	64.0	65.6	68.5	72.5	77.6	80.9	83.5	88.1	91.6
15	65.2	66.1	67.5	70.3	74.0	78.6	81.7	84.1	88.2	91.3
16	66.7	67.5	68.9	71.5	75.0	79.4	82.2	84.5	88.3	91.3
17	67.5	68.3	69.7	72.2	75.6	79.8	82.6	84.8	88.6	91.6
18	68.2	69.0	70.3	72.7	76.1	80.3	83.1	85.3	89.1	92.1

**Table VI**  
Arm circumference percentile values (cm) for Bolivian adolescents 12<sup>th</sup> to 18<sup>th</sup> years by age and gender

Percentile	3 <sup>th</sup>	5 <sup>th</sup>	10 <sup>th</sup>	25 <sup>th</sup>	50 <sup>th</sup>	75 <sup>th</sup>	85 <sup>th</sup>	90 <sup>th</sup>	95 <sup>th</sup>	97 <sup>th</sup>
<i>Age</i>										
<i>Girls</i>										
12	18.2	18.6	19.3	20.6	22.2	24.1	25.3	26.1	27.5	28.4
13	19.1	19.5	20.2	21.4	23.0	24.8	26.0	26.8	28.2	29.1
14	19.9	20.3	20.9	22.1	23.6	25.4	26.5	27.4	28.7	29.7
15	20.4	20.8	21.4	22.6	24.0	25.8	26.9	27.7	29.0	30.0
16	20.7	21.1	21.7	22.8	24.3	26.0	27.1	28.0	29.3	30.3
17	20.9	21.3	21.9	23.0	24.4	26.2	27.4	28.2	29.7	30.8
18	21.1	21.5	22.1	23.2	24.7	26.5	27.7	28.6	30.1	31.3
<i>Boys</i>										
12	17.9	18.3	19.0	20.2	21.8	23.8	25.0	26.0	27.5	28.6
13	18.7	19.2	19.8	21.1	22.7	24.6	25.7	26.6	28.0	28.9
14	19.5	19.9	20.6	21.8	23.4	25.2	26.3	27.1	28.3	29.2
15	20.1	20.5	21.2	22.4	24.0	25.7	26.7	27.4	28.6	29.4
16	20.7	21.1	21.8	23.0	24.5	26.2	27.2	27.9	29.0	29.7
17	21.1	21.5	22.2	23.5	25.1	26.7	27.7	28.4	29.5	30.2
18	21.5	22.0	22.7	24.0	25.6	27.3	28.3	29.0	30.1	30.8

For adolescents it is also useful to determine biological maturity and health risks. To the knowledge of the authors, this is the largest anthropometric survey carried out in Bolivia, providing nationally representative data. Bolivian health providers have for the first time locally developed anthropometric tables and charts at their disposal, to assess the nutritional status of Bolivian adolescents 12 to 18 y which can also be used for other clinical applications.

Several characteristics make this tool valuable and reliable: they came from a large set of data, in which all the 327 Counties of Bolivia had similar opportunity to be selected, rural and urban areas were represented at

the same level of distribution as the general population, all ages and gender of adolescents from 12 to 18<sup>th</sup> year were assessed in the same study, following a standard protocol, using the same equipment and carried out by the same research team.

It is important to mention that percentile values and charts came from a descriptive study that portrays the present situation of adolescents, and does not claim to be a standard that describes a population that followed healthy recommendations for all parameters that could affect normal growth and body composition, or a population that have developed its full genetic potential. Values and charts must be used for this reason with

**Table VII**  
Wrist percentile values (cm) for Bolivian adolescents 12<sup>th</sup> to 18<sup>th</sup> years by age and gender

Percentile	3 <sup>th</sup>	5 <sup>th</sup>	10 <sup>th</sup>	25 <sup>th</sup>	50 <sup>th</sup>	75 <sup>th</sup>	85 <sup>th</sup>	90 <sup>th</sup>	95 <sup>th</sup>	97 <sup>th</sup>
<i>Age</i>										
<i>Girls</i>										
12	13.0	13.3	13.6	14.3	15.0	15.8	16.2	16.5	17.0	17.3
13	13.4	13.6	14.0	14.6	15.3	16.0	16.4	16.7	17.1	17.4
14	13.7	13.9	14.2	14.8	15.5	16.2	16.6	16.8	17.3	17.5
15	13.9	14.1	14.4	14.9	15.6	16.2	16.6	16.9	17.3	17.6
16	13.9	14.1	14.4	14.9	15.6	16.3	16.7	17.0	17.4	17.7
17	13.9	14.1	14.4	14.9	15.6	16.3	16.8	17.1	17.5	17.8
18	13.8	14.1	14.4	15.0	15.7	16.4	16.9	17.2	17.7	18.1
<i>Boys</i>										
12	13.1	13.4	13.8	14.4	15.2	16.1	16.6	17.0	17.5	17.9
13	13.6	13.9	14.2	14.9	15.7	16.5	17.0	17.3	17.8	18.1
14	14.0	14.3	14.7	15.3	16.0	16.8	17.2	17.5	18.0	18.3
15	14.3	14.6	14.9	15.5	16.2	16.9	17.3	17.6	18.0	18.3
16	14.6	14.8	15.1	15.7	16.4	17.1	17.4	17.7	18.1	18.3
17	14.7	14.9	15.3	15.9	16.5	17.2	17.6	17.8	18.2	18.4
18	14.8	15.0	15.4	16.0	16.7	17.3	17.7	17.9	18.3	18.5




Fig. 1.—Height curves for the 3<sup>rd</sup>, 5<sup>th</sup>, 10<sup>th</sup>, 25<sup>th</sup>, 50<sup>th</sup>, 75<sup>th</sup>, 85<sup>th</sup>, 90<sup>th</sup>, 95<sup>th</sup> and 97<sup>th</sup> percentile of Bolivian girls 12<sup>th</sup>-18<sup>th</sup> years.




Fig. 3.—Weight curves for the 3<sup>rd</sup>, 5<sup>th</sup>, 10<sup>th</sup>, 25<sup>th</sup>, 50<sup>th</sup>, 75<sup>th</sup>, 85<sup>th</sup>, 90<sup>th</sup>, 95<sup>th</sup> and 97<sup>th</sup> percentile of Bolivian girls 12<sup>th</sup>-18<sup>th</sup> years.




Fig. 2.—Height curves for the 3<sup>rd</sup>, 5<sup>th</sup>, 10<sup>th</sup>, 25<sup>th</sup>, 50<sup>th</sup>, 75<sup>th</sup>, 85<sup>th</sup>, 90<sup>th</sup>, 95<sup>th</sup> and 97<sup>th</sup> percentile of Bolivian boys 12<sup>th</sup>-18<sup>th</sup> years.




Fig. 4.—Weight curves for the 3<sup>rd</sup>, 5<sup>th</sup>, 10<sup>th</sup>, 25<sup>th</sup>, 50<sup>th</sup>, 75<sup>th</sup>, 85<sup>th</sup>, 90<sup>th</sup>, 95<sup>th</sup> and 97<sup>th</sup> percentile of Bolivian boys 12<sup>th</sup>-18<sup>th</sup> years.




Fig. 5.—Body Mass Index (BMI) curves for the 3<sup>rd</sup>, 5<sup>th</sup>, 10<sup>th</sup>, 25<sup>th</sup>, 50<sup>th</sup>, 75<sup>th</sup>, 85<sup>th</sup>, 90<sup>th</sup>, 95<sup>th</sup> and 97<sup>th</sup> percentile of Bolivian girls 12<sup>th</sup>-18<sup>th</sup> years.




Fig. 8.—Abdominal circumference curves for the 3<sup>rd</sup>, 5<sup>th</sup>, 10<sup>th</sup>, 25<sup>th</sup>, 50<sup>th</sup>, 75<sup>th</sup>, 85<sup>th</sup>, 90<sup>th</sup>, 95<sup>th</sup> and 97<sup>th</sup> percentile of Bolivian girls 12<sup>th</sup>-18<sup>th</sup> years.




Fig. 6.—Body Mass Index (BMI) curves for the 3<sup>rd</sup>, 5<sup>th</sup>, 10<sup>th</sup>, 25<sup>th</sup>, 50<sup>th</sup>, 75<sup>th</sup>, 85<sup>th</sup>, 90<sup>th</sup>, 95<sup>th</sup> and 97<sup>th</sup> percentile of Bolivian boys 12<sup>th</sup>-18<sup>th</sup> years.




Fig. 9.—Arm circumference curves for the 3<sup>rd</sup>, 5<sup>th</sup>, 10<sup>th</sup>, 25<sup>th</sup>, 50<sup>th</sup>, 75<sup>th</sup>, 85<sup>th</sup>, 90<sup>th</sup>, 95<sup>th</sup> and 97<sup>th</sup> percentile of Bolivian girls 12<sup>th</sup>-18<sup>th</sup> years.




Fig. 7.—Abdominal circumference curves for the 3<sup>rd</sup>, 5<sup>th</sup>, 10<sup>th</sup>, 25<sup>th</sup>, 50<sup>th</sup>, 75<sup>th</sup>, 85<sup>th</sup>, 90<sup>th</sup>, 95<sup>th</sup> and 97<sup>th</sup> percentile of Bolivian girls 12<sup>th</sup>-18<sup>th</sup> years.



Fig. 10.—Arm circumference curves for the 3<sup>rd</sup>, 5<sup>th</sup>, 10<sup>th</sup>, 25<sup>th</sup>, 50<sup>th</sup>, 75<sup>th</sup>, 85<sup>th</sup>, 90<sup>th</sup>, 95<sup>th</sup> and 97<sup>th</sup> percentile of Bolivian boys 12<sup>th</sup>-18<sup>th</sup> years.

caution by the medical community. Once an international or a national reference that considers these aspects is developed, the BAP must be replaced by it.

Clearly there will be quantitative and qualitative dimensions to consider with the introduction of the newly developed Bolivian Adolescent Percentiles




Fig. 11.—Wrist circumference curves for the 3<sup>rd</sup>, 5<sup>th</sup>, 10<sup>th</sup>, 25<sup>th</sup>, 50<sup>th</sup>, 75<sup>th</sup>, 85<sup>th</sup>, 90<sup>th</sup>, 95<sup>th</sup> and 97<sup>th</sup> percentile of Bolivian girls 12<sup>th</sup>-18<sup>th</sup> years.




Fig. 12.—Wrist circumference curves for the 3<sup>rd</sup>, 5<sup>th</sup>, 10<sup>th</sup>, 25<sup>th</sup>, 50<sup>th</sup>, 75<sup>th</sup>, 85<sup>th</sup>, 90<sup>th</sup>, 95<sup>th</sup> and 97<sup>th</sup> percentile of Bolivian boys 12<sup>th</sup>-18<sup>th</sup> years.

(BAP) Reference in replacement of the CDC or other international available references such as the IOTF. There is a need to compare the performance of these different References in their ability to classify individuals according to their nutritional status or to diagnose the risk of chronic disease in this population.

It may be that the criteria to identify adolescents at risk of overweight or a biochemical imbalance needs to be revised when differences are observed in sensitivity and specificity between the different references for a variety of outcome parameters.

### Acknowledgments

The authors wish to thank the adolescents, their parents, and school directors from the 32 locations where the studies were conducted. Also to students and the staff of the Faculty of Medicine of the Universidad del Valle as well as the personnel of other institutions for their cooperation during different steps of the study. Special thanks go to the Prince Leopold Institute of Tropical Medicine who has supported this study and to Nutrition Third World for making it possible through funding it.

### References

1. WHO. Physical status : The use and interpretation of anthropometry. Report of a WHO Expert Committee. 1995; 854: 1-453. Geneva, WHO. WHO Technical Report Series. Ref Type: Report.
2. The World Health Organization MONICA Project (monitoring trends and determinants in cardiovascular disease): a major international collaboration. WHO MONICA Project Principal Investigators. *J Clin Epidemiol* 1988; 41: 105-14.
3. De Onis M, Garza C, Onyango AW, Martorell R. WHO Child growth standards. Taylor and Francis, 2006.
4. Pérez-Cueto A, Almanza M, Kolsteren P. Female gender and wealth are associated to overweight among adolescents in La Paz, Bolivia. *Eur J Clin Nutr* 2005; 59: 82-7.

5. Ogden CL, Kuczmarski RJ, Flegal KM et al. Centers by Disease Control and Prevention 200 growth charts for the United States: Improvements to the 1977 National Center for Health Statistical version. *Pediatrics* 2002; 109: 45-60.
6. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ* 2000; 320: 1240-3.
7. Flegal KM, Nicholls D, Jackson AA. Body mass index cut offs to define thinness in children and adolescents: international survey. *BMJ* 2007; 8: 299-306.
8. Pérez-Cueto FJA, Naska A, Monterrey J, Almanza-López M, Trichopoulos A, Kolsteren P. Monitoring food and nutrient availability in a nationally representative sample of Bolivian households. *BJN* 2006; 95: 555-567.
9. Pérez-Cueto FJ, Kolsteren P. Changes in the nutritional status of Bolivian women 1994-1998: demographic and social predictors. *Eur J Clin Nutr* 2004; 58: 660-6.
10. Benefice E, López R, Monroy SL, Rodríguez S. Fatness and overweight in women and children from riverine Amerindian communities of the Beni River (Bolivian Amazon). *Am J Hum Biol* 2007; 19: 61-73.
11. Zimmet P, Alberti KG, Kaufman F et al. The metabolic syndrome in children and adolescents-and IDF consensus report. *Pediatr Diabetes* 2007; 8: 299-306.
12. De Onis M, Onyango AW, Borghi E, Garza C, Yang H. Comparison of the World Health Organization (WHO) Child Growth Standards and the National Center for Health Statistics/WHO international growth reference: implications for child health programmes. *Public Health Nutr* 2006; 9: 942-7.
13. Goran MI, Ball GD, Cruz ML. Obesity and risk of type 2 diabetes and cardiovascular disease in children and adolescents. *J Clin Endocrinol Metab* 2003; 88: 1417-27.
14. Hauner H. Insulin resistance and the metabolic syndrome-a challenge of the new millennium. *Eur J Clin Nutr* 2002; 56 (Supl. 1): S25-S29.
15. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA* 2000; 284: 3043-5.
16. Cole TJ. The LMS method for constructing normalized growth standards. *Eur J Clin Nutr* 1990; 44: 45-60.
17. Instituto Nacional de Estadística. Bolivia Indicadores Sociodemográficos por provincial y secciones de provincial 1992-2001. Instituto Nacional de Estadística, 2006.
18. Instituto Nacional de Estadística. Bolivia Proyecciones de población por provincias y municipios, según sexo y grupos de edad, periodo 2000-2010. 2005.