

## Quaternionic wave packets

Stefano De Leo and Gisele C. Ducati

Citation: *J. Math. Phys.* **48**, 052111 (2007); doi: 10.1063/1.2738362

View online: <http://dx.doi.org/10.1063/1.2738362>

View Table of Contents: <http://jmp.aip.org/resource/1/JMAPAQ/v48/i5>

Published by the [AIP Publishing LLC](http://www.aip.org).

---

### Additional information on *J. Math. Phys.*

Journal Homepage: <http://jmp.aip.org/>

Journal Information: [http://jmp.aip.org/about/about\\_the\\_journal](http://jmp.aip.org/about/about_the_journal)

Top downloads: [http://jmp.aip.org/features/most\\_downloaded](http://jmp.aip.org/features/most_downloaded)

Information for Authors: <http://jmp.aip.org/authors>

## ADVERTISEMENT



**physicstoday**

Comment on any  
*Physics Today* article.

Physics Today / Volume 65 / Issue 7 / July 2012  
Previous Article | Next Article

**Measured energy in Japan**  
David von Seggern  
([dvs@seismo.unr.edu](mailto:dvs@seismo.unr.edu)) University of Nevada  
July 2012, page 10  
DIGITAL OBJECT IDENTIFIER  
<http://dx.doi.org/10.1063/PT.3.1619>

The article by Thorne Lay and Hiroo Kanamori (2012) is an excellent review of the 1964 Chilean earthquake. The authors estimate that the earthquake released approximately five times as much energy as the atomic bomb dropped on Nagasaki. The authors also mention that the 1964 Chilean earthquake had still more energy by a factor of about 3, or 15 times as much energy as the atomic bomb dropped on Nagasaki. I believe the authors used the relation for seismic energy release rather than total strain energy release. The seismic energy underestimates the total strain energy release by a variable that depends on the fault plane. Accounting for total strain energy release would increase the earthquake energy number by orders of magnitude.

Despite the catastrophic damage potential of nuclear bombs, the forces of nature occasionally unleash much larger energy releases. Although the nuclear bombs are under our control, earthquakes, volcanic eruptions, and extreme weather events are not. However, by judicious preparation and avoidance measures, humans can significantly diminish the damage of natural events.

This article does not have any references.

**Comment on this article**  
By the act of hitting a ball with a bat, one calculates the force energy to deliver the ball to its new location, but one must also take into account that the ball extended its energy release to that location which became struck by the ball as its momentum ceased and passed energy to the struck team. Therefore the parameters of the damage extend into the future when the received energy to that pushed upon, later becomes released in a new event. Perhaps calculations of one added that in while another's calculations did not. E.M.C.  
Written by Edgar Mocarvill, 14 July 2012 19:59

## Quaternionic wave packets

Stefano De Leo<sup>a)</sup>

*Department of Applied Mathematics, University of Campinas, P.O. Box 6065, Campinas, São Paulo 13083-970, Brazil*

Gisele C. Ducati<sup>b)</sup>

*Department of Mathematics, University of Parana, P.O. Box 19081, Curitiba, Paraná 81531-970, Brazil*

(Received 28 November 2006; accepted 18 April 2007; published online 31 May 2007)

We compare the behavior of a wave packet in the presence of a complex and a pure quaternionic potential step. This analysis, done for a Gaussian convolution function, sheds new light on the possibility to recognize quaternionic deviations from standard quantum mechanics. © 2007 American Institute of Physics. [DOI: [10.1063/1.2738362](https://doi.org/10.1063/1.2738362)]

### I. INTRODUCTION

This article represents the third work of the authors on the Schrödinger equation in the presence of a quaternionic potential step. In the first paper,<sup>1</sup> we have shown that, for such a potential, it is possible to calculate an analytic plane wave solution. This represents, to the best of our knowledge, the first case in which an analytic solution has been given for a quaternionic quantum mechanical potential problem (previous plane wave studies, regarding the barrier<sup>2-4</sup> and well<sup>5</sup> potentials, required numerical calculations). The possibility to work with an analytic plane wave solution allowed, in our second paper,<sup>6</sup> a detailed discussion of the quaternionic diffusion through the stationary phase method. The motivation to write a third paper on this topic is mainly due to the old wish of the authors to find qualitative differences between complex and quaternionic formulations of quantum mechanics which could be useful in identifying the evidence of quaternionic potentials in the case in which such potentials really exist. In this spirit, the quaternionic diffusion, at the moment discussed from a general point of view by using the stationary phase method, should be investigated by directly analyzing the motion of the incident, reflected, and transmitted wave packets in a potential step problem. The numerical results of this study can then be interpreted by looking for analytical approximations of the incident, reflected, and transmitted wave packets in the case of Gaussian convolution functions. The comparison between the complex and pure quaternionic case for the potential step seems to be the best starting point to analyze the qualitative differences between quaternionic and complex quantum mechanics. Indeed, more complicated potentials can often be seen as successive potential steps. For example, a potential barrier can be successfully studied as a two-step problem.<sup>7,8</sup> The investigation proposed in this paper could also give a final answer to the old question concerning the possibility to fit a pure quaternionic potential by a complex one and consequently, to never recognize quaternionic deviations from standard quantum mechanics.

### II. PLANE WAVE ANALYSIS

Before we give the explicit plane wave solution for a pure quaternionic potential step, let us show how, by a simple rephasing of the wave function, we can transform, without loss of generalization, the  $(j, k)$  part of the quaternionic potential,

---

<sup>a)</sup>Electronic mail: [deleo@ime.unicamp.br](mailto:deleo@ime.unicamp.br)

<sup>b)</sup>Electronic mail: [ducati@mat.ufpr.br](mailto:ducati@mat.ufpr.br)

$$iV_1 + jV_2 + kV_3,$$

into a pure  $j$  part, i.e.,

$$iV_1 + j\sqrt{V_2^2 + V_3^2}.$$

This will greatly simplify our presentation. Let us begin by considering the quaternionic Schrödinger equation,<sup>9</sup>

$$i\frac{\hbar^2}{2m}\Phi_{,xx}(x,t) - (iV_1 + jV_2 + kV_3)\Phi(x,t) = \hbar\Phi_t(x,t). \quad (1)$$

This partial differential equation can be reduced by using the well-known separation of variables,

$$\Phi(x,t) = \varphi(x)\exp[-iEt/\hbar],$$

to the following second order ordinary differential equation,<sup>10</sup>

$$i\frac{\hbar^2}{2m}\varphi''(x) - (iV_1 + jV_2 + kV_3)\varphi(x) = -\varphi(x)Ei. \quad (2)$$

By multiplying (from the left) the previous equation by an unitary complex number  $e^{i\alpha}$  and rewriting the pure quaternionic part of the potential as

$$jV_2 + kV_3 = j\sqrt{V_2^2 + V_3^2}e^{-i\theta},$$

where  $\theta = \arctan[V_3/V_2]$ , we find

$$e^{i\alpha}\left[i\frac{\hbar^2}{2m}\varphi''(x) - (iV_1 + j\sqrt{V_2^2 + V_3^2}e^{-i\theta})\varphi(x)\right] = -e^{i\alpha}\varphi(x)Ei. \quad (3)$$

By observing that  $e^{i\alpha}j = je^{-i\alpha}$ , we can rewrite the previous equation as follows:

$$i\frac{\hbar^2}{2m}[e^{i\alpha}\varphi(x)]'' - iV_1[e^{i\alpha}\varphi(x)] - j\sqrt{V_2^2 + V_3^2}[e^{-i(\alpha+\theta)}\varphi(x)] = -[e^{i\alpha}\varphi(x)]Ei. \quad (4)$$

Consequently, the choice  $\alpha = -\theta/2$  conduces to

$$i\frac{\hbar^2}{2m}\psi''(x) - (iV_1 + j\sqrt{V_2^2 + V_3^2})\psi(x) = -\psi(x)Ei, \quad (5)$$

where

$$\psi(x) = e^{-i(\theta/2)}\varphi(x).$$

It is important to observe that constant phases play no role in the stationary phase method and, thus, such a rephasing has no physical effect on the motion of the incident, reflected, and transmitted wave packets. For the diffusion problem,  $E > V_0$ , we shall compare the complex case,  $V_0 = V_1$  ( $V_2 = V_3 = 0$ ), with the pure quaternionic case,  $V_0 = \sqrt{V_2^2 + V_3^2}$  ( $V_1 = 0$ ).

### A. Complex potential step

By setting  $V_1 = V_0$  and  $V_2 = V_3 = 0$  in Eq. (5), we find the standard Schrödinger equation,

$$i\frac{\hbar^2}{2m}\psi_c''(x) - iV_0\psi_c(x) = -\psi_c(x)Ei, \quad (6)$$

whose analytic plane wave solution is given by<sup>11</sup>

$$\begin{aligned}\psi_{c,I}(\epsilon, x) &= \exp[i\epsilon x] + r_c(\epsilon)\exp[-i\epsilon x] \quad \text{[(I) free region } (x < 0)], \\ \psi_{c,II}(\epsilon, x) &= t_c(\epsilon)\exp[i\sigma x] \quad \text{[(II) potential region } (x > 0)],\end{aligned}\quad (7)$$

where

$$\epsilon = \sqrt{2mE/\hbar}, \quad \sigma = \sqrt{2m(E - V_0)/\hbar},$$

and

$$r_c(\epsilon) = (\epsilon - \sigma)/(\epsilon + \sigma), \quad t_c(\epsilon) = 2\epsilon/(\epsilon + \sigma).$$

### B. Pure quaternionic potential step

By setting in Eq. (5)  $V_1=0$  and  $\sqrt{V_2^2+V_3^2}=V_0$ , we obtain

$$i\frac{\hbar^2}{2m}\psi_q''(x) - jV_0\psi_q(x) = -\psi_q(x)Ei. \quad (8)$$

The analytic plane wave solution for the pure quaternionic potential step reads

$$\begin{aligned}\psi_{q,I}(\epsilon, x) &= \exp[i\epsilon x] + r_q(\epsilon)\exp[-i\epsilon x] + j\tilde{r}_q(\epsilon)\exp[\epsilon x] \quad \text{[(I) free region],} \\ \psi_{q,II}(\epsilon, x) &= (1 + jw)t_q(\epsilon)\exp[i\rho x] + (\bar{w} + j)\tilde{t}_q(\epsilon)\exp[-\rho x] \quad \text{[(II) potential region],}\end{aligned}\quad (9)$$

where

$$\epsilon = \sqrt{2mE/\hbar}, \quad \rho = \sqrt{2m\sqrt{E^2 - V_0^2}/\hbar}, \quad w = -iV_0/(E + \sqrt{E^2 - V_0^2}),$$

and

$$r_q(\epsilon) = (\epsilon - \rho)\exp[i \arctan(\epsilon/\rho)]/\sqrt{\epsilon^2 + \rho^2},$$

$$\tilde{r}_q(\epsilon) = (1 + i)\epsilon w/(\epsilon + \rho),$$

$$t_q(\epsilon) = \epsilon/\rho,$$

$$\tilde{t}_q(\epsilon) = \sqrt{\epsilon^2 + \rho^2}\epsilon w \exp[-i \arctan(\rho/\epsilon)]/[\rho(\epsilon + \rho)].$$

For a detailed derivation of the plane wave solution for a quaternionic potential step, we refer the reader to the paper cited in Ref. 1

### III. WAVE PACKET ANALYSIS

Until now, we have been concerned only with plane waves. In this section, we are going to study the time evolution of quaternionic wave packets and deduce from them several important results. The principle of superposition guarantees that every *real* linear combination of the plane waves in region I and region II,

$$\Psi_{I,II}(x, t) = \int d\epsilon g(\epsilon)\psi_{I,II}(\epsilon, x)\exp[-iEt/\hbar] \quad [E = (\hbar\epsilon)^2/2m],$$

will satisfy the Schrödinger equation in the presence of a potential step. The reason why the use of the wave packet formalism is very interesting lies in the fact that the calculations, in the case of a Gaussian convolution function

$$g(\epsilon) = \exp[a^2(\epsilon - \epsilon_0^2)/4]/2\sqrt{\pi},$$

can be analytically approximated. This will allow us to check and interpret our numerical results. In Sec. III A, we first discuss the wave packets motion in the case of a standard (complex) potential step and, then, we analyze the special case of a pure quaternionic potential.

### A. Complex case

The wave packets in region I and region II are given by

$$\Omega_{c,I}(x,t) = \int_{a\epsilon_{\min}}^{+\infty} d(a\epsilon)g(\epsilon)\{\exp[i\epsilon x] + r_c(\epsilon)\exp[-i\epsilon x]\}\exp[-i\epsilon^2\hbar t/2m],$$

$$\Omega_{c,II}(x,t) = \int_{a\epsilon_{\min}}^{+\infty} d(a\epsilon)g(\epsilon)\{t_c(\epsilon)\exp[i\sigma x]\}\exp[-i\epsilon^2\hbar t/2m],$$

where

$$a\epsilon_{\min} = a\sqrt{2mV_0/\hbar}.$$

We find three wave packets: incident, reflected, and transmitted,

$$\Omega_{c,inc}(x,t) = \int_{a\epsilon_{\min}}^{+\infty} d(a\epsilon)g(\epsilon)\exp[i(\epsilon x - \epsilon^2\hbar t/2m)],$$

$$\Omega_{c,ref}(x,t) = \int_{a\epsilon_{\min}}^{+\infty} d(a\epsilon)g(\epsilon)r_c(\epsilon)\exp[-i(\epsilon x + \epsilon^2\hbar t/2m)],$$

$$\Omega_{c,tra}(x,t) = \int_{a\epsilon_{\min}}^{+\infty} d(a\epsilon)g(\epsilon)t_c(\epsilon)\exp[i(\sigma x - \epsilon^2\hbar t/2m)].$$

The choice of a Gaussian convolution function  $g(\epsilon)$  peaked in  $\epsilon_0$  ( $\epsilon_0 > \epsilon_{\min}$ ) and whose value is practically zero near to  $\epsilon_{\min}$ , i.e.,  $g(\epsilon_{\min}) \approx 0$ , allows us to legitimately approximate the incident wave packet as follows:

$$\begin{aligned} \Omega_{c,inc}(x,t) &\approx \int_{-\infty}^{+\infty} d(a\epsilon)g(\epsilon)\exp[i(\epsilon x - \epsilon^2\hbar t/2m)] \\ &= \exp[i(\epsilon_0 x - a^2\epsilon_0^2\pi/2) - i\arctan(2\tau)/2] \frac{\exp[-(x/a - a\epsilon_0\tau)^2/(1+2i\tau)]}{(1+4\tau^2)^{1/4}}, \end{aligned} \quad (10)$$

where  $\tau = \hbar t/ma^2$ . In the case in which the variation of  $r_c(\epsilon)$  can be neglected compared to that of  $g(\epsilon)$ , the reflected wave has the same form as the incident wave packet,<sup>11</sup>

$$\Omega_{c,ref}(x,t) \approx r_c(\epsilon_0) \int_{a\epsilon_{\min}}^{+\infty} d(a\epsilon)g(\epsilon)\exp[-i(\epsilon x + \epsilon^2\hbar t/2m)] = r_c(\epsilon_0)\Omega_{c,inc}(-x,t). \quad (11)$$

For the transmitted wave packet, by neglecting the variation of  $t_c(\epsilon)$  compared to that of  $g(\epsilon)$  and by approximating  $\sigma$  by its first order Taylor expansion,  $\sigma_0 + (\epsilon - \epsilon_0)\epsilon_0/\sigma_0$ , we find

$$\begin{aligned}\Omega_{c,\text{tra}}(x,t) &\approx t_c(\epsilon_0)\exp[i(\sigma_0^2 - \epsilon_0^2)x/\sigma_0] \int_{a\epsilon_{\min}}^{+\infty} d(a\epsilon)g(\epsilon)\exp[i(\epsilon\epsilon_0x/\sigma_0 - \epsilon^2\hbar t/2m)] \\ &= t_c(\epsilon_0)\exp[i(\sigma_0^2 - \epsilon_0^2)x/\sigma_0]\Omega_{c,\text{inc}}(\epsilon_0x/\sigma_0,t).\end{aligned}\quad (12)$$

The reflection probability (the ratio between the probabilities of finding the particle in the reflected packet, at a positive time  $t_0$ , and in the incident packet, at time  $-t_0$ ) is given by

$$\begin{aligned}P_{c,\text{ref}} &= \int_{-\infty}^0 dx |\Omega_{c,\text{ref}}(x,t_0)|^2 \Big/ \int_{-\infty}^0 dx |\Omega_{c,\text{inc}}(x,-t_0)|^2 \\ &\approx |r_c(\epsilon_0)|^2 \int_{-\infty}^0 dx |\Omega_{c,\text{inc}}(-x,t_0)|^2 \Big/ \int_{-\infty}^0 dx |\Omega_{c,\text{inc}}(x,-t_0)|^2 = |r_c(\epsilon_0)|^2.\end{aligned}\quad (13)$$

Similarly, the transmission probability is given by

$$\begin{aligned}P_{c,\text{tra}} &= \int_0^{+\infty} dx |\Omega_{c,\text{tra}}(x,t_0)|^2 \Big/ \int_{-\infty}^0 dx |\Omega_{c,\text{inc}}(x,-t_0)|^2 \\ &\approx |t_c(\epsilon_0)|^2 \int_0^{+\infty} dx |\Omega_{c,\text{inc}}(\epsilon_0x/\sigma_0,t_0)|^2 \Big/ \int_{-\infty}^0 dx |\Omega_{c,\text{inc}}(x,-t_0)|^2 \\ &= |t_c(\epsilon_0)|^2 \frac{\sigma_0}{\epsilon_0} \int_0^{+\infty} dx |\Omega_{c,\text{inc}}(x,t_0)|^2 \Big/ \int_{-\infty}^0 dx |\Omega_{c,\text{inc}}(x,-t_0)|^2 = \frac{\sigma_0}{\epsilon_0} |t_c(\epsilon_0)|^2.\end{aligned}\quad (14)$$

It is easy to verify that

$$P_{c,\text{ref}} + P_{c,\text{tra}} = \left[ \left( \frac{\epsilon - \sigma}{\epsilon + \sigma} \right)^2 + \frac{\sigma}{\epsilon} \frac{4\epsilon^2}{(\epsilon + \sigma)^2} \right]_0 = 1.$$

## B. Pure quaternionic case

The wave packets in region I and region II are given by

$$\Omega_{q,\text{I}}(x,t) = \int_{a\epsilon_{\min}}^{+\infty} d(a\epsilon)g(\epsilon)\{\exp[i\epsilon x] + r_q(\epsilon)\exp[-i\epsilon x] + j\tilde{r}_q(\epsilon)\exp[\epsilon x]\}\exp[-i\epsilon^2\hbar t/2m],$$

$$\begin{aligned}\Omega_{q,\text{II}}(x,t) &= \int_{a\epsilon_{\min}}^{+\infty} d(a\epsilon)g(\epsilon)\{t_q(\epsilon)\exp[i\rho x] + \bar{w}\tilde{t}_q(\epsilon)\exp[-\rho x]\}\exp[-i\epsilon^2\hbar t/2m] \\ &\quad + j \int_{a\epsilon_{\min}}^{+\infty} d(a\epsilon)g(\epsilon)\{w t_q(\epsilon)\exp[i\rho x] + \tilde{t}_q(\epsilon)\exp[-\rho x]\}\exp[-i\epsilon^2\hbar t/2m].\end{aligned}$$

For sufficiently large negative  $t$ , only the (complex) incident wave packet

$$\Omega_{q,\text{inc}}(x,t \ll 0) = \int_{a\epsilon_{\min}}^{+\infty} d(a\epsilon)g(\epsilon)\exp[i(\epsilon x - \epsilon^2\hbar t/2m)] = \Omega_{c,\text{inc}}(x,t)$$

exists. For sufficiently large positive  $t$ , the evanescent wave packets disappear and we find only one (complex) reflected wave packet,

$$\Omega_{q,\text{ref}}(x, t \geq 0) = \int_{a\epsilon_{\min}}^{+\infty} d(a\epsilon)g(\epsilon)r_q(\epsilon)\exp[-i(\epsilon x + \epsilon^2\hbar t/2m)] \approx r_q(\epsilon_0)\Omega_{c,\text{inc}}(-x, t),$$

and one (quaternionic) transmitted wave packet,

$$\begin{aligned}\Omega_{q,\text{tra}}(x, t \geq 0) &= \int_{a\epsilon_{\min}}^{+\infty} d(a\epsilon)g(\epsilon)(1+jw)t_q(\epsilon)\exp[i(\rho x - \epsilon^2\hbar t/2m)] \\ &\approx (1+jw_0)t_q(\epsilon_0)\exp[i(\rho_0^4 - \epsilon_0^4)x/\rho_0^3]\Omega_{c,\text{inc}}(\epsilon_0^3x/\rho_0^3, t).\end{aligned}$$

A simple calculation shows that

$$P_{q,\text{ref}} = \int_{-\infty}^0 dx |\Omega_{q,\text{ref}}(x, t_0)|^2 \Big/ \int_{-\infty}^0 dx |\Omega_{q,\text{inc}}(x, -t_0)|^2 \approx |r_q(\epsilon_0)|^2 \quad (15)$$

and

$$P_{q,\text{tra}} = \int_0^{+\infty} dx |\Omega_{q,\text{tra}}(x, t_0)|^2 \Big/ \int_{-\infty}^0 dx |\Omega_{q,\text{inc}}(x, -t_0)|^2 \approx \frac{\rho_0^3}{\epsilon_0^3} (1 + |w_0|^2) |t_q(\epsilon_0)|^2. \quad (16)$$

Finally,

$$P_{q,\text{ref}} + P_{q,\text{tra}} = \left[ \frac{(\epsilon - \rho)^2}{\epsilon^2 + \rho^2} + \frac{\rho^3}{\epsilon^3} \left( 1 + \frac{\epsilon^2 - \rho^2}{\epsilon^2 + \rho^2} \right) \frac{\epsilon^2}{\rho^2} \right]_0 = 1.$$

#### IV. NUMERICAL RESULTS AND INTERPRETATION

The results of our numerical study are plotted in Fig. 1, where the probability densities  $|\Omega_c(x, t)|^2$  (complex case) and  $|\Omega_q(x, t)|^2$  (pure quaternionic case) are drawn as a function of  $x/a$  for different values of  $\tau = \hbar t / ma^2$ . The incident wave packets are centered in  $E_0 = 2V_0$  (diffusion) and the potential is given by  $a\sqrt{2mV_0} = 10^2\hbar$ . The data show an interesting phenomenon. For the same potential and the same incident energy, the reflected waves have different amplitudes in the case of a complex or a pure quaternionic potential step, and the transmitted waves move with different velocities. The quaternionic transmitted wave packet moves faster than the complex one. Let us try to understand these results by using the analytic approximations given in Sec. III for the incident, reflected, and transmitted wave packets and by using the fact that for small times,  $\tau \ll 1$ , no spreading effect is present. This allows us to simplify the expressions given in Sec. III for the complex and quaternionic wave packets. The amplitude of the incident wave is then given by

$$|\Omega_{q,\text{inc}}(x, t)| \approx |\Omega_{c,\text{inc}}(x, t)| \approx \exp[-(x/a - a\epsilon_0\tau)^2].$$

For the reflected waves, we have

$$|\Omega_{c,\text{ref}}(x, t)| \approx |r_c(\epsilon_0)| \exp[-(x/a + a\epsilon_0\tau)^2],$$

$$|\Omega_{q,\text{ref}}(x, t)| \approx |r_q(\epsilon_0)| \exp[-(x/a + a\epsilon_0\tau)^2].$$

This implies

$$|\Omega_{c,\text{ref}}^{\max}(x, t)|^2 / |\Omega_{c,\text{inc}}^{\max}(x, t)|^2 \approx |r_c(\epsilon_0)|^2 \approx 2.94 \times 10^{-2},$$

$$|\Omega_{q,\text{ref}}^{\max}(x, t)|^2 / |\Omega_{q,\text{inc}}^{\max}(x, t)|^2 \approx |r_q(\epsilon_0)|^2 \approx 2.58 \times 10^{-3},$$




FIG. 1. Diffusion ( $E_0=2V_0$ ) of a wave packet at a complex and pure quaternionic potential step. The complex ( $V_0=V_1$ ) and quaternionic ( $V_0=\sqrt{V_2^2+V_3^2}$ ) potentials act for positive  $x$ . For negative time ( $\tau < 0$ ) only the incident packet is present and it moves toward the step. After a certain time, we find four packets. The reflected packets,  $|\Omega_{c,inc}(x, t)|^2$  and  $|\Omega_{q,inc}(x, t)|^2$ , are returning to the left moving with the same velocity. The transmitted packets,  $|\Omega_{c,tra}(x, t)|^2$  and  $|\Omega_{q,tra}(x, t)|^2$ , propagate toward the right and move with different velocities.

$$|\Omega_{c,ref}(x, t)|^2 / |\Omega_{q,ref}(x, t)|^2 \approx |r_c(\epsilon_0)|^2 / |r_q(\epsilon_0)|^2 \approx 11.4,$$

which confirms the numerical results given in Fig. 1 (negative  $x$  axis). We can also see that the reflected waves move with the same velocity,

$$v_{c,ref} = v_{q,ref} = -\hbar \epsilon_0 / m. \tag{17}$$

For the transmitted waves, by observing that

$$|\Omega_{c,tra}(x, t)| \approx |t_c(\epsilon_0)| \exp[-(\epsilon_0/\sigma_0)^2(x/a - a\sigma_0\tau)^2],$$

$$|\Omega_{q,tra}(x, t)| \approx |(1 + jw_0)t_q(\epsilon_0)| \exp[-(\epsilon_0/\rho_0)^6(x/a - a\rho_0^3\tau/\epsilon_0^2)^2],$$

we find that the centers of the complex and quaternionic transmitted wave packets move, respectively, with velocities

$$v_{c,tra} = \hbar, \sigma_0 / m \quad \text{and} \quad v_{q,tra} = \hbar \rho_0^3 / m \epsilon_0^2. \tag{18}$$

Consequently,



$$\frac{x_{c,tra}^{\max}(t)}{a} = \sqrt{\frac{E_0}{V_0} - 1} \frac{a\sqrt{2mV_0}}{\hbar} \frac{\hbar t}{ma^2},$$

$$\frac{x_{q,tra}^{\max}(t)}{x_{c,tra}^{\max}(t)} = \left[ \left( \frac{E_0}{V_0} - 1 \right) \left( \frac{E_0}{V_0} + 1 \right)^3 \left( \frac{V_0}{E_0} \right)^4 \right]^{1/4},$$

and by using

$$E_0 = 2V_0, \quad \frac{a\sqrt{2mV_0}}{\hbar} = 10^2, \quad \tau = \frac{\hbar t}{ma^2},$$

we find

$$\frac{x_{c,tra}^{\max}(\tau)}{a} = 10^2 \tau,$$

$$\frac{x_{q,tra}^{\max}(\tau)}{x_{c,tra}^{\max}(\tau)} \approx 1.14.$$

This agrees with our numerical calculations as shown by the motion of the complex and quaternionic transmitted wave packets plotted in Fig. 1 (positive  $x$  axis). For example, at time  $\tau=0.15$  the maximum of the complex transmitted wave is found at  $x=15a$ , whereas the center of the quaternionic wave packet (which moves faster) reaches at the same time the point  $x \approx 17.1a$ .

## V. CONCLUSIONS

In the last few years, the Schrödinger equation in the presence of quaternionic potentials with constant and space-dependent phases has been a matter of study and discussion in the literature.<sup>4-6</sup> Some properties of this class of potentials are discussed in detail in the Adler book<sup>9</sup> which represents a milestone in such a field. If quaternionic quantum mechanics represents a possible way to describe the nature, then it becomes relevant to examine how the predictions of standard theories may be affected by changing from complex to quaternionic potentials. The first theoretical analysis of quaternionic potential barriers was developed in Refs. 2 and 3 and showed that in contrast with the standard complex case where the left and right transmission probabilities are equal in magnitude and phase, in the quaternionic quantum mechanics only the magnitude are equal. So, the measurement of a phase shift should be an indicator of quaternionic effects *and* space-dependent phase. Nevertheless, as remarked in Ref. 9, experiments to detect a phase shift are equivalent to detecting time-reversal violation and, consequently, cannot be seen in neutron-optical experiments.<sup>12-14</sup> A more complete phenomenology of the quaternionic potential barrier is given in Ref. 4. With respect to previous works regarding potential barrier diffusion, in this paper, we have introduced the quaternionic wave packet formalism. This allow us to study new *qualitative* differences between standard quantum mechanics and theoretical solutions obtained by solving the Schrödinger equation in the presence of a quaternionic step. For a detailed discussion of experimental proposals on quaternionic potentials and quantum mechanical systems in which a Cahrge-Parity (CP) violation is interpreted by potentials with a space-dependent phase, we refer the reader to Refs. 1 and 4.

In this paper, we have seen that for incoming particles with an energy spectrum centered in  $E_0=2V_0$  the complex and quaternionic transmitted wave packets move with different velocities,

$$\frac{v_{q,\text{tra}}}{v_{c,\text{tra}}} = \frac{\sqrt{3}\sqrt{3}}{2}.$$

Due to the fact that the complex and quaternionic incident wave packets coincide for large negative times (incoming particles), we could try to fit the motion of the quaternionic wave packet by using a different complex potential step. For example, by choosing a *new* complex potential step  $W_0$  such that

$$\sqrt{\frac{E_0}{W_0} - 1} = \left[ \left( \frac{E_0}{V_0} \right)^2 - 1 \right]^{3/4} \frac{V_0}{E_0},$$

we find  $x_{q,\text{tra}}^{\text{max}}(t) = x_{c,\text{tra}}^{\text{max}}(t)$ . In the particular case investigated in this paper,  $E_0 = 2V_0$ , this implies

$$\frac{E_0}{W_0} = 1 + \frac{3\sqrt{3}}{4}.$$

In this scenario, we have the same incident wave and complex or quaternionic transmitted wave packets which move with the same velocity (note that the reflected waves, for the same incoming energy spectrum, always move with the same velocity because they propagate in the free potential region). Once we have guaranteed the same velocities in the potential region, we have to check the probability of transmission (or equivalently the probability of reflection). A simple calculation (see Sec. III) shows that

$$P_{c,\text{ref}} \approx \left( \frac{\sqrt{E_0} - \sqrt{E_0 - W_0}}{\sqrt{E_0} + \sqrt{E_0 - W_0}} \right)^2 \approx 2.01 \times 10^{-2},$$

$$P_{q,\text{ref}} \approx \frac{[\sqrt{E_0} - (E_0^2 - V_0^2)^{1/4}]^2}{E_0 - \sqrt{E_0^2 - V_0^2}} \approx 2.58 \times 10^{-3}.$$

This gives an interesting answer to the old question concerning the possibility to fit a quaternionic potential by a complex one and, consequently, to never recognize deviations from standard quantum mechanics. The study presented for a very simple potential, i.e., the potential step, shows that incoming particles with a given energy spectrum behave differently if they are diffused by a complex or a pure quaternionic potential step. This result, obtained by numerical calculations and interpreted by our Gaussian analytical approximations, cannot be seen in the plane wave analysis. In such a limit, we only have reflection and transmission probabilities, and the quaternionic case can *always* be interpreted in terms of a complex potential step which appropriately fits the quaternionic probabilities. This paper stimulates further investigations; for example, it should be interesting to study within the wave packet formalism, diffusion, and tunneling by complex and pure quaternionic potential barriers which, surely, represents a more realistic potential to be tested by an experimental analysis. In the case of the potential barrier, it is of great interest to examine the tunneling phenomena. In particular, in a forthcoming paper we aim to analyze in detail the Hartman effect<sup>15,16</sup> in the presence of a pure quaternionic potential. The possibility to treat the potential barrier problem as a two-step problem<sup>7,8</sup> gives a more solid interest to this paper. How it could appear as an academic quantum mechanical exercise plays a fundamental role in understanding the behavior of wave packets in the presence of a more complicate potential. The analysis done in this paper explicitly shows that a *qualitative* difference exists between complex and quaternionic quantum mechanics.

<sup>1</sup> S. De Leo, G. C. Ducati, and T. M. Madureira, J. Math. Phys. **47**, 082106 (2006).

<sup>2</sup> A. J. Davies and B. H. McKellar, Phys. Rev. A **40**, 4209 (1989).

<sup>3</sup> A. J. Davies and B. H. McKellar, Phys. Rev. A **46**, 3671 (1992).

<sup>4</sup> S. De Leo, G. C. Ducati, and C. C. Nishi, J. Phys. A **35**, 5411 (2002).

<sup>5</sup> S. De Leo and G. C. Ducati, J. Phys. A **38**, 3443 (2005).

- <sup>6</sup>S. De Leo and G. C. Ducati, J. Math. Phys. **47**, 102104 (2006).
- <sup>7</sup>A. Bernardini, S. De Leo, and P. Rotelli, Mod. Phys. Lett. A **19**, 2717 (2004).
- <sup>8</sup>S. De Leo and P. Rotelli, Eur. Phys. J. C **46**, 551 (2006).
- <sup>9</sup>S. L. Adler, *Quaternionic Quantum Mechanics and Quantum Fields*, (Oxford University Press, New York, 1995).
- <sup>10</sup>S. De Leo and G. C. Ducati, J. Math. Phys. **42**, 2236 (2001).
- <sup>11</sup>C. Choen-Tannoudji, B. Diu, and F. Lalöe, *Quantum Mechanics*, (Wiley, New York, 1977).
- <sup>12</sup>A. Peres, Phys. Rev. Lett. **42**, 683 (1979).
- <sup>13</sup>H. Kaiser, E. A. George, and S. A. Werner, Phys. Rev. A **29**, 2276 (1984).
- <sup>14</sup>A. G. Klein, Physica B **151**, 44 (1989).
- <sup>15</sup>T. E. Hartman, J. Appl. Phys. **33**, 3427 (1962).
- <sup>16</sup>V. S. Olkhovsky, E. Recami, and J. Jakiel, Phys. Rep. **398**, 133 (2004).