



Genetics and Molecular Biology, 28, 2, 321-327 (2005)
Copyright by the Brazilian Society of Genetics. Printed in Brazil
www.sbg.org.br

Research Article

In silico identification of potential chaperone genes that belong to type III and type IV secretion systems in *Xanthomonas axonopodis* pv *citri*

Letícia Khater^{1,2}, Túlio M. Santos¹, Marcos C. Alegria³, Cassia Docena³, Ana C.R. da Silva³ and Carlos H.I. Ramos^{1,2}

¹Laboratório Nacional de Luz Síncrotron, Centro de Biologia Molecular Estrutural, Campinas SP, Brazil.

²Universidade Estadual de Campinas, Instituto de Biologia, Departamento de Bioquímica, Campinas, SP, Brazil.

³Universidade de São Paulo, Instituto de Química, Departamento de Bioquímica, São Paulo, SP, Brazil.

Abstract

The secretion of bacterial virulence factors and flagellar components requires the assistance of specific type III and flagellar chaperones. Standard computational annotation of the genome of *Xanthomonas axonopodis* pv *citri*, a plant pathogen that causes citrus canker, initially did not identify any genes belonging to these chaperone categories since the primary sequence homology between them was very low. However, in a search for hypothetical proteins with characteristics similar to these chaperones, we have now identified 30 chromosomal and 10 plasmidial potential genes encoding chaperones belonging to types III/IV, and flagellar secretion systems in this organism. The significance of these findings is discussed.

Key words: chaperone, protein secretion, type III and type IV secretion systems, *Xanthomonas*.

Received: August 7, 2004; Accepted: December 8, 2004.

Introduction

Bacterial pathogens use the type III and type IV secretion systems to transfer virulence proteins to animal or plant cells (Wattiau *et al.*, 1996; Winans *et al.*, 1996; Hueck, 1998). Hence, these secretion systems are critical for bacterial pathogenicity and the initiation of disease (Gálan and Collmer, 1999; Cornelis and Van Gijsegem, 2000). The type III and type IV secretion systems require cell-to-cell contact and molecular chaperones for secretion (Deng *et al.*, 1999), and are involved in the assembly of extracellular filaments or pili (Christie, 1997). The flagellum-specific secretion pathway is considered to be the archetype of the type III secretion system in bacteria that interact with plants and animals, and is thus part of the type III superfamily. Thus, the proteins that form the type III apparatus are homologous to proteins that are essential for the assembly of surface flagella required for bacterial motility (Aizawa, 1996; Stephens and Shapiro, 1996; Hueck, 1998; Minamino and MacNab, 1999; MacNab,

2003). The type IV system resembles bacterial conjugation systems and is involved in host cell contact-mediated protein translocation (Covacci *et al.*, 1999).

Although the details of the signaling pathway involved in delivery to the target are still uncertain (see, for instance, Anderson and Schneewind, 1997; Aldridge and Hughes, 2001; Lloyd *et al.*, 2001), it is generally accepted that chaperone binding prevents non-productive pre-secretory associations with its substrates (Hueck, 1998; Wattiau and Cornelis, 1993). The main role suggested for type III chaperones is to maintain the bound proteins in a partially folded conformation, or secretion-competent state, that is able to engage the secretory machinery and travel through the secretory pathway (Woestyn *et al.*, 1996; Bennett and Hughes, 2000; Stebbins and Gálan, 2001).

In general, chaperones of the effector class are responsible for only one secreted protein, the effector, whereas chaperones of the translocator class bind two related proteins (Cornelis and Van Gijsegem, 2000). To date, three classes of type III chaperones have been distinguished based on their target category (Page and Parsot, 2002). The SycE-like family (alpha-beta class) associates with only one effector protein, the SycD/LcrH family associates with

Send correspondence to Carlos H.I. Ramos. Laboratório Nacional de Luz Síncrotron, Centro de Biologia Molecular Estrutural, Caixa Postal 6192, 13084-971 Campinas SP, Brazil. E-mail: cramos@lnls.br

two translocator proteins, and members of the Spa 15 associates with several effector proteins (Page *et al.*, 2002; Pallen *et al.*, 2003 and references therein). The SycD/LcrH family is distinguished by a tetratricopeptide repeat (TPR) motif (Pallen *et al.*, 2003), which is an imperfect 34 amino acid repeat often arranged in tandem arrays (Goebl and Yanagida, 1991).

The importance of secretion chaperones to bacteria is enormous since intact chaperones are required for substrate secretion (Wattiau *et al.*, 1994; Iriarte and Cornelis, 1998) and may act as regulators of virulence gene expression (Lloyd *et al.*, 2001). Thus, to understand the secretion systems of bacteria, the chaperones involved in each pathway need to be identified. *Xanthomonas axonopodis* pv *citri* is the causal agent of citrus canker, a serious but still poorly understood disease that affects most commercial citrus cultivars in more than 30 countries (for a recent review of *Xanthomonas*, see Brunings and Gabriel, 2003). This bacterium has the type III cluster present in its chromosome, and two type IV secretion systems, one present in the chromosome and the other in the plasmid pXAC64 (da Silva *et al.*, 2002). Bonas and coworkers (Fenselau *et al.*, 1992; Buttner and Bonas, 2002 and references therein) showed that certain *Xanthomonas* species use their type III system to inject virulence effector proteins into eukaryotic cells. The genome of *Xanthomonas axonopodis* pv *citri* has been fully sequenced (da Silva *et al.*, 2002) and its initial annotation, based on DNA sequence homology, showed no type III chaperone genes. We have reanalyzed the genome of this bacterium using bioinformatic tools to search for hypothetical proteins with predominant type III and flagellar chaperone characteristics (see Bennett and Hughes, 2000 and others for discussion of these characteristics). The results of this analysis indicate the presence of secretion system chaperone genes in the genome of *Xanthomonas axonopodis* pv *citri*.

Material and Methods

Prevalent structural characteristics of type III chaperones used to analyse the genome of *Xanthomonas axonopodis* pv *citri*

The genes encoding particular type III chaperones are often located adjacent to the gene encoding their particular effector in the genome. Thus, the co-expression of these genes increases the performance of the system. Although primary sequence identity is low or not evident among type III and flagellar chaperones, these proteins share several features that can be used to classify them in their respective classes. Such features include a low molecular mass (< 20 kDa), a generally low isoelectric point, and a secondary structure predicted to be predominantly helical with a putative amphipathic α -helix located at the N- or C-terminus (for a discussion of these features see Wattiau *et al.*, 1996; Gygi *et al.*, 1997; Fraser *et al.*, 1999; Bennett and

Hughes, 2000). The presence of such features in a hypothetical gene sequence provides the first indication of the function of the protein until more data become available.

Homology search strategy

To find potential type III and flagellar chaperone genes in *Xanthomonas axonopodis* pv *citri*, we used two independent strategies to search the genome of this pathogen (Figure 1). In the first round, a database containing sequences belonging to type III and flagellar chaperones from several species of bacteria was generated using the Entrez databank (<http://www.ncbi.nlm.nih.gov/Entrez/>). These sequences were aligned using the DIALIGN 2 (Morgenstern, 1999) and CLUSTAL W (Thompson *et al.*, 1994) programs. This alignment allowed redundant sequences in the type III chaperones database to be discarded and decreased their number. BlastP software (Altschul *et al.*, 1990) was then used to compare the generated database with the *Xanthomonas axonopodis* pv *citri* genome in order to find potential type III and flagellar chaperone genes. The default parameters used in this analysis included BLOSUM62 matrix, an e-value for inclusion in subsequent iterations of 0.005, a word size of 3, gap costs of existence 10 and extension 1, and no filter. The ANTHEPROT (ANalyse THE PROTeins; Deleage *et al.*, 1988, 2001) and Compute pI/Mw (Bjellqvist *et al.*, 1993, 1994) software packages,




Figure 1 - Schematic representation of the two strategies used to search the *Xanthomonas axonopodis* pv *citri* genome for potential type III chaperone genes. The figure schematically represents the two strategies used to search the *Xanthomonas axonopodis* pv *citri* genome (da Silva *et al.*, 2002) for potential type III chaperone genes. See Material and Methods for details and software references.

which perform large scale protein sequence analyses to predict protein secondary structure and physico-chemical parameters, were used to determine whether the potential *Xanthomonas axonopodis* pv *citri* type III and flagellar chaperones genes found had the global type III chaperone characteristics described above. In the second round of analysis, the genes in the *Xanthomonas axonopodis* pv *citri* genome (da Silva *et al.*, 2002) located within 1.5 kb of pathogenic and virulence clusters, were analysed individually with the ANTHEPROT and Compute pI/Mw packages to identify potential physico-chemical characteristics similar to type III chaperones.

PSI-BLAST search for conserved domains in secretion chaperones.

A PSI-BLAST search (Altschul *et al.*, 1997) was performed on the NCBI site (<http://www.ncbi.nlm.nih.gov/BLAST/>) using the established default parameters (see above) in order to verify whether the genes selected have characteristic secretion chaperone domains, such as tetratricopeptide (TPR)-like repeats (Pallen *et al.*, 2003), Spa15 family characteristic residues (Page *et al.*, 2002), or similarity to the SycE family (Woestyn *et al.*, 1996).

Results

The analysis for redundancy in the type III chaperone database generated from the Entrez databank reduced the number of sequences belonging to type III and flagellar system chaperones to 120. The *Xanthomonas axonopodis* pv *citri* genome was searched for genes homologous to each of these 120 sequences. The other hypothetical genes close to pathogenic and virulence clusters in the *Xanthomonas axonopodis* pv *citri* genome were analyzed individually for potential physico-chemical characteristics similar to those of type III chaperones using the programs described in the Material and Methods section. The sequences selected by these two approaches were then ana-

lyzed for their cellular location in order to eliminate sequences not predicted to be in the periplasm. Five of the putative secretion chaperone genes found in the first round were located on the chromosome and one in the plasmid pXAC33, and had type III and flagellar chaperone characteristics, as described in Material and Methods. This approach allowed the identification of proteins with $\geq 45\%$ similarity to type III and flagellar chaperones of other bacteria (Table 1). The second approach selected 25 chromosomal genes and 9 plasmidial genes with secretion chaperones characteristics (Table 2). A total of 40 potential type III/IV or flagellar chaperone genes were identified in the *Xanthomonas axonopodis* pv *citri* genome, with 30 located on the chromosome and 10 located in the two plasmids, pXAC33 and pXAC64 (da Silva *et al.*, 2002).

As an example of this approach, the analysis for two potential type III chaperones in *Xanthomonas axonopodis* pv *citri* is shown in Figure 2. This figure illustrates the amphipathic α -helix wheel predicted from the primary amino acid sequences of the potential chaperones NP_640774 (Figure 2a) and NP_640641 (Figure 2b). The amphipathic α -helix has predominantly nonpolar amino acid residues along one side and polar residues along the other. The presence of an amphipathic α -helix is one of the characteristics of secretion chaperones.

Discussion

Secretory systems are of fundamental importance for several species of bacteria that use the systems to assemble the flagellum and to secrete important, mainly pathogenic factors, into the environment (Hueck, 1998; Gálan and Collmer, 1999; Cornelis and Van Gijsegem, 2000; MacNab, 2003). The acquisition of a type III secretory system allows many gram-negative bacteria to become pathogens since mutants with a disabled type III system are nonpathogenic (Hueck, 1998; Neyt and Cornelis, 1999).

Table 1 - Potential type III chaperone genes identified by sequence homology and predicted protein physico-chemical characteristics. The *Xanthomonas axonopodis* pv *citri* genome was searched for homology to type III chaperones. Five of the identified genes are localized in the chromosome and one in the plasmid pXAC33 (da Silva *et al.*, 2002). All 6 were confirmed for type III chaperone characteristics as described in Material and Methods. This method allowed the identification of genes with 45% (or higher) homology to type III and flagellar chaperones.

Putative secretion chaperone		Predicted physico-chemical parameters			Protein homologue / gene number	S (%)
<i>Xanthomonas axonopodis</i> pv <i>citri</i> Gene ID	Genome localization	MW (kDa)	pI	Residues involved in the Amphipathic domain		
NP_642798.1	C	17.8	5.8	98-111	FliS / gi2276420	49
NP_641681.1	C	10.9	5.6	88-100	FliJ / gi1518878	51
NP_642046.1	C	12.7	6.8	100-117	YscB / gi2635354	63
NP_643331.1	C	20.2	4.5	158-175	FliJ / gi1518878	45
NP_641682.1	C	17.1	5.5	92-102	FliJ / gi1518878	48
NP_644711.1	P	21.3	6.0	152-162	Spa15/gi13449095	56

C, chromosomal gene; P, plasmidial gene; S, similarity.

Table 2 - Potential type III chaperone genes determined by analysis of sequences localized near pathogenic and virulence clusters. Hypothetical genes in the *Xanthomonas axonopodis* pv *citri* genome (da Silva *et al.*, 2002) localized near (< 1.5 kb) pathogenic and virulence clusters, were individually analyzed with the programs described in the Material and Methods section for potential physico-chemical characteristics similar to type III chaperones. This method allowed the identification of 34 potential secretion system chaperone genes, 25 localized at the chromosome and 9 localized at the two plasmids, pXAC33 and pXAC64.

Putative secretion chaperone				
<i>Xanthomonas axonopodis</i> pv <i>citri</i> Gene ID	Genome localization	Predicted physico-chemical parameters		
		MW (kDa)	pI	Residues involved in the Amphipathic domain
Near avirulence genes (avr)				
NP_640641.1	C	18.6	7.0	69-97
NP_643529.1	C	14.7	9.1	69-84
NP_641699.1	C	16.3	5.5	72-83
NP_643565.1	C	12.9	6.9	54-66
NP_641577.1	C	18.1	6.9	61-76
Near hypersensitivity response and pathogenicity genes (Type III cluster)				
NP_641599.1	C	14.3	8.5	54-79
NP_640774.1	C	10.7	7.0	67-91
NP_642316.1	C	12.1	6.7	22-38
Near host cell wall degradation genes				
NP_640498.1	C	11.3	10.3	1-18
NP_640495.1	C	17.1	6.2	16-30
NP_640494.1	C	18.6	7.7	147-165
NP_640523.1	C	14.0	11.3	36-52
NP_643871.1	C	16.6	10.8	80-96
NP_643873.1	C	12.3	7.7	100-114
NP_640383.1	C	15.7	5.62	21-36
NP_643815.1	C	13.3	11.3	1-14
NP_642683.1	C	10.5	4.5	61-77
In the type IV secretion system cluster				
NP_641701.1	C	22.6	6.2	148-165
NP_642723.1	C	17.8	5.2	42-58
NP_642935.1	C	14.7	6.0	55-71
NP_640924.1	C	15.4	4.7	42-60
NP_642924.1	C	17.1	8.1	71-84
NP_642919.1	C	17.2	8.4	53-65
NP_644761.1	P	8.4	5.6	01-13
NP_644763.1	P	15.8	7.9	110-122
NP_644770.1	P	14.2	9.8	123-129
NP_644771.1	P	9.4	9.1	28-41
NP_644776.1	P	13.9	8.6	87-97
NP_644777.1	P	13.6	8.5	71-82
Nearby virB6 protein				
NP_642918.1	C	16.6	5.5	126-142
NP_642914.1	C	16.4	10.2	95-108
Near pathogenicity, virulence and adaptation genes				
NP_644726.1	P	16.1	10.9	96-110
NP_644791.1	P	17.2	12.1	5-21
NP_644792.1	P	16.2	11.7	110-126

C. chromosomal gene; P. plasmidial gene.




Figure 2 - Predicted amphipathic α -helical wheel of potential secretion chaperone. The genes NP_640774 (a) and NP_640641 (b) were selected as potential secretion chaperones because they were located near pathogenic and virulence clusters and because they had predicted protein physico-chemical characteristics similar to those of type III chaperones (see Table 2). Analysis with the ANTHEPROT software package (ANalyse THE PROTEins; Deleage *et al.*, 1988; Deleage *et al.*, 2001) allowed the prediction of a potential amphipathic α -helical wheel at the protein C-terminus. Amphipathic α -helices have predominantly nonpolar amino acid residues along one side and polar residues along the other side.

Type III secretory systems have been found in animal and plant pathogens, and show varying degrees of homology, from highly conserved (some proteins of the secretory apparatus) to quite divergent (chaperones and translocators) or very diverse (effectors) (Bennett and Hughes, 2000; Subtil *et al.*, 2000). The secreted proteins must be targeted to the secretory apparatus in order to be secreted. One of the most important mechanisms for labeling and delivering the proteins to be secreted involves chaperones that recognize the target proteins and direct them to the secretory apparatus (Wattiau and Cornelis, 1993; Hueck, 1998; Bennett and Hughes, 2000; Stebbins and Gálan, 2001). Curiously, no type III chaperone genes were annotated in the fully sequenced genome of *Xanthomonas axonopodis* pv *citri* (da Silva *et al.*, 2002).

Large-scale DNA sequencing efforts produce a huge amount of data that require detailed analysis to be well understood. The analysis of sequences and their annotation by homology is a suitable method to identify potential proteins in a sequenced genome. Since standard techniques did not annotate type III and type IV chaperones in the *Xanthomonas axonopodis* pv *citri* genome, we undertook a more specific analysis of this genome to search for these chaperones. For this, we used homology to search for specific sequences followed by a search for potential physico-chemical properties of the proteins. Several hypothetical genes in the *Xanthomonas axonopodis* pv *citri* genome were analyzed in order to identify those belonging to the secretion chaperone categories. These approaches have been used to show the presence of proteins in certain organisms (see for instance Lawson, 1999; Subtil *et al.*, 2000; Searls, 2000; and others). The comparison of multiple sequences can reveal gene functions that are not detected by simple sequence homology searches.

Secretion system chaperones vary in their amino acid sequences, but share a few common properties, including genome localization close to the gene encoding the target protein, a low isoelectric point, a low molecular weight, and a putative amphipathic α -helix at the N- or C-terminus (Bennett and Hughes, 2000; Parsot *et al.*, 2003). Our analyses of the *Xanthomonas axonopodis* pv *citri* genome using the tools described here allowed the identification of 40 potential secretion system chaperone genes, 30 which were located on the chromosome and 10 in the two plasmids. The first round of the search for potential type III chaperones identified several hypothetical genes that shared similarity with type III chaperones from other bacteria. Two potential chaperones with similarity to the chaperone families SycE and Spa15 were found. The *Xanthomonas axonopodis* pv *citri* gene NP_642046.1 was similar to the secretion chaperone YscB that belongs to the SycE family and functions as a chaperone for the protein YopN in *Yersinia pestis* (Day and Plano 1998). *Xanthomonas axonopodis* pv *citri* gene NP_644711.1 was similar to the secretion chaperone Spa15 from *Shigella flexneri* and also to InvB from other bacteria that belong to a third class of chaperones in the type III secretion pathway (Page *et al.*, 2002). The potential chaperones identified here did not have the TPR domain characteristic of the SycD family (Pallen *et al.*, 2003). Genes belonging to the flagellum assembly apparatus were also identified in *Xanthomonas axonopodis* pv *citri*. Several of these sequences showed homology with FliJ from *Rhodobacter sphaeroides* and FliS from *Pseudomonas aeruginosa*. FliJ is a general flagellar chaperone related to the export process itself and FliS plays a role in protecting the target proteins before exportation (Minamino *et al.*, 2000).

Our results also indicated the presence of potential chaperone genes that may belong to the type IV secretion system because most of the genes identified in the second

round of searching were located close to the type IV system cluster. However, this characteristic may not be enough to support this conclusion and since some type IV chaperones share most of the type III chaperone characteristics (for instance VirE1; Deng *et al.*, 1999) we can only state that the annotated chaperones are related to type III/IV systems.

Type III secretion system occurs in other *Xanthomonas* species (Fenselau *et al.*, 1992; Buttner and Bonas, 2002) and our results indicated the presence of chaperones belonging to this system, and to the flagellar and type IV systems in *Xanthomonas axonopodis* pv *citri*. Together, the results presented here are important for understanding the pathogenicity of *Xanthomonas axonopodis* pv *citri* and may be useful in studies of the mechanism of disease caused by this pathogen in orange plantations. To date, few reports have described a requirement for chaperones in the type III secretion systems of plant pathogens (see for instance, Gaudriault *et al.*, 1997; van Dijk *et al.*, 2002). In conclusion, the study of *Xanthomonas axonopodis* will benefit from the identification of secretion system chaperone genes since uncovering the precise molecular events controlling the delivery of effector proteins should eventually allow the design of compounds that specifically interfere with these processes, hopefully without deleterious side effects to the host plant (Staskawicz, 2001).

Acknowledgments

This work was supported by Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq). LK, TMS, MCA, and CD were supported by fellowship from FAPESP.

References

- Aizawa SI (1996) Flagellar assembly in *Salmonella typhimurium*. *Mol Microbiol* 19:1-5.
- Aldridge P and Hughes KT (2001) How and when are substrates selected for type III secretion? *Trends Microbiol* 9:209-214.
- Altschul SF, Gish W, Miller W, Myers EW and Lipman DJ (1990) Basic local alignment search tool. *J Mol Biol* 215:403-410.
- Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W and Lipman DJ (1997) Gapped BLAST and PSI-BLAST: A new generation of protein database search programs. *Nucleic Acids Res* 25:3389-3402.
- Anderson DM and Schneewind O (1997) A mRNA signal for the type III secretion of Yop proteins by *Yersinia enterocolytica*. *Science* 278:1140-1143.
- Bennett JCQ and Hughes C (2000) From flagellum assembly to virulence. *Trends Microbiol* 8:202-204.
- Bjellqvist B, Basse B, Olsen E and Celis JE (1994) Reference points for comparisons of two-dimensional maps of proteins from different human cell types defined in a pH scale where isoelectric points correlate with polypeptide compositions. *Electrophoresis* 15:529-539.
- Bjellqvist B, Hughes GJ, Pasquali C, Paquet N, Ravier F, Sanchez J-C, Frutiger S and Hochstrasser DF (1993) The focusing positions of polypeptides in immobilized pH gradients can be predicted from their amino acid sequences. *Electrophoresis* 14:1023-1031.
- Blasco F, Dos Santos JP, Magalon A, Frixon C, Guigliarelli B, Santini CL and Giordano G (1998) NarJ is a specific chaperone required for molybdenum cofactor assembly in nitrate reductase A of *Escherichia coli*. *Mol Microbiol* 28:435-447.
- Brunings AM and Gabriel DW (2003) *Xanthomonas citri*: Breaking the surface. *Mol Plant Pathol* 4:41-57.
- Buttner D and Bonas U (2002) Getting across-bacterial type III effector proteins on their way to the plant cell. *EMBO J* 21:5313-5322.
- Christie PJ (1997) *Agrobacterium tumefaciens* T-complex transport apparatus: A paradigm for a new family of multifunctional transporters in eubacteria. *J Bacteriol* 179:3085-3094.
- Cornelis GR and Van Gijsegem F (2000) Assembly and function of type III secretory systems. *Annu Rev Microbiol* 54:735-774.
- Cornelis GR and Wolf-Watz H (1997) The *Yersinia* Yop virulon: A bacterial system for subverting eukaryotic cells. *Mol Microbiol* 23:861-867.
- Covacci A, Telford JL, Del Giudice G, Parsonnet J and Rappuoli R (1999) *Helicobacter pylori* virulence and genetic geography. *Science* 284:1328-1333.
- da Silva ACR, Ferro JA, Reinach FC, Farah CS, Furlan LR, Quaggio RB, Monteiro-Vitorello CB, Van Sluys MA, Almeida NF, Alves LM, do Amaral AM, Bertolini MC, Camargo LE, Camarotte G, Cannavan F, Cardozo J, Chamberg F, Ciapina LP, Cicarelli RM, Coutinho LL, Cursino-Santos JR, El-Dorry H, Faria JB, Ferreira AJ, Ferreira RC, Ferro MI, Formighieri EF, Franco MC, Greggio CC, Gruber A, Katsuyama AM, Kishi LT, Leite RP, Lemos EG, Lemos MV, Locali EC, Machado MA, Madeira AM, Martinez-Rossi NM, Martins EC, Meidanis J, Menck CF, Miyaki CY, Moon DH, Moreira LM, Novo MT, Okura VK, Oliveira MC, Oliveira VR, Pereira HA, Rossi A, Sena JA, Silva C, de Souza RF, Spinola LA, Takita MA, Tamura RE, Teixeira EC, Tezza RI, Trindade dos Santos M, Truffi D, Tsai SM, White FF, Setubal JC and Kitajima JP (2002) Comparison of the genomes of two *Xanthomonas* pathogens with differing host specificity. *Nature* 417:459-463.
- Day JB and Plano GV (1998) A complex composed of SycN and YscB functions as a specific chaperone for YopN in *Yersinia pestis*. *Mol Microbiol* 30:777-788.
- Deleage G, Clerc FF, Roux B and Gautheron DC (1988) ANTHEPROT: A package for protein sequence analysis using a microcomputer. *Comput Appl Biosci* 4:351-356.
- Deleage G, Combet C, Blanchet C and Geourjon C (2001) ANTHEPROT: An integrated protein sequence analysis software with client/server capabilities. *Comput Biol Med* 31:259-267.
- Deng W, Chen L, Peng WT, Liang X, Sekiguchi S, Gordon MP, Comai L and Nester EW (1999) VirE1 is a specific molecular chaperone for the exported single-stranded-DNA-binding protein VirE2 in *Agrobacterium*. *Mol Microbiol* 31:1795-1807.
- Fenselau S, Balbo I and Bonas U (1992) Determinants of pathogenicity in *Xanthomonas campestris* pv *vesicatoria* are related

- to proteins involved in secretion in bacterial pathogens of animals. *Mol Plant Microbe Interact* 5:390-396.
- Fraser GM, Bennett JCQ and Hughes C (1999) Substrate-specific binding of hook-associated proteins by FlgN and FliT, putative chaperones for flagellum assembly. *Mol Microbiol* 32:569-580.
- Gálán JE and Collmer A (1999) Type III secretion machines: Bacterial devices for protein delivery into host cells. *Science* 284:1322-1328.
- Gaudriault S, Malandrin L, Paulin JP and Barny, MA (1997) DspA, an essential pathogenicity factor of *Erwinia amylovora* showing homology with AvrE of *Pseudomonas syringae*, is secreted via the Hrp secretion pathway in a DspB-dependent way. *Mol Microbiol* 26:1057-1069.
- Gygi D, Fraser, G, Dufour A and Hughes C (1997) A motile but non-swarming mutant of *Proteus mirabilis* lacks FlgN, a facilitator of FljC export and flagella assembly. *Mol Microbiol* 25:597-604.
- Goebel M and Yanagida M (1991) The TPR snap helix: A novel protein repeat motif from mitosis to transcription. *Trends Biochem Sci* 16:173-177.
- Hueck CJ (1998) Type III protein secretion systems in bacterial pathogens of animals and plants. *Microbiol Mol Biol Rev* 62:379-433.
- Iriarte M and Cornelis GR (1998) YopT, a new *Yersinia* Yop effector protein, affects the cytoskeleton of host cells. *Mol Microbiol* 29:915-929.
- Lawson D (1999) Data mining parasite genomes: Haystack searching with a computer. *Parasitology* 118:S15-S18.
- Lloyd SA, Forsberg A, Wolf-Watz H and Francis MS (2001) Targeting exported substrates to the *Yersinia* TTSS: Different functions for different signals? *Trends Microbiol* 9:367-371.
- MacNab RM (2003) How bacteria assemble flagella. *Annu Rev Microbiol* 57:77-100.
- Minamino T, Chu R, Yamaguchi S and Macnab RM (2000) Role of FljI in *Salmonella*. *J Bacteriol* 182:4207-4215.
- Minamino T and Macnab RM (1999) Components of the *Salmonella* flagellar export apparatus and classification of export substrates. *J Bacteriol* 181:1388-1394.
- Morgenstern B (1999) DIALIGN 2: Improvement of the segment-to-segment approach to multiple sequence alignment. *Bioinformatics* 15:211-218.
- Nakai K and Kanehisa M (1991) Expert system for predicting protein localization sites in Gram-negative bacteria. *Proteins* 11:95-110.
- Neyt C and Cornelis GR (1999) Insertion of a Yop translocation pore into the macrophage plasma membrane by *Yersinia enterocolitica*: Requirement for translocators YopB and YopD, but not LcrG. *Mol Microbiol* 33:971-981.
- Page A-L, Sansonetti P and Parsot C (2002) Spa15 of *Shigella flexneri*, a third type of chaperone in the type III secretion pathway. *Mol Microbiol* 43:1533-1542.
- Page A-L and Parsot C (2002) Chaperones of the type III secretion pathway: Jacks of all trades. *Mol Microbiol* 46:1-11.
- Pallen MJ, Francis MS and Fütterer K (2003) Tetratricopeptide-like repeats in type-III secretion chaperones and regulators. *FEMS Microbiol Lett* 223:53-60.
- Parsot C, Hamiaux C and Page A-L (2003) The various and varying roles of specific chaperones in type III secretion systems. *Curr Opin Microbiol* 6:7-14.
- Searls DB (2000) Using bioinformatics in gene and drug discovery. *Drug Discov Today* 5:135-143.
- Staskawicz BJ, Mudgett MB, Dangl JL and Galan JE (2001) Common and contrasting themes of plant and animal diseases. *Science* 22:2285-2289.
- Stebbins CE and Galán JE (2001) Maintenance of an unfolded polypeptide by a cognate chaperone in bacterial type III secretion. *Nature* 414:77-81.
- Stephens C and Shapiro L (1996) Bacterial pathogenesis: Delivering the payload. *Curr Biol* 6:927-930.
- Subtil A, Blocker A, and Dautry-Varsat A (2000) Type III secretion system in *Chlamydia species*: Identified members and candidates. *Microbes and Infection* 2:367-369.
- Thompson JD, Higgins DG and Gibson TJ (1994) CLUSTAL W: Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, positions-specific gap penalties and weight matrix choice. *Nucleic Acids Research* 22:4673-4680.
- van Dijk K, Tam VC, Records AR, Petnicki-Ocwieja T and Alfano JR (2002). The ShcA protein is a molecular chaperone that assists in the secretion of the HopPsyA effector from the type III (Hrp) protein secretion system of *Pseudomonas syringae*. *Mol Microbiol* 44:1469-81.
- Wattiau P and Cornelis GR (1993) SycE, a chaperone-like protein of *Yersinia enterocolitica* involved in the secretion of YopE. *Mol Microbiol* 8:123-131.
- Wattiau P, Bernier B, Deslée P, Michiels T and Cornelis GR (1994) Individual chaperones required for Yop secretion by *Yersinia*. *Proc Natl Acad Sci USA* 91:10493-10497.
- Wattiau P, Woestyn S and Cornelis GR (1996) Customized secretion chaperones in pathogenic bacteria. *Mol Microbiol* 20:255-262.
- Winans SC, Burns DL and Christie PJ (1996) Adaptation of a conjugal transfer system for the export of pathogenic macromolecules. *Trends Microbiol* 4:64-68.
- Woestyn S, Sory MP, Boland A, Lequenne O and Cornelis GR (1996). The cytosolic SycE and SycH chaperones of *Yersinia* protect the region of YopE and YopH involved in translocation across eukaryotic cell membranes. *Mol Microbiol* 20:1261-1271.

Associate Editor: Sérgio Olavo Pinto da Costa