

Delay time in quaternionic quantum mechanics

Stefano De Leo and Gisele Ducati

Citation: *J. Math. Phys.* **53**, 022102 (2012); doi: 10.1063/1.3684747

View online: <http://dx.doi.org/10.1063/1.3684747>

View Table of Contents: <http://jmp.aip.org/resource/1/JMAPAQ/v53/i2>

Published by the [AIP Publishing LLC](http://www.aip.org).

Additional information on *J. Math. Phys.*


Journal Homepage: <http://jmp.aip.org/>

Journal Information: http://jmp.aip.org/about/about_the_journal

Top downloads: http://jmp.aip.org/features/most_downloaded

Information for Authors: <http://jmp.aip.org/authors>

ADVERTISEMENT



Give us your dirty old books!
The Niels Bohr Library & Archives is looking
for book donations

Delay time in quaternionic quantum mechanics

Stefano De Leo^{1,a)} and Gisele Ducati^{2,b)}

¹*Department of Applied Mathematics, State University of Campinas, Brazil*

²*CMCC, Universidade Federal do ABC, Brazil*

(Received 20 July 2011; accepted 20 January 2012; published online 13 February 2012)

In looking for quaternionic violations of quantum mechanics, we discuss the delay time for pure quaternionic potentials. Our study shows the energy region which amplifies the difference between quaternionic and complex quantum mechanics.

© 2012 American Institute of Physics. [<http://dx.doi.org/10.1063/1.3684747>]

I. INTRODUCTION

Many papers have introduced the general framework of quaternionic quantum mechanics.^{1–10} The book published in 1995 under the title *Quaternionic quantum mechanics and quantum fields* by Adler¹¹ is, today, recognized to be indispensable to a deeper understanding of the subject. It seems, however, that a more practical side has been somewhat neglected. In other words, the investigation of quantitative and qualitative differences between complex and quaternionic quantum mechanics and the consequent experimental proposals seem to be in the early stage. It was in the hope of filling this gap that the authors have recently reviewed scattering problems in quaternionic quantum mechanics.^{12–14} The more elementary problems, such as square-well potentials, have been carefully re-considered with the *new* tendency to arrive at more practical results. The main purpose was to emphasize quaternionic violations of complex quantum mechanics which could be tested in laboratory experiments. This should make the subject more useful to and accessible over the worldwide community of scientists interested in looking for the existence of quaternionic potentials.

Contrary to what is predicted by classical mechanics, in complex quantum mechanics a particle with energy E_0 is *not* instantaneously reflected by a potential step of height V_0 greater than the incoming energy E_0 .^{15,16} Let us briefly recall the standard introduction to delay times. The center of the incident wave packet arrives at the potential discontinuity, $x = 0$, at time $t = 0$. During a certain interval of time around $t = 0$, the wave packet is localized in the region where the potential discontinuity is found. For sufficiently large times the incident wave packet disappears and the reflected wave packet propagates towards the left at a speed $\sqrt{2 E_0/m}$. By using the stationary phase condition

$$\left[\frac{d}{dE} \left(2\theta_c - \frac{\sqrt{2mE}}{\hbar} x_r - \frac{Et}{\hbar} \right) \right]_0 = 0,$$

where $2\theta_c$ is the phase of the reflection coefficient,

$$R_c = \frac{\sqrt{E} - i\sqrt{V_0 - E}}{\sqrt{E} + i\sqrt{V_0 - E}} = \exp[2i\theta_c], \quad (1)$$

we can calculate the position of the center of the reflected wave packet,

$$\sqrt{\frac{m}{2E_0}} x_r = -t + 2\hbar \left[\frac{d\theta_c}{dE} \right]_0.$$

a) Author to whom correspondence should be addressed. Electronic mail: deleo@ime.unicamp.br.

b) Electronic mail: ducati@ufabc.edu.br.

This clearly shows a *delay time* in reflection given by

$$2\hbar \left[\frac{d\theta_c}{dE} \right]_0 = \frac{\hbar}{\sqrt{E_0(V_0 - E_0)}}. \quad (2)$$

This delay time is due to the fact that, for times close to zero, the probability of presence of the particle in the classically forbidden region is *not* zero. The study of the delay time in quaternionic quantum mechanics will be the subject matter of this paper.

II. ONE-DIMENSIONAL POTENTIALS AND ONE-DIMENSIONAL MOTION

In order to understand the essential features of quaternionic quantum mechanics, we focus our attention on quaternionic one-dimensional problems. They may be studied deriving the equation of motion from the three-dimensional quaternionic Schrödinger equation,¹¹

$$\hbar \partial_t \Phi(\mathbf{r}, t) = \left[i \frac{\hbar^2}{2m} \nabla^2 - i V_1(\mathbf{r}, t) - j V_2(\mathbf{r}, t) - k V_3(\mathbf{r}, t) \right] \Phi(\mathbf{r}, t). \quad (3)$$

If the potential only depends on one of the three spatial coordinates, say x , and the motion is one-dimensional, $p_y = p_z = 0$, the previous equation becomes

$$i \frac{\hbar^2}{2m} \varphi''(x) = [i V_1(x) + j V_q(x) e^{-i\rho(x)}] \varphi(x) - E \varphi(x) i, \quad (4)$$

where

$$V_q(x) = \sqrt{V_2^2(x) + V_3^2(x)} \quad \text{and} \quad \tan \rho(x) = V_3(x) / V_2(x).$$

Let us now analyze the stationary states for the case of a quaternionic step potential defined by

$$\mathbf{h} \cdot \mathbf{V}(x) := \{\mathbf{0} \text{ for } x < 0 \quad \text{and} \quad \mathbf{h} \cdot \mathbf{V} \text{ for } x > 0\}, \quad (5)$$

where $\mathbf{h} = (i, j, k)$. To shorten notation, it is convenient to introduce the following adimensional quantities

$$\frac{\sqrt{2m V_0}}{\hbar} x = \xi, \quad \frac{V_q}{V_0} = v_q, \quad \text{and} \quad \frac{E}{V_0} = \epsilon,$$

where $V_0 = \sqrt{V_1^2 + V_q^2}$. In terms of this new adimensional quantities, the differential equation for the wave function in the free potential region $\xi < 0$ becomes

$$i \varphi_1''(\xi) = -\epsilon \varphi_1(\xi) i. \quad (6)$$

For the case of incident particles coming from the left, the free potential plane wave solution is the region $\xi < 0$ is^{4,13}

$$\varphi_1(\xi) = \exp[i \sqrt{\epsilon} \xi] + R \exp[-i \sqrt{\epsilon} \xi] + j \tilde{R} \exp[\sqrt{\epsilon} \xi], \quad (7)$$

where R and \tilde{R} are complex coefficients to be determined by the matching conditions. The wave function in the potential region, $\xi > 0$, satisfies the following differential equation

$$i \varphi_1''(\xi) = \left(i \sqrt{1 - v_q^2} + j v_q e^{-i\rho} \right) \varphi_1(\xi) - \epsilon \varphi_1(\xi) i, \quad (8)$$

whose solution is^{4,13}

$$\varphi_1(\xi) = (1 + j \gamma) T \exp[i \alpha_- \xi] + (\beta + j) \tilde{T} \exp[-\alpha_+ \xi]. \quad (9)$$

where

$$\alpha_{\pm} = \sqrt{\epsilon^2 - v_q^2} \pm \sqrt{1 - v_q^2}, \quad \beta = i \frac{v_q e^{i\rho}}{\epsilon + \sqrt{\epsilon^2 - v_q^2}}, \quad \gamma = -i \frac{v_q e^{-i\rho}}{\epsilon + \sqrt{\epsilon^2 - v_q^2}}, \quad (10)$$

and T and \tilde{T} are complex coefficients to be determined by the matching conditions.

For diffusion, $E > V_0$, we have $\alpha_{\pm} \in \mathbb{R}_+$. For the case of total reflection, $E < V_0$, we find

$$v_q < \epsilon < 1 : \alpha_- = i \sqrt{\sqrt{1 - v_q^2} - \sqrt{\epsilon^2 - v_q^2}} \in i \mathbb{R}_+ \quad \text{and} \quad \alpha_+ \in \mathbb{R}_+,$$

$$0 < \epsilon < v_q : \alpha_- = i (1 - \epsilon^2)^{1/4} \exp \left[-\frac{i}{2} \arctan \sqrt{\frac{v_q^2 - \epsilon^2}{1 - v_q^2}} \right] \in \mathbb{C}_+ \quad \text{and} \quad \alpha_+ = i \alpha_-^*.$$

III. TOTAL REFLECTION: PHASE CALCULATION

To obtain the reflection coefficient, we have to use the matching conditions at $x = 0$, i.e., $\varphi_I(0) = \varphi_{II}(0)$ and $\varphi_I'(0) = \varphi_{II}'(0)$. These conditions lead to the following quaternionic system,

$$1 + R + j \tilde{R} = (1 + j \gamma) T + (\beta + j) \tilde{T},$$

$$\sqrt{\epsilon} [i(R - 1) - j \tilde{R}] = -(1 + j \gamma) T i \alpha_- + (\beta + j) \tilde{T} \alpha_+. \quad (11)$$

Multiplying the first equation for $\sqrt{\epsilon}$ and then summing it with the second equation, we find

$$\sqrt{\epsilon} [1 + R + i(R - 1)] = (1 + j \gamma) T (\sqrt{\epsilon} - i \alpha_-) + (\beta + j) \tilde{T} (\sqrt{\epsilon} + \alpha_+).$$

The left-hand side of this equation is complex. Consequently, the pure quaternionic part of the right-hand side of the equation has to be zero. This immediately implies the following relation between T and \tilde{T} ,

$$\tilde{T} = -\gamma \frac{\sqrt{\epsilon} - i \alpha_-}{\sqrt{\epsilon} + \alpha_+} T.$$

By using this relation in Eqs. (11) and taking the ratio of the complex parts, after simple algebraic manipulations, we find the following reflection coefficient

$$R = \frac{(\sqrt{\epsilon} + \alpha_+)(\sqrt{\epsilon} - \alpha_-) - \beta \gamma (\sqrt{\epsilon} - i \alpha_-)(\sqrt{\epsilon} - i \alpha_+)}{(\sqrt{\epsilon} + \alpha_+)(\sqrt{\epsilon} + \alpha_-) - \beta \gamma (\sqrt{\epsilon} - i \alpha_-)(\sqrt{\epsilon} + i \alpha_+)}. \quad (12)$$

We now explicitly calculate the phase for each of the two cases which characterize total reflection. Let us begin with $v_q < \epsilon < 1$. In this case, we have $(\beta \gamma)^* = \beta \gamma$, $\alpha_-^* = -\alpha_-$ and $\alpha_+^* = \alpha_+$. This implies

$$R_{>}(v_q) := R(v_q < \epsilon < 1) = \exp [2i \theta_{>}(v_q)],$$

where

$$\theta_{>}(v_q) = \arctan \left[\frac{\beta \gamma \alpha_+ (\sqrt{\epsilon} - i \alpha_-) + i \alpha_- (\sqrt{\epsilon} + \alpha_+)}{\sqrt{\epsilon} (\sqrt{\epsilon} + \alpha_+) - \beta \gamma \sqrt{\epsilon} (\sqrt{\epsilon} - i \alpha_-)} \right]. \quad (13)$$

The case $0 < \epsilon < v_q$ is a little bit more complicated. Let us first observe that $\beta \gamma = \exp [-2i \omega]$ with $\tan \omega = \sqrt{v_q^2 - \epsilon^2} / \epsilon$ and recall that in this case $\alpha_+ = i \alpha_-^*$. We then find

$$R_{<}(v_q) := R(0 < \epsilon < v_q) = -\exp [2i \theta_{<}(v_q)],$$

where

$$\theta_{<}(v_q) = \arctan \left(\frac{\epsilon \sin \omega + \sqrt{\epsilon} \operatorname{Im} [i \alpha_-^* e^{i \omega}]}{|\alpha_-|^2 \sin \omega - \sqrt{\epsilon} \operatorname{Re} [\alpha_- e^{i \omega}]} \right). \quad (14)$$

The phase for a *pure complex potential* (standard quantum mechanics) is obtained from Eq. (13) by setting $v_q = 0$,

$$\theta_c := \theta_{>}(0) = -\arctan \sqrt{\frac{1 - \epsilon}{\epsilon}} \quad (15)$$

For a *pure quaternion potential*, by using Eq. (14) for $v_q = 1$, we find

$$\theta_q := \theta_<(1) = \arctan \left(\frac{2\alpha\epsilon + 2\alpha^2\sqrt{\epsilon} + \epsilon^{3/2}}{4\alpha^3 + 2\alpha^2\sqrt{\epsilon} - \epsilon^{3/2}} \right). \quad (16)$$

IV. TOTAL REFLECTION: DELAY TIME CALCULATION

The quaternionic wave function which determines the particle dynamics in region I, $\xi < 0$, is given by

$$\varphi_I(\xi, \tau) = \left(\underbrace{\exp[i\sqrt{\epsilon}\xi]}_{\varphi_{1,inc}} + \underbrace{R \exp[-i\sqrt{\epsilon}\xi] + j\tilde{R} \exp[\sqrt{\epsilon}\xi]}_{\varphi_{1,ref}} \right) \exp[-i\epsilon\tau], \quad (17)$$

where $\tau = V_0 t/\hbar$. As observed in the introduction, for sufficiently large times the incident wave packet ($\varphi_{1,inc}$) disappears and the reflected wave packet ($\varphi_{1,ref}$) propagates towards the left. Observing that the pure quaternionic part decreases exponentially, the stationary phase method can be directly applied to the complex part of the reflected quaternionic wave function,

$$\left[\frac{d}{d\epsilon} (2\theta - \sqrt{\epsilon}\xi_r - \epsilon\tau) \right]_0 = 0.$$

The *delay time* is then given by

$$\tau_0 = 2 \left[\frac{d\theta}{d\epsilon} \right]_0. \quad (18)$$

From this equation, by using the phase given in Eq. (15), we obtain the *standard* delay time for complex potentials, i.e.

$$\tau_{0,c}[\epsilon_{0,c}] = 2 \left[\frac{d\theta_c}{d\epsilon} \right]_0 = \frac{1}{\sqrt{\epsilon_{0,c}(1 - \epsilon_{0,c})}}, \quad (19)$$

where $\epsilon_{0,c} = E_0/V_1$. The dependence of $\tau_{0,c} = V_1 t_0/\hbar$ upon the ratio between the incoming energy, E_0 , and the complex potential, V_1 , is plotted in Fig. 1 (continuous line). The minimum in the plane $\epsilon_{0,c} - \tau_{0,c}$ is found at

$$\left\{ \frac{\tilde{E}_{0,c}}{V_1}, \frac{V_1 \tilde{t}_{0,c}}{\hbar} \right\} = \{0.5, 2\}. \quad (20)$$

Consequently,

$$\left\{ \frac{\tilde{E}_0 \tilde{t}_0}{\hbar} \right\}_c = 1. \quad (21)$$

For a *pure quaternionic potential*, we have to use the phase given in Eq. (16). The expression for the quaternionic delay time is a little bit more complicated,

$$\tau_{0,q}[\epsilon_{0,q}] = 2 \left[\frac{d\theta_q}{d\epsilon} \right]_0 = \frac{2 + \frac{1}{\sqrt{2\epsilon_{0,q}}(1 - \epsilon_{0,q}^2)^{3/4}} + \frac{2\epsilon_{0,q}}{(1 - \epsilon_{0,q}^2)^{1/2}} + \frac{2\sqrt{2\epsilon_{0,q}}}{(1 - \epsilon_{0,q}^2)^{1/4}}}{\epsilon_{0,q} + \sqrt{2\epsilon_{0,q}}(1 - \epsilon_{0,q}^2)^{1/4} + (1 - \epsilon_{0,q}^2)^{1/2}}, \quad (22)$$

where $\epsilon_{0,q} = E_0/V_q$. The plot of $\tau_{0,q} = V_q t_0/\hbar$ as a function of the ratio between the incoming energy, E_0 , and the modulus of the pure quaternionic potential, V_q , is shown in Fig. 1 (dotted line).




FIG. 1. Delay times for complex ($\tau_{0,c}$) and pure quaternionic ($\tau_{0,q}$) potentials in terms of the adimensional incoming energies \tilde{E}_0/V_1 ($\epsilon_{0,c}$) and E_0/V_q ($\epsilon_{0,q}$).

Due to the fact that the minimum of the quaternionic delay time,

$$\left\{ \frac{\tilde{E}_{0,q}}{V_q}, \frac{V_q \tilde{t}_{0,q}}{\hbar} \right\} = \{0.365, 2.763\}, \quad (23)$$

is different from the complex case, it seems simple to recognize a pure quaternionic potential by calculating the quantity $\tilde{E}_0 \tilde{t}_0 / \hbar$. Surprisingly,

$$\left\{ \frac{\tilde{E}_0 \tilde{t}_0}{\hbar} \right\}_q \approx 1.0085, \quad (24)$$

which is very close to the result found for the complex case. Thus, an experiment involving this measurement does not represent the best choice to see quaternionic potentials.

V. CONCLUSIONS

As observed in Sec. IV, a calculation of $\tilde{E}_0 \tilde{t}_0 / \hbar$ done by two observers which are, respectively, working with a complex and a pure quaternionic potential step practically gives the same result. In looking for the energy region which could amplify the difference between complex and quaternionic quantum mechanics, it is interesting to introduce new energy and time variables defined in terms of the incoming energy, \tilde{E}_0 , for which we have a minimal delay time. In terms of these new variables,

$$\tilde{\epsilon}_0 = E_0 / \tilde{E}_0 \quad \text{and} \quad \tilde{\tau}_0 = \tilde{E}_0 t_0 / \hbar,$$




FIG. 2. Delay times for complex and pure quaternionic potentials in terms of the adimensional incoming energy E_0 / \tilde{E}_0 . The difference between quaternionic and complex quantum mechanics is amplified for incoming energies closed to $2 \tilde{E}_0$.

TABLE I. Reflection delay times for a pure quaternionic potential of height $V_q = 20$ keV (second column). The complex potentials (third, fourth, and fifth columns) only mimic the quaternionic potential for particular incoming energies (starred entries). Consequently, a complex potential cannot reproduce the quaternionic results.

E_0/keV	$V_q = 20$ keV keV t_0/\hbar	$V_1 = 18.5$ keV keV t_0/\hbar	$V_1 = 16.5$ keV keV t_0/\hbar	$V_1 = 14.6$ keV keV t_0/\hbar
2	0.174	★ 0.174	0.186	0.199
4	0.147	0.131	0.141	0.154
6	0.139	0.115	0.126	0.140
8	0.138	0.109	0.121	★ 0.138
10	0.142	0.108	0.124	0.147
12	0.152	0.113	0.136	0.179
14	0.169	0.126	★ 0.169	0.345

we find

$$\begin{aligned}\epsilon_{0,c} &= \frac{\tilde{E}_0}{V_1} \cdot \frac{E_0}{\tilde{E}_0} = 0.5 \tilde{\epsilon}_0, \\ \tilde{\tau}_{0,c} &= \frac{\tilde{E}_0}{V_1} \cdot \tau_{0,c}[0.5 \tilde{\epsilon}_0] = 0.5 \tau_{0,c}[0.5 \tilde{\epsilon}_0],\end{aligned}\quad (25)$$

and

$$\begin{aligned}\epsilon_{0,q} &= \frac{\tilde{E}_0}{V_q} \cdot \frac{E_0}{\tilde{E}_0} = 0.365 \tilde{\epsilon}_0, \\ \tilde{\tau}_{0,q} &= \frac{\tilde{E}_0}{V_q} \cdot \tau_{0,q}[0.365 \tilde{\epsilon}_0] = 0.365 \tau_{0,q}[0.365 \tilde{\epsilon}_0].\end{aligned}\quad (26)$$

Figure 2 clearly shows that the best choice to amplify the difference between complex and pure quaternionic potentials is achieved for incoming energies closed to $2\tilde{E}_0$ (this energy value represents the limit case between tunneling and diffusion for complex potentials). It is also interesting to observe that a complex potential *cannot* completely mimic a pure quaternionic potential. Indeed, as shown in Fig. 2 due to the different shapes of the delay time curves for complex and pure quaternionic potentials, a complex potential can perfectly mimic a pure quaternionic potential at most in two cases (see intersections between the two curves shown in Fig. 2). In Table I, we explicitly shown the delay times for reflection as function of the incoming energy (first column) for a pure quaternionic potential of height 20 keV (second column). In such a table, we also find the reflection delay times for complex potentials of height 18.5 keV (third column), 16.5 keV (fourth column) and 14.5 keV (fifth column). These complex potentials only mimic the quaternionic potential, respectively, for incoming energy of 2, 14, and 8 keV. This clearly shows that the reflection delay times obtained in presence of a quaternionic potential step cannot be reproduced by a complex potential step. The energy dependence of the delay times in reflection by a step potential can be then used to determine whether the step as a quaternionic nature.

The study presented in this paper represents a preliminary analysis of quaternionic delay times. Further investigations imply a generalization from plane waves to wave packets and/or from one-dimensional to three-dimensional problems.

ACKNOWLEDGMENTS

The authors thank an anonymous referee for his interesting comments and useful suggestions. One of the authors (S.d.L.) also thanks the CNPq (*Bolsa de Produtividade em Pesquisa*2010/13) and the FAPESP (Grant No. 10/02213-3) for financial support.

- ¹ A. J. Davies and B. H. McKellar, *Phys. Rev. A* **40**, 4209 (1989).
- ² A. J. Davies and B. H. McKellar, *Phys. Rev. A* **46**, 3671 (1992).
- ³ S. De Leo and G. Sclarici, *J. Phys. A* **33**, 2971 (2000).
- ⁴ S. De Leo and G. Ducati, *J. Math. Phys.* **42**, 2236 (2001).
- ⁵ S. De Leo, G. Sclarici, and L. Solombrino, *J. Math. Phys.* **43**, 5812 (2002).
- ⁶ S. De Leo, G. Ducati, and C. Nishi, *J. Phys. A* **35**, 5411 (2002).
- ⁷ S. De Leo and G. Ducati, *J. Math. Phys.* **44**, 2224 (2003).
- ⁸ S. De Leo and G. Ducati, *Comput. Math. Appl.* **48**, 1893 (2004).
- ⁹ S. De Leo and G. Ducati, *J. Math. Phys.* **47**, 082106 (2006).
- ¹⁰ S. De Leo, G. Ducati, V. Leonardi, and K. Pereira, *J. Math. Phys.* **51**, 113504 (2010).
- ¹¹ S. L. Adler, *Quaternionic Quantum Mechanics and Quantum Fields* (Oxford University Press, New York, 1995).
- ¹² S. De Leo and G. Ducati, *J. Phys. A* **38**, 3443 (2005).
- ¹³ S. De Leo and G. C. Ducati, *J. Math. Phys.* **47**, 102104 (2006).
- ¹⁴ S. De Leo and G. C. Ducati, *J. Math. Phys.* **48**, 052111 (2007).
- ¹⁵ C. Cohen-Tannoudji, B. Diu and F. Lalöe, *Quantum Mechanics* (Wiley, New York, 1977).
- ¹⁶ E. Merzbacher, *Quantum Mechanics* (Wiley, New York, 1970).