

Original article (full paper)

Association of inflammation, dyslipidemia, obesity and physical activity status in children

Juliano Magalhães Guedes

José Bontempo Mamêde Neto

Alyne Christian Ribeiro Andaki

Universidade Federal do Triângulo Mineiro, Uberaba, MG, Brasil

Patrícia Feliciano Pereira

Michelle Dias de Oliveira

Roberto Sousa Dias

Sérgio Oliveira de Paula

Antônio José Natali

Universidade Federal de Viçosa, Viçosa, MG, Brasil

Alexandre de Paula Rogério

Edmar Lacerda Mendes

Universidade Federal do Triângulo Mineiro, Uberaba, MG, Brasil

Abstract—The aim of this study was to verify the association between inflammatory biomarkers, dyslipidemia, obesity and physical activity status in 10-years old children. Ninety-four children participated in this study and were classified into eutrophic (n=36), overweight (n=34) or obese (n=24) according to their body mass index (BMI). The genic expression of interleukin 6 (IL-6), tumor necrosis factor alpha (TNF- α) and chemokine C-C motif ligand 2 (CCL-2) mRNA; the serum concentration of high-density lipoprotein cholesterol (HDL-c) and triglycerides; BMI, percentage of body fat (% BF) and waist circumference; and the number of steps per day were determined. The expression of IL-6, TNF- α and CCL-2 were associated ($p < 0.05$) positively with serum triglycerides, BMI, % BF and waist circumference, and negatively with serum HDL-c. No association ($p > 0.05$) between pro-inflammatory biomarkers and number of steps per day was found.

Keywords: child, inflammatory biomarkers, physical activity status, obesity

Introduction

Over 2.8 billion people die in consequence of overweight and obesity worldwide (World Health Organization [WHO], 2014). In Brazil, obesity triggers serious public health problems and demands high social costs (Duncan *et al.*, 2012). Inflammation plays a key role in the pathophysiology of obesity (Geraldo & Alfenas, 2008). The white adipocytes exhibit altered physiology due to excessive fat storage and release of pro-inflammatory cytokines such as interleukin 6 (IL-6), tumor necrosis factor alpha (TNF- α) and chemokine C-C motif ligand 2 (CCL-2), also referred to as monocyte chemoattractant protein-1 (Leite, Rocha, & Brandão-Neto, 2009). Such framework results in a state of chronic low-grade inflammation (Calder *et al.*, 2011; Herder *et al.*, 2007; Tam, Clement, Baur, & Tordjman, 2010).

Chronic low-grade inflammation is directly related to health problems such as insulin resistance, dyslipidemia, type 2 diabetes and atherosclerosis in obese children (Codoner-Franch *et al.*, 2011; Elshorbagy, Valdivia-Garcia, Refsum, & Butte, 2012; Garanty-Bogacka *et al.*, 2011; Kim *et al.*, 2010). Indeed, a recent

study showed that body mass index (BMI) and fasting insulin correlated positively with insulin resistance, serum lipids, and pro-inflammatory biomarkers, but negatively with adiponectin, an anti-inflammatory mediator (Chang *et al.*, 2015). Despite the positive associations between obesity and chronic low-grade inflammation, the expression of IL-6, TNF- α and CCL-2 in obese children shows conflicting results (Breslin *et al.*, 2012; Dixon, Meng, Goldberg, Schneiderman, & Delamater, 2009; Kim *et al.*, 2010; Maffei *et al.*, 2007).

Regular physical activity has been stimulated, in order, to promote a healthy lifestyle as well as to counteract overweight and obesity in the childhood. Indeed, regular physical activity reduces obesity, the production of pro-inflammatory biomarkers as well as the prevalence of cardiovascular diseases in children (Ben Ounis *et al.*, 2010; Chae *et al.*, 2010; Garanty-Bogacka *et al.*, 2011; Vos, Wit, Pijl, & Houdijk, 2011). To achieve the benefits of physical activity, it was proposed that 13000 to 15000 steps per day for boys and 11000 to 12000 steps per day for girls should be performed by 7 to 12-years old children (Tudor-Locke *et al.*, 2011). To our knowledge, no study tested the associations

between IL-6, TNF- α , CCL-2 with the number of steps per day in 10-years old children.

Taking into consideration that the prevalence of overweight and obesity in children and adolescents is high (Espin Rios, Perez Flores, Sanchez Ruiz, & Salmeron Martinez, 2013; Ferreira, 2006; Instituto Brasileiro de Geografia e Estatística (IBGE), 2004), the early diagnostic of the association of pro-inflammatory biomarkers with modifiable lifestyle factors are of interest as well as could preventing the outcomes of chronic diseases in the future. Therefore, we designed this cross-sectional study to examine the association between inflammatory biomarkers, dyslipidemia, obesity and physical activity status in 10-years old children.

Methods

Participants

Ninety-four children (44 boys and 50 girls) aged 10.03 ± 0.74 years (mean \pm SD) were selected and classified as eutrophic ($n = 19$ girls, $n = 17$ boys), overweight ($n = 18$ girls, $n = 16$ boys) and obese ($n = 13$ girls, $n = 11$ boys), according to their BMI. This study was approved by the Federal University of Triângulo Mineiro Research Ethics Committee (CEP/UFTM n° 2388) and authorized by the Municipal Secretary of Education and Culture of Uberaba, Minas Gerais, Brazil (CRAFT GAB / SEMEC / No. 0898). All participants agreed to take part in the study, and either parents or legal guardians signed the statement of consent.

Anthropometric measures

Body mass index calculated using the formula [BMI = body mass (kg)/height (m²)]. Waist circumference (WC) was measured at the midpoint between the iliac crest and the last rib (WHO, 2005). Triceps (TS) and subscapular (SS) skinfolds were measured in triplicate using a caliper (Lange Skinfold Caliper, Cambridge, USA) according to Guedes (2006) protocol. The mean value of the three measures was used. The percentage of body fat percentage (% BF) was determined according to Slaughter *et al.* (1998). Boys and girls were classified as eutrophic, overweight and obese using BMI values according to Cole, Bellizzi, Flegal and Dietz (2000).

Physical activity status

The habitual physical activity status was determined by measuring the number of steps per day using the uni-axial motion sensor, pedometer (Yamax SW200 Digiwalker, Japan). After explanations and familiarization with the pedometer, the participants registered the number of steps for five consecutive days (Brusseau *et al.*, 2011). Participants were instructed to position the pedometer above the iliac crest in the midline of the right thigh, attached to an elastic waistband during all wake time. They were advised to perform their routine activities and remove the pedometer only when lying down, riding a bicycle, bathing or practicing water activities. The participants were told that the pedometer should be placed back to the waist

immediately after these activities. Children were instructed to record the number of steps shown on the pedometer on a paper given at the end of each day before lying down to sleep. On the next day, after wake up, children were told to reset the pedometer and verify if the display showed zero before placed back on the waist. For analyzes, the average of steps performed during the 5 days were considered. For classification the physical activity status, we adopted the cut-offs of 11000 steps/day for girls and 13000 steps/day for boys (Tudor-Locke *et al.*, 2011).

Serum lipid profile

After 12-14 hours fasting, blood samples were collected in appropriate tubes (BD Vacutainer®, São Paulo, Brazil) containing gel and clot activator. Serum, plasma and buffy-coat were separated from other blood components by centrifugation at 3,400 rpm for eight minutes and stored in Eppendorf tubes (1.5 mL) at -80°C. Serum triglycerides and HDL-c were analyzed by an enzymatic colorimetric method using a semiautomatic biochemical analyzer BIO 200F (Bioplus, São Paulo, Brazil) and commercial kits (LABTEST, System Diagnostics Ltda. Lagoa Santa, Brazil). The cut-offs used for dyslipidemia diagnosis were: triglycerides ≥ 150 mg / dL and HDL-c ≤ 35 mg / dL (Kavey *et al.*, 2014).

Gene expression of inflammatory biomarkers

The buffy-coat aqueous and organic phases were separate by adding 100 μ L of phosphate buffered saline (PBS) and 750 μ L of Trizol (Invitrogen - Sao Paulo, Brazil). After incubation at room temperature for five minutes, it was added 200 μ L of chloroform (Merck, Rio de Janeiro, Brazil) and centrifuged for 15 minutes at 4° C. The supernatant (aqueous phase RNA) was then transferred to another eppendorf tube in which was added 380 μ L of isopropanol. Subsequently, the samples were centrifuged for 4 minutes at 4° C and stored at -80° C for 12 hours to precipitate the RNA. Then, the precipitate was washed with 750 μ L of ethanol (75%) and stored for 10 minutes at room temperature to dry. Thereafter, it was added 50 μ L of ultrapure water, free RNase. The concentration of purified total RNA was quantified by spectrophotometer (Nanodrop 2000c, Wilmington, North Carolina - United States) at the concentrations of 260/280 ratio nM.

For the synthesis of c-DNA reverse transcription reaction was performed from total RNA purified. A mixture of 1 g of RNA, 1 ml of enzyme DNase and 12 μ l of ultrapure H₂O was homogenized and incubated at 65° C for 5 minutes and, subsequently, cooled on ice for 1 minute. Then, it was added 2 μ l of physiological buffer (10X concentrate), 2 μ l of 10 mM dNTP Mix, 1 ml of the enzyme reverse transcriptase (Vivantis) and 2 μ l of ultrapure H₂O. The samples were incubated again at 42° C for 60 minutes.

For the analysis of IL-6, TNF- α and CCL-2, the ECO Real Time PCR was carried out (Polymerase Chain Reaction System, Uniscience, São Paulo - Brazil).

For amplification of inflammatory biomarkers gene, it was added in a solution of 0.4 µL of c-DNA 400 nm of each primer (IL-6, TNF-α and CCL-2) and 10 µL of EVAgreen master mix (Invitrogen - Sao Paulo, Brazil). The gene expressions of IL-6, TNF-α and CCL-2 were demonstrated in values of standard dissociation curve obtained through relative endogenous control gene β-actin. The sequence of the primers is shown in Table 1.

Table 1. Primers used for the analysis of inflammatory biomarkers.

Biomarker	primer sense sequence primer anti-sense sequence
IL-6	TCCAGTTGCCTTCTTGGGAC GTACTCCAGAAGACCAGAGG
TNF - α	AAGCCTGTAGCCCATGTTGT CAGATAGATGGGCTCATACC
CCL2	AGGAAGATCTCAGTG CAGAGG AGTCTTCGGAGTTTGCCTTTG
Beta-actin	ATGTTTGAGACCTTCAACAC CACGTCADACTTCATGATGG

Legend: IL-6 – Interleukin 6; TNF-α – Tumor necrosis factor alpha; CCL-2 – Chemokine (C-C motif.) ligand 2.

Statistical analysis

The Shapiro-Wilk test was used to test data normality. For comparisons, one-way analysis of variance (ANOVA) followed by post-hoc Tukey test was used. Pearson’s correlation test was

used to test associations between inflammatory biomarkers, triglycerides, HDL-c, BMI, % BF, waist circumference and physical activity status. We adopted confidence interval (CI) of 95% and significance level of $\alpha \leq 5\%$.

Results

Anthropometric measurements

The obese group had higher measures for body mass, BMI, waist circumference, TS, SS, Σ 2SF (TS + SS) and %BF as compared with eutrophic and overweight (Table 2). Likewise, the overweight group differed statistically from the eutrophic group in all anthropometric variables analyzed. No gender differences were found ($p > 0.05$) in all anthropometric measurements.

Physical activity status

The number of steps per day was not statistically different among eutrophic, overweight and obese children (Figure. 1). Likewise, no gender difference between groups was observed. The recommendation of 11000 and 13000 steps/day was not reached by 58.8% of girls and 83.6% of boys, respectively. Among these individuals, 42.2% were eutrophic, 32.8% overweight and 25% obese.

Table 2. Anthropometric measures and body composition.

	Eutrophic			Overweight			Obesity		
	Boys (n = 19)	Girls (n = 17)	Total (n=36)	Boys (n = 18)	Girls (n = 16)	Total (n=34)	Boys (n = 13)	Girls (n = 11)	Total (n=24)
Height (cm)	137 ± 10	137 ± 9	137 ± 9,5	137 ± 10	138 ± 8	137,5 ± 9	137 ± 9	137 ± 8	137 ± 8,5
BM (kg)	29,7 ± 2,0	29,3 ± 2,04	29,5 ± 2,03	39,1 ± 2,8	39,7 ± 3,4	39,4 ± 3,1*	53,9 ± 3,0	53,7 ± 3,6	53,8 ± 3,3**
BMI (kg/m ²)	16,1 ± 1,7	15,8 ± 1,8	15,9 ± 1,7	20,2 ± 1,1	21,6 ± 0,9	20,9 ± 1,0*	26,4 ± 1,4	27,1 ± 2,0	26,9 ± 1,7**
WC (cm)	58,1 ± 1,9	56,9 ± 1,1	57,7 ± 2,6	65 ± 1,5	63 ± 1,9	64 ± 2,4*	79 ± 3,2	77 ± 2,6	78 ± 5,9**
TS (mm)	10,9 ± 3,8	12,1 ± 4,0	11,08 ± 3,9	19,1 ± 4,0	20,1 ± 4,4	19,6 ± 4,2*	27 ± 3,9	27,6 ± 5,1	27,3 ± 4,5**
SS (mm)	8 ± 3,3	8,4 ± 4,3	8,2 ± 3,8	16,2 ± 6	16,6 ± 6,4	16,4 ± 6,2*	29,6 ± 5,0	30,4 ± 6,4	30 ± 5,8**
Σ 2SF (mm)	18,9 ± 7,0	20,5 ± 7,6	20 ± 7,3	35,3 ± 7,1	36,7 ± 13,1	36,04 ± 10,1*	56,6 ± 8,2	60 ± 9,8	57,3 ± 9,0**
BF (%)	20,9 ± 7,5	21,1 ± 7,9	21 ± 7,7	32,1 ± 4,0	32,9 ± 5,4	32,5 ± 4,7*	41,9 ± 3,4	43,7 ± 4,4	42,8 ± 3,9**

Data are expressed as mean ± Standard deviation (SD). Legend: BM = body mass; BMI = Body mass index; WC = Waist circumference; TS = Triceps skinfold; SS = Subscapular skinfold; Σ2SF= Sum of two skinfold; BF = Body fat; Test: Analysis of variance (ANOVA) plus Tukey.

* Significantly different from eutrophic individuals ($p < .05$);

† Significantly different from overweight individuals ($p < .05$).

Lipid profile

The concentration of HDL-c was higher in eutrophic children as compared with overweight and obese (Figure. 2A). However, no differences were found ($p < 0.05$) in HDL-c between overweight and obese children. Obese children had higher ($p > 0.05$) triglycerides concentrations as compared with eutrophic and overweight (Figure. 2B).

Gene expression of inflammatory biomarkers

The genic expressions of IL-6 (Figure. 3A), TNF-α (Figure. 3B) and CCL-2 (Figure. 3C) were higher in obese children as compared with eutrophic group. However, no statistical difference in the mRNA gene expressions of these biomarkers was found between overweight and obese group.




Figure 1. Number of steps per day. Data expressed as mean \pm standard deviation. Eutrophic (n=19 boys and 17 girls). Overweight (n=18 boys and 16 girls). Obese (13 boys and 11 girls). *, significantly different from eutrophic individuals ($p < .05$)




Figure 2. Serum lipid profile. (A) High-density lipoprotein cholesterol (HDL-c). (B) Triglycerides. Data expressed as mean \pm standard deviation. Eutrophic (n = 36). Overweight (n = 34). Obese (n = 24).*, significantly different from eutrophic individuals ($p < 0.05$).




Figure 3. Genic expression of inflammatory biomarkers. (A) Tumor necrosis factor alpha (TNF- α). (B) Interleukin 6 (IL-6). (C) Chemokine C-C motif ligand 2 (CCL-2). Data are expressed as mean \pm standard deviation. Eutrophic (n = 36). Overweight (n = 34). Obese (n = 24). *, significantly different from eutrophic individuals ($p < 0.05$).

Associations analysis

Table 3 shows the correlations analysis. IL-6, TNF- α and CCL-2 presented strong, positive and statistically significant associations with BMI and waist circumference. In addition, IL-6, TNF- α and CCL-2 showed moderate, positive and

statistically significant associations with triglycerides and %BF; but moderate, negative and statistically significant association with HDL-c. Moreover, there was a weak, but not statistically significant association between inflammatory biomarkers and the number of steps per day. However, there was no gender difference in these correlation analysis.

Table 3. Correlation between inflammatory biomarkers, triglycerides and HDL-c, anthropometric measures, body composition and physical activity status.

	IL- 6			TNF- α			CCL- 2		
	Boys (n = 19)	Girls (n = 17)	Total (n=36)	Boys (n = 18)	Girls (n = 16)	Total (n=34)	Boys (n = 13)	Girls (n = 11)	Total (n=24)
IL-6	0.894*	0.913*	1*	0.304*	0.473*	0.929*	0.378*	0.478*	0.884*
TNF- α	0.304*	0.473*	0.929*	0.918*	0.944*	1*	0.421*	0.437*	0.908*
Triglycerides	0.394*	0.397 *	0.377 *	0.368 *	0.392 *	0.407 *	0.321 *	0.363 *	0.398 *
HDL-c	- 0.306*	- 0.394 *	- 0.324 *	- 0.412 *	- 0.394 *	- 0.348 *	- 0.385 *	- 0.348 *	- 0.359 *
Body Mass Index	0.626 *	0.785 *	0.718 *	0.669 *	0.809 *	0.735 *	0.677 *	0.841 *	0.784 *
Waist circumference	0.876*	0.815*	0.833*	0.894*	0.795*	0.851*	0.860*	0.776*	0.813*
Percentage of body fat	0.586 *	0.720 *	0.620 *	0.648 *	0.743 *	0.593 *	0.655 *	0.767 *	0.592 *
Steps per day	0.045	- 0.029	0.046	0.191	0.038	0.046	0.128	- 0.036	0.052

IL-6 = Interleukin 6; TNF- α = Tumor necrosis factor alpha; HDL-c = High-density lipoprotein cholesterol; CCL2= chemokine C-C motif ligand 2 ; * = statistically significant ($p < 0.05$).

Discussion

In the present study, we tested the association of inflammatory biomarkers with dyslipidemia, obesity and physical activity status in 10 years-old children. Our data showed that the genic expressions of IL-6, TNF- α , CCL-2 and serum triglycerides concentrations were higher in obese than in eutrophic and overweight children while serum HDL-c was higher in eutrophic children. In addition, it was observed that these pro-inflammatory biomarkers were positively associated with triglycerides, %BF, BMI and waist circumference; but negatively associated with HDL-c. Nevertheless, no association of these biomarkers with physical activity status was found.

Positive association of dyslipidemia (i.e. LDL-C, VLDL-C and triglycerides) with IL-6 (Alvarez *et al.*, 2009; Bugge *et al.*, 2012a; Chirico *et al.*, 2013; Elshorbagy *et al.*, 2012; Garanty-Bogacka *et al.*, 2011; Kim *et al.*, 2010; Pyrzak *et al.*, 2009; Roth, Kratz, Ralston, & Reinehr, 2011) and TNF- α (Breslin *et al.*, 2012; Bugge *et al.*, 2012a; Codoner-Franch *et al.*, 2011; Dixon *et al.*, 2009; Jermendy *et al.*, 2010; Kim *et al.*, 2010; Pyrzak, Wisniewska, Popko, Demkow, & Kucharska, 2010; Roth *et al.*, 2011) in children have been demonstrated by others. However, either positive (Breslin *et al.*, 2012; Roth *et al.*, 2011) or negative (Kim *et al.*, 2010; Wasilewska, Tenderenda, Taranta-Janusz, Tobolczyk, & Stypulkowska, 2012) associations between CCL-2 and dyslipidemia has been reported.

High serum triglycerides and low HDL-c concentrations have been found in obese children which increased risk of cardiovascular disease in the adulthood (Duncan *et al.*, 2012; Friedemann *et al.*, 2012). According to Morrison (2003), serum cholesterol values are similar in boys and girls in the childhood, but in post-menarche girls the sex hormones, particularly estradiol, deeply influence the levels of HDL-c and LDL-c. cholesterol. Thus, it appears that sex hormones do not have great influences the relationship between inflammatory biomarkers and obesity in boys and girls before pubertal phase.

The associations of IL-6, TNF- α , CCL-2 with BMI, waist circumference and %BF observed in the present study support the global concern for the control of obesity, especially during childhood. An increase in pro-inflammatory biomarkers triggers the chronic low grade inflammation that is linked to cardiovascular disorders such as atherosclerosis (Dessi *et al.*, 2013; Gustafson, 2010; Poitou *et al.*, 2011), insulin resistance (Charles *et al.*, 2011; Elshorbagy *et al.*, 2012; Kim *et al.*, 2010) and diabetes (DeBoer, 2013; Donath & Shoelson, 2011; Sobti, Kler, Sharma, Talwar, & Singh, 2012). Therefore, obesity in the childhood can lead to chronic low-grade inflammation as well as increase of susceptibility to cardio metabolic diseases.

Concerning gender, it was observed that the genic expressions of pro-inflammatory biomarkers were not different between genders in 10-years old children. A longitudinal study by Bugge *et al.* (2012b) showed no significant differences between gender for IL-6, TNF- α and CCL-2 concentrations. However, for insulin resistance, the concentrations of IL-6 in 9-years old and 13-years old were associated with girls and not with boys.

Regarding obesity, positive associations between obesity, IL-6, TNF- α (Alvarez *et al.*, 2009; Tsaoussoglou *et al.*, 2010; Chirico *et al.*, 2013) and CCL-2 (Breslin *et al.*, 2012; Roth *et al.*,

2011) have been shown in 10-years old children. Nevertheless, no association of obesity with IL-6, TNF- α (Andersen *et al.*, 2010; Dixon, Meng, Goldberg, Schneiderman & Delamater, 2009) and CCL-2 (Elshorbagy *et al.*, 2012; Kim *et al.*, 2010) was found by others in this population.

Our data showed that the number of steps per day did not differ significantly between boys and girls and among eutrophic, overweight and obese children. The study by Rosa *et al.* (2011) found no association of the number of steps per day with BMI and %BF in children and adolescents 10 to 18-years old. However, higher number of steps per day in boys than in girls was reported by others (Raustorp, Svenson, & Perlinger, 2007; Tudor-Locke, Ainsworth, & Popkin, 2001). It seems that boys generally have higher family support and encouragement to practice moderate to vigorous physical activities than girls (Goncalves, Hallal, Amorim, Araujo, & Menezes, 2007). Therefore, the family can be instrumental in influencing the physical activity status in the childhood. It is noteworthy that in the present study more than 50% of girls and 80% of boys did not reach the recommended 11000 and 13000 steps per day, respectively. In this sense, individuals studied here exhibited a low physical activity status.

Unexpectedly, our data showed that the number of steps day was not associated with the pro-inflammatory biomarkers. It has been reported that low physical activity status did not lead to decreased pro-inflammatory biomarkers, whereas moderate to vigorous physical activity status promoted reductions in pro-inflammatory and increases in anti-inflammatory biomarkers (Park, Schwarz, Willoughby & Koh, 2015). Thus, the absence of association between number of steps per day with pro-inflammatory biomarkers in the present study is consistent with the low physical status of the studied individuals. It is worthy to note that children with high physical activity status as well as low concentrations of pro-inflammatory biomarkers presents low risks for cardiovascular diseases such as obesity (Sobieska, Gajewska, Kalmus & Samborski, 2013; Seabra *et al.*, 2015).

The cut-off points for BMI classification were made according to Cole *et al.* (2000) and based on international studies. It is important to mention that BMI has been widely used in many studies with children in both national (Chiara, Sichieri, & Martins, 2003; Vieira, Alvarez, Marins, Sichieri, & Veiga, 2006; Vitolo, Campagnolo, Barros, Gama, & Lopez, 2007) and international investigations (Conde & Monteiro, 2006; Freedman *et al.*, 2004; Neovius, Linne, Barkeling, & Rossner, 2004; Zimmermann, Gubeli, Puntener, & Molinari, 2004). In addition, BMI shows strong associations with gold standard methods (Jimenez-Pavon, Kelly, & Reilly, 2010). According to WHO (1995), the use of BMI is appropriate inasmuch as it has good correlation with %BF and allows comparisons to other studies. However, the determination of specific cut-off points to classify the obesity degree is complex since there are physiological differences between populations and lack of validation for the adopted criteria (DaSilva, Lopes, & DaSilva, 2010).

Finally, our results have clinical relevance as the early detection of high concentrations of pro-inflammatory biomarkers and dyslipidemia in 10-years old children pointing out the need for the adoption of preventive strategies.

Study limitations: pedometers are limited devices to measure walking because they capture movement only of the lower body in the vertical plane and cannot distinguish the intensity of walking, load carriage as well as between walking on different surfaces or gradients. This cross-sectional study design does not allow to affirm that the reported associations are causal.

References

- Alvarez, J.A., Higgins, P.B., Oster, R.A., Fernandez, J.R., Darnell, B.E., & Gower, B.A. (2009). Fasting and postprandial markers of inflammation in lean and overweight children. *The American Journal of Clinical Nutrition*, 89(4).
- Andersen, L.B., Muller, K., Eiberg, S., Froberg, K., Andersen, J.F., Bugge, A., Hermansen, B., McMurray, R.G. (2010). Cytokines and clustered cardiovascular risk factors in children. *Metabolism*, 59(4), 561-566.
- Ben Ounis, O., Elloumi, M., Zouhal, H., Makni, E., Denguezli, M., Amri, M., Lack, G., Tabka, Z. (2010). Effect of individualized exercise training combined with diet restriction on inflammatory markers and IGF-1/IGFBP-3 in obese children. *Annals of Nutrition and Metabolism*, 56(4), 260-266.
- Breslin, W.L., Johnston, C.A., Strohacker, K., Carpenter, K.C., Davidson, T.R., Moreno, J.P., Foreyt, J.P., McFarlin, B.K. (2012). Obese Mexican American children have elevated MCP-1, TNF-alpha, monocyte concentration, and dyslipidemia. *Pediatrics*, 129(5), e1180-1186.
- Brusseau, T.A., Kulinna, P.H., Tudor-Locke, C., Ferry, M., van der Mars, H., & Darst, P. W. (2011). Pedometer-determined segmented physical activity patterns of fourth- and fifth-grade children. *The Journal of Physical Activity and Health*, 8(2), 279-286.
- Bugge, A., El-Naaman, B., McMurray, R.G., Froberg, K., Nielsen, C.H., Muller, K., Andersen, L.B. (2012a). Inflammatory markers and clustered cardiovascular disease risk factors in Danish adolescents. *Hormone Research in Paediatrics*, 78(5-6), 288-296.
- Bugge, A., El-Naaman, B., McMurray, R.G., Froberg, K., Nielsen, C. H., Muller, K., & Andersen, L. B. (2012b). Sex differences in the association between level of childhood interleukin-6 and insulin resistance in adolescence. *Experimental Diabetes Research*, 859186.
- Calder, P.C., Ahluwalia, N., Brouns, F., Buetler, T., Clement, K., Cunningham, K., Esposito, K., Jönsson, L.S., Kolb, H., Lansink, M., Marcos, A., Margioris, A., Matusheski, N., Nordmann, H., O'Brien, J., Pugliese, G., Rizkalla, S., Schalkwijk, C., Tuomilehto, J., Wärnberg, J., Watzl, B., Winkhofer-Roob, B.M. (2011). Dietary factors and low-grade inflammation in relation to overweight and obesity. *British Journal of Nutrition*, 106(3), S5-78.
- Chae, H.W., Kwon, Y.N., Rhie, Y.J., Kim, H.S., Kim, Y.S., Paik, I.Y. Suh, S.H., Kim, D.H. (2010). Effects of a structured exercise program on insulin resistance, inflammatory markers and physical fitness in obese Korean children. [Randomized Controlled Trial. *Journal of Pediatric Endocrinology & Metabolism* 23(10), 1065-1072.
- Chang, C.J., Jian, D.Y., Lin, M.W., Zhao, J.Z., Ho, L.T., Juan, C. (2015). Evidence in obese childre: Contribution of Hiperlipidemia, Obesity-Inflammation and Insulin Sensitivity. *Plos One*, 10(5), E0125935.doi: 10.1371.
- Charles, B.A., Doumatey, A., Huang, H., Zhou, J., Chen, G., Shriner, D. (2011). The roles of IL-6, IL-10, and IL-1RA in obesity and insulin resistance in African-Americans. *The Journal of Clinical Endocrinology & Metabolism*, 96(12), E2018-2022. doi: 10.1210/jc.2011-1497.
- Chiara, V., Sichiari, R., & Martins, P. D. (2003). Sensibilidade e especificidade de classificação de sobrepeso em adolescentes. *Revista de Saúde Pública*, 37(2), 226-231.
- Chirico, V., Cannavo, S., Lacquaniti, A., Salpietro, V., Mandolino, M., Romeo, P.D., Cotta, O., Munafò, C., Giorgianni, G., Salpietro, C., Arrigo, T. (2013). Prolactin in obese children: a bridge between inflammation and metabolic-endocrine dysfunction. *Clinical Endocrinology*, 79(4), 537-44.
- Codoner-Franch, P., Tavares-Alonso, S., Murria-Estal, R., Megias-Vericat, J., Tortajada-Girbes, M., & Alonso-Iglesias, E. (2011). Nitric oxide production is increased in severely obese children and related to markers of oxidative stress and inflammation. *Atherosclerosis*, 215(2), 475-480.
- Cole, T.J., Bellizzi, M.C., Flegal, K.M., & Dietz, W.H. (2000). Establishing a standard definition for child overweight and obesity worldwide: international survey. *British Medical Journal*, 320(7244), 1240-1243.
- Conde, W.L., & Monteiro, C.A. (2006). Valores críticos do índice de massa corporal para classificação do estado nutricional de crianças e adolescentes brasileiros. *Journal of Pediatrics* 82(4), 266-272.
- DaSilva, K.S., Lopes, A.S., & DaSilva, F.M. (2010). Sensibilidade e especificidade de diferentes critérios de classificação do excesso de peso em escolares de João Pessoa, Paraíba, Brasil. *Revista de Nutrição, Campinas*, 23(1), 27-35.
- DeBoer, M.D. (2013). Obesity, systemic inflammation, and increased risk for cardiovascular disease and diabetes among adolescents: a need for screening tools to target interventions. *Nutrition*, 29(2), 379-386.
- Dessi, M., Noce, A., Bertucci, P., Manca di Villahermosa, S., Zenobi, R., Castagnola, V. (2013). Atherosclerosis, Dyslipidemia, and Inflammation: The Significant Role of Polyunsaturated Fatty Acids. [Review]. *International Scholarly Research Network Inflammation*, 191823.
- Dixon, D., Meng, H., Goldberg, R., Schneiderman, N., & Delamater, A. (2009). Stress and body mass index each contributes independently to tumor necrosis factor-alpha production in prepubescent Latino children. *Journal of Pediatric Nursing*, 24(5), 378-388.
- Donath, M.Y., & Shoelson, S.E. (2011). Type 2 diabetes as an inflammatory disease. *Nature Reviews Immunology*, 11(2), 98-107.
- Elshorbagy, A. K., Valdivia-Garcia, M., Refsum, H., & Butte, N. (2012). The association of cysteine with obesity, inflammatory cytokines and insulin resistance in Hispanic children and adolescents. *PLoS One*, 7(9), e44166.
- Espin Rios, M.I., Perez Flores, D., Sanchez Ruiz, J. F., & Salmeron Martinez, D. (2013). Prevalence of childhood obesity in the Murcia Region; an assessment of different references for body mass index. *Anales de Pediatria (Barc)*, 78(6), 374-381.
- Ferreira, A.P. (2006). Síndrome metabólica e fatores de risco cardiovascular em crianças pré-púberes de diferentes classificações nutricionais e níveis de resistência à insulina. *Brasília: Universidade Católica de Brasília*.

- Freedman, D.S., Thornton, J.C., Mei, Z., Wang, J., Dietz, W. H., Pierson, R.N., Jr. (2004). Height and adiposity among children. *Obesity Research*, 12(5), 846-853.
- Friedemann, C., Heneghan, C., Mahtani, K., Thompson, M., Perera, R., Ward, A.M. (2012). Cardiovascular disease risk in healthy children and its association with body mass index: systematic review and meta-analysis. *British Medical Association*, 345(e4759), 1-116.
- Garanty-Bogacka, B., Syrenicz, M., Goral, J., Krupa, B., Syrenicz, J., Walczak, M., & Syrenicz, A. (2011). Changes in inflammatory biomarkers after successful lifestyle intervention in obese children. *Endokrynologia Polska*, 62(6), 499-505.
- Geraldo, J.M., & Alfenas, R.C.G. (2008). Papel da dieta na prevenção e no controle da inflamação crônica - evidências atuais. *Arquivos Brasileiros de Endocrinologia & Metabologia*, 5(6), 951-967.
- Goncalves, H., Hallal, P.C., Amorim, T.C., Araujo, C.L., & Menezes, A.M. (2007). Sociocultural factors and physical activity level in early adolescence. *Revista Panamericana de Salud Publica*, 22(4), 246-253.
- Guedes, D. P. G., J. E.R.P. (2006). *Manual Prático para Avaliação em Educação Física*. Barueri-SP.
- Gustafson, B. (2010). Adipose tissue, inflammation and atherosclerosis. *Journal of Atherosclerosis and Thrombosis*, 17(4), 332-341.
- Herder, C., Schneitler, S., Rathmann, W., Haastert, B., Schneitler, H., Winkler, H., Bredahl, R., Hahnloser, E., Martin, S (2007). Low-grade inflammation, obesity, and insulin resistance in adolescents. *The Journal of Clinical Endocrinology & Metabolism*, 92(12), 4569-4574.
- Instituto Brasileiro de Geografia e Estatística. (2004). Pesquisas de orçamentos familiares, 2002-2003. Antropometria e análise do estado nutricional de crianças e adolescentes no Brasil. In: Ministério do Planejamento. *Rio de Janeiro*.
- Jermendy, A., Korner, A., Kovacs, M., Kaszas, E., Balazsovics, J., Szocs, A. (2010). Association between toll-like receptor polymorphisms and serum levels of tumor necrosis factor-alpha and its soluble receptors in obese children. *Medical Science Monitor*, 16(4).
- Jimenez-Pavon, D., Kelly, J., & Reilly, J.J. (2010). Associations between objectively measured habitual physical activity and adiposity in children and adolescents: Systematic review. *International Journal of Pediatric Obesity*, 5(1), 3-18.
- Kavey, R.W., Daniels, S.R., Lauer, R.M., Atkins, D.L., Hayman, L.L., & Taubert, K. (2014). American Heart Association Guidelines for Primary Prevention of Atherosclerotic Cardiovascular Disease Beginning in Childhood. *American Heart Association*, 107, 1562-1563.
- Kim, J., Bhattacherjee, R., Kheirandish-Gozal, L., Khalyfa, A., Sans Capdevila, O., Tauman, R. et al. (2010). Insulin sensitivity, serum lipids, and systemic inflammatory markers in school-aged obese and nonobese children. *International Journal of Pediatrics*, 2010, 846098.
- Leite, L. D., Rocha, E.D.D.M., & Brandão-Neto, J. (2009). Obesidade: uma doença inflamatória. *Revista Ciência & Saúde, Porto Alegre*, 2(2), 85-95.
- Maffei, C., Silvagni, D., Bonadonna, R., Grezzani, A., Banzato, C., & Tato, L. (2007). Fat cell size, insulin sensitivity, and inflammation in obese children. *The Journal of Pediatrics*, 151(6), 647-652.
- Morrison, J. A. (2003). A longitudinal evaluation of the NCEP-Peds guidelines for elevated total and LDL cholesterol in adolescent girls and boys. *Progress in Pediatric Cardiology*, 17, 159-168.
- Neovius, M.G., Linne, Y.M., Barkeling, B.S., & Rossner, S.O. (2004). Sensitivity and specificity of classification systems for fatness in adolescents. *Am J Clin Nutr*, 80(3), 597-603.
- Park, J., Schwarz, N., Willoughby, D., Koh, Y. (2015). Role of High-intensity Resistance Exercise in Vascular Inflammation in Recreationally Trained Men. *International Journal of Exercise Science*. 2(7), 7-65.
- Poitou, C., Dalmás, E., Renovato, M., Benhamo, V., Hajdúch, F., Abdennour, M. (2011). CD14dimCD16+ and CD14+CD16+ monocytes in obesity and during weight loss: relationships with fat mass and subclinical atherosclerosis. *Arteriosclerosis, Thrombosis and Vascular Biology*, 31(10).
- Pyrzak, B., Wisniewska, A., Majcher, A., Popko, K., Wasik, M., & Demkow, U. (2009). Association between metabolic disturbances and G-174C polymorphism of interleukin-6 gene in obese children. *European Journal of Medical Research*, 14 Suppl 4, 196-200.
- Pyrzak, B., Wisniewska, A., Popko, K., Demkow, U., & Kucharska, A. M. (2010). Association between anthropometric measures of obesity, metabolic disturbances and polymorphism G-308A of the tumor necrosis factor-alpha gene in children. *European Journal of Medical Research*, 15 Suppl 2, 141-146.
- Raustorp, A., Svenson, K., & Perlinger, T. (2007). Tracking of pedometer-determined physical activity: a 5-year follow-up study of adolescents in Sweden. *Pediatric Exercise Science*, 19(2), 228-238.
- Rosa, C.S.C., Messias, K.P., Fernandes, R.A., DaSilva, C.B., Monteiro, H. L., & Júnior, I. F.F. (2011). Atividade física habitual de crianças e adolescentes mensurada por pedômetro e sua relação com índices nutricionais. *Revista Brasileira de Cineantropometria & Desempenho Humano*, 13(1), 22-28.
- Roth, C.L., Kratz, M., Ralston, M.M., & Reinehr, T. (2011). Changes in adipose-derived inflammatory cytokines and chemokines after successful lifestyle intervention in obese children. *Metabolism*, 60(4), 445-452.
- Seabra, A.,Katzmarzyk, P., Facsm., Carvalho, M. J., Seabra, A.,Coelho-E-Silva, Abreu, S., Vale, S., Nascimento, H., Belo, L., Torres, S., Oliveira, J., Mota, J., Santos-Sila, A., Rêgo, C., Malina, M. (2015). Effects Of 6-month Soccer And Traditional Physical Activity Programs On Body Composition, Cardiometabolic, Inflammatory And Oxidative Markers In Obese Boys. *62nd American College of Sports Medicine*, 216(1), 1423.
- Slaughter, M.H., Lohman, T.G., Boileau, R.A., Horswill, C.A., Stillman, R.J., Van Loan, M. (1988). Skinfold equations for estimation of body fatness in children and youth. *Human Biology*, 60(5), 709-723.
- Sobieska, M.,Gajewska, E.,Kalmus, G., Samborski, W. Obesit, physical fitness and inflammatory markers in Polish children. (2013). *Medical Science. Monitor*. 19(1), 493-500.
- Sobti, R.C., Kler, R., Sharma, Y.P., Talwar, K.K., & Singh, N. (2012). Risk of obesity and type 2 diabetes with tumor necrosis factor-alpha 308G/A gene polymorphism in metabolic syndrome and coronary artery disease subjects. *Molecular and Cellular Biochemistry*, 360(1-2), 1-7.

- Tam, C.S., Clement, K., Baur, L.A., & Tordjman, J. (2010). Obesity and low-grade inflammation: a paediatric perspective. *Obesity Review*, 11(2), 118-126.
- Tsaoussoglou, M., Bixler, E.O., Calhoun, S., Chrousos, G.P., Sauder, K., & Vgontzas, A. N. (2010). Sleep-disordered breathing in obese children is associated with prevalent excessive daytime sleepiness, inflammation, and metabolic abnormalities. *The Journal of Clinical Endocrinology & Metabolism*, 95(1), 143-150.
- Tudor-Locke, C., Ainsworth, B.E., & Popkin, B.M. (2001). Active commuting to school: an overlooked source of childrens' physical activity? *Sports Medicine*, 31(5), 309-313.
- Tudor-Locke, C., Craig, C.L., Aoyagi, Y., Bell, R. C., Croteau, K. A., De Bourdeaudhuij, I. (2011). How many steps/day are enough? For older adults and special populations. *International Journal of Behavioral Nutrition and Physical Activity*, 8, 80. doi: 10.1186/1479-5868-8-80
- Tudor-Locke, C., Craig, C.L., Beets, M.W., Belton, S., Cardon, G.M., Duncan, S. (2011). How many steps/day are enough? for children and adolescents. *International Journal of Behavioral Nutrition and Physical Activity*, 8, 78.
- Vieira, A.C.R., Alvarez, M.M., Marins, V.M.R., Sichieri, R., & Veiga, G.V. (2006). Desempenho de pontos de corte do índice de massa corporal de diferentes referências na predição de gordura corporal em adolescentes. *Caderno de Saúde Pública*, 22(8), 1681-1690.
- Vitolo, M.R., Campagnolo, P.D.B., Barros, M.E., Gama, C.M., & Lopez, F.A. (2007). Avaliação de duas classificações para excesso de peso em adolescentes brasileiros. *Revista de Saúde Pública*, 41(4), 653-656.
- Vos, R.C., Wit, J.M., Pijl, H., & Houdijk, E.C. (2011). Long-term effect of lifestyle intervention on adiposity, metabolic parameters, inflammation and physical fitness in obese children: a randomized controlled trial. *Nutrition & Diabetes*, 1, e9.
- Wasilewska, A., Tenderenda, E., Taranta-Janusz, K., Tobolczyk, J., & Stypulkowska, J. (2012). Markers of systemic inflammation in children with hyperuricemia. *Acta Paediatr*, 101(5), 497-500.
- World Health Organization. (1995). Physical status: the use and interpretation of anthropometry. Geneva: World Health Organization; (Technical Report Series, 854).
- World Health Organization. (2000). Obesity: preventing and managing the global epidemic. Report of a WHO consultation. World Health Organ Tech Rep Ser 894: i-xii, 1-253.
- World Health Organization (2010). Obesity and overweight. What are overweight and obesity? World Health Organization. [Acesso em 8 mai. 2010].
- World Health Organization (2014). 10 facts on obesity: World Health Organization. Resource Document. London.
- Zimmermann, M.B., Gubeli, C., Puntener, C., & Molinari, L. (2004). Detection of overweight and obesity in a national sample of 6-12-y-old Swiss children: accuracy and validity of reference values for body mass index from the US Centers for Disease Control and Prevention and the International Obesity Task Force. *The American Journal of Clinical Nutrition*, 79(5), 838-843.

Author's note

Juliano Magalhães Guedes, José Bontempo Mamêde Neto, Alynne Christian Ribeiro Andaki and Edmar Lacerda Mendes are affiliated with the Federal University of Triângulo Mineiro, Department of Sport Science, Uberaba, MG, Brazil.

Alexandre de Paula Rogério is affiliated with the Federal University of Triângulo Mineiro, Department of Biomedicine, Uberaba, MG, Brazil.

Patrícia Feliciano Pereira is affiliated with the Federal University of Viçosa, Department of Nutrition Science, Viçosa, MG, Brazil.

Michelle Dias de Oliveira, Roberto Sousa Dias and Sérgio Oliveira de Paula are affiliated with the Federal University of Viçosa, Department of General Biology, Viçosa, MG, Brazil.

Antônio José Natali is affiliated with the Federal University of Viçosa, Department of Physical Education, Viçosa, MG, Brazil.

Acknowledgments

I would like to express the deepest appreciation to my teacher Dr. Edmar Lacerda Mendes for sharing attitude and substance of a genius, guidance, friendship, dedication and competence.

Corresponding author

Juliano Magalhães Guedes
Federal University of Triângulo Mineiro, Uberaba, Brazil. Department of Sports Sciences. 70, Nicolau Chereim, downtown, 37.200-000, Lavras, MG, Brazil.
Email: juliano_mguedes@yahoo.com.br

Manuscript accepted on February 10, 2015

Manuscript received on January 11, 2016



Motriz. The Journal of Physical Education. UNESP. Rio Claro, SP, Brazil
- eISSN: 1980-6574 – under a license Creative Commons - Version 3.0