

CompCytogen 10(4): 753–767 (2016)  
doi: 10.3897/CompCytogen.v10i4.9972  
<http://compcytogen.pensoft.net>

RESEARCH ARTICLE



COMPARATIVE  
Cytogenetics  
A peer-reviewed open-access journal  
International Journal of Plant & Animal Cytogenetics,  
Karyosystematics, and Molecular Systematics

# Comparative cytogenetics of tree frogs of the *Dendropsophus marmoratus* (Laurenti, 1768) group: conserved karyotypes and interstitial telomeric sequences

Lívia S. R. Teixeira<sup>1</sup>, Karin Regina Seger<sup>1</sup>, Cíntia Pelegrineti Targueta<sup>1</sup>,  
Victor G. Dill Orrico<sup>2</sup>, Luciana Bolsoni Lourenço<sup>1</sup>

<sup>1</sup> Departamento de Biologia Estrutural e Funcional, Instituto de Biologia, Universidade Estadual de Campinas, 13083-863 Campinas, São Paulo, Brasil <sup>2</sup> Departamento de Ciências Biológicas, Universidade Estadual de Santa Cruz, 45662-900 Ilhéus, Bahia, Brasil

Corresponding author: Luciana Bolsoni Lourenço ([bolsoni@unicamp.br](mailto:bolsoni@unicamp.br))

Academic editor: R. Noleto | Received 22 July 2016 | Accepted 4 November 2016 | Published 14 December 2016

<http://zoobank.org/298BF317-CA8B-4CAF-9A01-2604312656AA>

**Citation:** Teixeira LSR, Seger KG, Targueta CP, Orrico VGD, Lourenço LB (2016) Comparative cytogenetics of tree frogs of the *Dendropsophus marmoratus* (Laurenti, 1768) group: conserved karyotypes and interstitial telomeric sequences. Comparative Cytogenetics 10(4): 753–767. <https://doi.org/10.3897/CompCytogen.v10i4.9972>

## Abstract

The diploid number  $2n = 30$  is a presumed synapomorphy of *Dendropsophus* Fitzinger, 1843, although a noticeable variation in the number of biarmed/telocentric chromosomes is observed in this genus. Such a variation suggests that several chromosomal rearrangements took place after the evolutionary origin of the hypothetical ancestral 30-chromosome karyotype; however, the inferred rearrangements remain unknown. Distinct numbers of telocentric chromosomes are found in the two most cytogenetically studied species groups of *Dendropsophus*. In contrast, all three species of the *Dendropsophus marmoratus* (Laurenti, 1768) group that are already karyotyped presented five pairs of telocentric chromosomes. In this study, we analyzed cytogenetically three additional species of this group to investigate if the number of telocentric chromosomes in this group is not as variable as in other *Dendropsophus* groups. We described the karyotypes of *Dendropsophus seniculus* (Cope, 1868), *D. soaresi* (Caramaschi & Jim, 1983) and *D. novaisi* (Bokermann, 1968) based on Giemsa staining, C-banding, silver impregnation and *in situ* hybridization with telomeric probes. *Dendropsophus seniculus*, *D. soaresi* and *D. novaisi* presented five pairs of telocentric chromosomes, as did the remaining species of the group previously karyotyped. Though the species of this group show a high degree of karyotypic similarity, *D. soaresi* was unique in presenting large blocks

of het-ITSs (heterochromatic internal telomeric sequences) in the majority of the centromeres. Although the ITSs have been interpreted as evidence of ancestral chromosomal fusions and inversions, the het-ITSs detected in the karyotype of *D. soaresi* could not be explained as direct remnants of ancestral chromosomal rearrangements because no evidence of chromosomal changes emerged from the comparison of the karyotypes of all of the species of the *D. marmoratus* group.

### Keywords

Chromosomes, Anura, telomeric sequence

## Introduction

Faivovich et al. (2005) resurrected the genus *Dendropsophus* Fitzinger, 1843 to accommodate all Neotropical hylid species known or suspected to have a diploid chromosome number  $2n = 30$ . This cytogenetic character state was later confirmed as a synapomorphy for this genus by Suárez et al. (2013) after the description of a  $2n = 24$  karyotype for *Xenohyla* Izecksohn, 1998, the sister genus of *Dendropsophus* (see Faivovich et al. 2005, Pyron and Wiens 2011, Duellman et al. 2016). Based on preliminary data, Bogart (1973) hypothesized that centric fission events may have been involved in the origin of an ancestral 30-chromosome karyotype, which is a hypothesis that was also considered by Suárez et al. (2013). However, the chromosomes that are putatively involved in these events have not yet been recognized, and this hypothesis remains to be validated.

Although all of the *Dendropsophus* species karyotyped to date show  $2n = 30$  (see review in Catroli and Kasahara 2009, Medeiros et al. 2013, Suárez et al. 2013, Oliveira et al. 2016), a noticeable variation in the number of biarmed/telocentric chromosomes is observed among them, suggesting that several chromosomal rearrangements took place after the evolutionary origin of the hypothetical ancestral 30-chromosome karyotype. Karyotypes with only biarmed chromosomes [as in *D. minutus* (Peters, 1872) (Gruber et al. 2005) and *D. leali* (Bokermann, 1964) (Bogart 1973)] and karyotypes with up to five pairs of telocentric/subtelocentric chromosomes [as in *D. labialis* (Peters, 1863) (Bogart 1973), *D. sanborni* (Schmidt, 1944) and *D. jimi* (Napoli & Caramaschi, 1999) (Medeiros et al. 2013)] may be observed. However, the chromosomes and events involved in these rearrangements also remain undiscovered because most *Dendropsophus* species karyotypes are not yet described, and few chromosomal markers are available for the known karyotypes, preventing reliable hypotheses of chromosome homeology.

Of the nine species groups recognized in *Dendropsophus* (for a review of the *Dendropsophus* groups, see Faivovich et al. 2005), the *D. microcephalus* (Cope, 1886) group is the most species-rich (currently with 40 species—Frost 2016) and the most studied cytogenetically (17 species karyotyped—review by Catroli and Kasahara 2009, Medeiros et al. 2013, Oliveira et al. 2016). It is noteworthy that karyotypes without any telocentric chromosome (in *D. leali*—Bogart 1973) and with one [as in *D. bipunctatus* (Spix, 1824)—Bogart 1973], two [as in *D. phlebodes* (Steineger, 1906)—Kaiser et al. 1996], three [as in *D. cruzi* (Pombal & Bastos, 1998)—Gruber et al. 2005], four [as in

*D. nanus* (Boulenger, 1889)—Medeiros et al. 2003] or five (as in *D. jimi*—Medeiros et al. 2013) telocentric chromosome pairs are observed in this group. Karyotypes with distinct numbers of telocentric chromosomes were also found in the *D. leucophyllatus* (Beireis, 1783) group (Bogart 1973, Kaiser et al. 1996, Gruber et al. 2005), which currently has 11 species (see Frost 2016) and is the second most cytogenetically studied species group of *Dendropsophus* (four of the named species are karyotyped). In contrast, all of the three species of the *D. marmoratus* group that are already karyotyped (i.e. *D. marmoratus*, *D. melanargyreus* and *D. nahdereri*) present five pairs of telocentric chromosomes (Bogart 1973, Gruber et al. 2005, Suárez et al. 2013).

Chromosomal sites composed of telomeric repeats localized apart from the telomeres, also known as interstitial or intrachromosomal telomeric sequences (ITSs) or repeats (ITRs), have been detected in several animals (Meyne et al. 1990, Nanda et al. 2002, Rovatsos et al. 2015, Schmid and Steinlein 2016) and plants (Tek and Jiang 2004, He et al. 2013). Based on the genomic location and sequence organization, especially in the number of telomeric repeats, Ruiz-Herrera et al. (2008) classified the ITSs in short ITSs (s-ITSs) and heterochromatic ITS (het-ITS). The s-ITSs [called short interstitial telomeres, short ITs, by Azzalin et al. (2001)] are short stretches of telomeric hexamers distributed at internal chromosomal positions, presumably present in all vertebrate species, whereas het-ITSs are large blocks of telomeric-like repeats localized mainly in centromeric and pericentromeric regions (Ruiz-Herrera et al. 2008). The s-ITSs probably originated from the insertion of telomeric repeats during the repair of DNA double-strand breaks, as was originally proposed by Nergadze et al. (2004, 2007). On the other hand, the het-ITSs have been widely considered to be remnants of ancestral chromosomal rearrangements as fusions (e.g., Lee et al. 1993, Slijepcevic 1998, Ropiquet et al. 2010, Paço et al. 2013, Young et al. 2013) and inversions (e.g., Farré et al. 2009, Ocalewicz et al. 2013, Paço et al. 2013). Recently, Schmid and Steinlein (2016) proposed an additional category of ITS, named euchromatic-ITSs (eu-ITSs), to accommodate the large ITSs that are not revealed as heterochromatic sites by C-banding or staining with base-specific fluorochromes.

The *Dendropsophus marmoratus* group currently includes eight species, i.e., *D. marmoratus*, *D. acreanus* (Bokermann, 1964), *D. dutrai* (Gomes & Peixoto, 1996), *D. melanargyreus*, *D. nahdereri*, *D. novaisi*, *D. seniculus* and *D. soaresi* (Faivovich et al. 2005). Some adult and larval morphological synapomorphies of this species group may be recognized (Faivovich et al. 2005); however, its cladistic proximity with other *Dendropsophus* species groups as well as the internal phylogenetic relationships of this group remain unclear (Faivovich et al. 2005; Pyron and Wiens 2011, Fouquet et al. 2011, Medeiros et al. 2013). To date, up to three of the eight species of the *D. marmoratus* group have been included in phylogenetic analysis (Fouquet et al. 2011).

In this study, we analyzed cytogenetically three additional species of the *Dendropsophus marmoratus* group to investigate if the number of telocentric chromosomes in this group is not as variable as in other *Dendropsophus* groups. Because karyotypic variation in number of telocentric chromosomes may result from rearrangements involving telomeric sequences (review in Ruiz-Herrera et al. 2008), we included here

the mapping of telomeric sequences in the karyotypes of two of the analyzed species. Additionally, we provided the nucleotide sequence of a fragment of the 16S ribosomal RNA gene of one exemplar for each of the species that were analyzed cytogenetically to yield a reliable association of the chromosomal data set with a DNA data set that has been remarkably useful for taxonomic and phylogenetic studies of anurans.

## Material and methods

### Specimens

Four male exemplars of *Dendropsophus seniculus* from Ribeirão Grande, state of São Paulo, Brazil, nine *D. soaresi* males from Barreiras, state of Bahia, Brazil and one female of *D. novaisi* from Jequié, state of Bahia, Brazil were analyzed cytogenetically. The specimens were collected under a permit issued by the Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis (IBAMA) (#32483), and deposited at the amphibian collection of the Museu de Zoologia “Prof. Adão José Cardoso” at the Institute of Biology – University of Campinas, Campinas, Brazil, under the accession numbers ZUEC 17225–17228 (*D. seniculus*), ZUEC 16867–16875 (*D. soaresi*) and ZUEC 17858 (*D. novaisi*).

### Cytogenetic analyses

Animals were injected intraperitoneally with 2% colchicine (Sigma – Aldrich; 0.02 mL per 1 g body weight) for an “in vivo” treatment that lasted at least 4 hours. The animals were deeply anesthetized with lidocaine gel 2% and their intestines were removed and used for obtaining chromosomal preparations according to the method of King and Rofe (1976). Chromosomes were conventionally stained with 10% Giemsa and sequentially submitted to C-banding (Sumner 1972) and silver staining by the Ag-NOR method (Howell and Black 1980).

To localize telomeric sequences, the karyotypes were *in situ* hybridized with the probe (CCCTAA)<sub>3</sub> (PNA – Peptid Nucleic Acid TelC-Cy3; PNA Bio Inc.), following the manufacturer’s instructions.

### Mitochondrial DNA sequences

Samples of genomic DNA were obtained from *Dendropsophus seniculus* (ZUEC 17225), *D. soaresi* (ZUEC 16867) and *D. novaisi* (ZUEC 17858) following the procedure reported by Medeiros et al. (2013). A fragment of approximately 1300 bp of the 16S ribosomal RNA gene was PCR-amplified using the primers 12L13(L) (Feller


and Hedges 1998) and 16Sbr(H) (Palumbi et al. 1991). The amplified products were purified with the GFX PCR and Gel Band DNA purification Kit (GE Healthcare) and directly sequenced in an automatic DNA ABI/Prism sequencer (Applied Biosystems) using BigDye Terminator kits (Applied Biosystems) and the primers 12L13 (Feller and Hedges 1998), TitusI(H) (Titus 1992), Hedges16L2a (Hedges 1994), Hedges16H10 (Hedges 1994), 16Sar(L) (Palumbi et al. 1991) and 16Sbr(H) (Palumbi et al. 1991). DNA sequences were aligned using ClustalW option implemented in BioEdit v. 7.2.5 (Hall 1999) and compared to each other and to the 16S rDNA sequence of *D. seniculus* available at GenBank (AY843666).

## Results


### Cytogenetic analyses

The karyotypes of *Dendropsophus seniculus*, *D. soaresi* and *D. novaisi* were very similar and presented three pairs (pairs 1, 2 and 4) of submetacentric chromosomes, seven pairs (pairs 3, 8–12 and 14) of metacentric chromosomes and five pairs (pairs 5–7, 13 and 15) of telocentric chromosomes (Figures 1–3). The nucleolus organizer region (NOR) was detected by silver staining in the long arm of chromosome 9 of the three species (insets in Figures 1–3). C-banding only detected the centromeric region of all the chromosomes of *D. seniculus* (Figure 1b), *D. soaresi* (Figure 2b) and *D. novaisi* (Figure 3b).


*In situ* hybridization detected telomeric sequences in all of the telomeres of *Dendropsophus seniculus* and *D. soaresi* (Figure 4). Additionally, interstitial telomeric sequences (ITSs) were detected in the centromeres of the chromosomes of *D. soaresi*, except in two of the five pairs of telocentric chromosomes (pairs 5 and 6) (Figure 4b).



**Figure 1.** Karyotype of *Dendropsophus seniculus* stained with Giemsa (a) and C-banded (b). In the inset in (a), the NOR-bearing chromosome pair 9 after silver staining. Bar = 5  $\mu$ m.



**Figure 2.** Karyotype of *Dendropsophus soaresi* stained with Giemsa (a) and C-banded (b). In the inset in (a), the NOR-bearing chromosome pair 9 after silver staining. Bar = 5  $\mu$ m.




**Figure 3.** Karyotype of *Dendropsophus novaisi* stained with Giemsa (a) and C-banded (b). In the inset in (a), the NOR-bearing chromosome pair 9 after silver staining. Bar = 5  $\mu$ m.

### Mitochondrial DNA sequences

The nucleotide sequence (1312 bp) (see Suppl. material 1) of the 16S rDNA of the specimen of *Dendropsophus seniculus* that we analyzed was highly similar (99.6%) to the corresponding sequence available at GenBank (AY843666; Faivovich et al. 2005) from a specimen of *D. seniculus* from Angra dos Reis, Rio de Janeiro State, Brazil. The sequences obtained from *D. soaresi* (1314 bp) and *D. novaisi* (1310 bp) (see Suppl. material 1) were 86.4% similar to each other, and 88.4% and 90.7% (average value) similar to the sequences of *D. seniculus*, respectively.





**Figure 4.** Karyotypes of *Dendropsophus seniculus* (a) and *Dendropsophus soaresi* (b) hybridized with telomeric probe.

## Discussion

### Karyotypic comparisons

The three species analyzed showed karyotypes composed of five pairs of telocentric chromosomes, similarly to the other three species of the *D. marmoratus* group previously studied cytogenetically [i.e. *D. marmoratus* (Bogart 1973, Suárez et al. 2013), *D. melanargyreus* (Suárez et al. 2013) and *D. nahdereri* (Gruber et al. 2005)]. The inter-specific morphological conservation of the karyotypes of the species of the *D. marmoratus* group contrasts with the variation found in other *Dendropsophus* groups. The *D. microcephalus* group, for instance, includes species with zero to five pairs of telocentric chromosomes (see Bogart 1973, Kaiser et al. 1996, Gruber et al. 2005, Medeiros et al. 2003). Variation in the number of telocentric chromosomes could also be found in the *D. leucophyllatus* group, although only four named species of this group have been karyotyped (see Bogart 1973, Kaiser et al. 1996, Gruber et al. 2005).

According to the estimated dates of divergence provided by Duellman et al. (2016), the *D. marmoratus*, *D. microcephalus* and *D. leucophyllatus* groups arose at similar times in the mid-Miocene (17.0, 17.2 and 18.7 Mya, respectively). Therefore, differential time for divergence does not justify the different levels of karyotypic variation observed among the three aforementioned species groups. Further analyses of the chromosomal rearrangements involved in the karyotypic variations in *Dendropsophus* combined with phylogeographic studies are still necessary to elucidate about the high conservation in the number of telocentric chromosomes in the *D. marmoratus* group.

With respect to the number and relative size of the telocentric chromosomes, the karyotypes of the species of the *D. marmoratus* group are similar to that of *D. labialis* (Bogart 1973), a species included in the *D. labialis* group. This morphological similarity suggests that the telocentric chromosomes of these karyotypes could be homeologous,

although a better characterization of these chromosomes is fundamental to test this hypothesis. According to the most comprehensive phylogenetic analysis of *Dendropsophus* (Duellman et al. 2016) and assuming the telocentric chromosomes of the species of the *D. marmoratus* group and *D. labialis* are homeologous, it is possible to hypothesize that this karyotype configuration is plesiomorphic with respect to those constituted by other numbers and/or relative sizes of the telocentric chromosomes. However, internal relationships within *Dendropsophus* are consistently poorly supported and small taxonomic additions cause huge impacts (e.g., Fouquet et al. 2015, Duellman et al. 2016).

Five pairs of telocentric chromosomes were also observed in the karyotypes of *D. jimi* and *D. sanborni* (Medeiros et al. 2013), which are species that belong to the *D. microcephalus* group. In these karyotypes, however, the telocentric pairs were classified as pairs 5, 6, 12, 13 and 15, whereas the telocentric chromosomes of the karyotypes of *D. labialis* and the species of the *D. marmoratus* group are numbered as pairs 5, 6, 7, 13 and 15. Because only a few chromosomal markers are available for a comparison of these karyotypes, it is still not possible to determine if the telocentric chromosomes of all of these species are homeologous. Therefore, we cannot discard the possibility that chromosomes 12 of *D. jimi* and *D. sanborni* are homeologous to chromosomes 7 of *D. labialis* and the species of the *D. marmoratus* group, although these chromosomes differ by the presence of NOR in chromosomes 12 of *D. jimi* and *D. sanborni* (Medeiros et al. 2013).

The similarities among the karyotypes of the species of the *Dendropsophus marmoratus* group are not restricted to the number of telocentric chromosomes. *Dendropsophus seniculus*, *D. soaresi* and *D. novaisi* also share with *D. marmoratus* and *D. melanargyreus* the location of the NOR at a distal site of the long arm of chromosome 9, which differs from *D. nahdereri*, whose NOR is located on the short arm of the submetacentric chromosome 1 (Gruber et al. 2005).

C-banding did not reveal any differential band that could be considered exclusive to the karyotypes of *D. seniculus*, *D. soaresi* or *D. novaisi*, since only the centromeric regions were detected by this technique (present work). Conspicuous non-centromeric C-bands were also absent in the karyotypes of *D. marmoratus* and *D. melanargyreus*, the other two species of the *D. marmoratus* group whose karyotypes were already C-banded, although Suárez et al. (2013) reported the presence of some distal and interstitial C-bands in those karyotypes.

Despite the high similarity of the karyotypes of the species of the *Dendropsophus marmoratus* group with respect to the number and morphology of the chromosomes, C-banding pattern and location of NOR (except for *D. nahdereri*), the karyotype of *D. soaresi* stands out because of the presence of internal telomeric sequences in addition to the terminal telomeric sequences.

### Interstitial telomeric sequences

Large and short ITSs are likely to play a role in karyotypic evolution. Several studies support the hypothesis that, in addition to possibly representing relics of chromosomal



changes, the het-ITSs may themselves induce chromosome breakage and subsequent chromosomal rearrangements (reviewed in Ruiz-Herrera et al. 2008 and Bolzán 2012). Similarly, experimental and associative studies have also suggested the involvement of s-ITSs with genomic instability or chromosomal hot spots of recombination (Akse-nova et al. 2013, Wood et al. 2015).

The het-ITSs detected in the present study in the karyotype of *Dendropsophus soaresi* cannot be explained as direct remnants of ancestral chromosomal rearrangements because no evidence of chromosomal changes has emerged from the comparison of the karyotypes of all species of the *D. marmoratus* group already known (Bogart 1973, Gruber et al. 2005, Suárez et al. 2013, present work). Although it is very similar to the karyotypes of the other species of the group, the karyotype of *D. soaresi* is unique in presenting large blocks of centromeric ITSs because the karyotypes of *D. seniculus* (present study), *D. melanargyreus* and *D. marmoratus* (Suárez et al. 2013) showed only telomeric sites hybridized with telomeric probes.

The occurrence of het-ITSs at the majority of the centromeres of the karyotype of *Dendropsophus soaresi* is also remarkable and suggests the expansion and homogenization of telomeric sequences throughout the repetitive elements that compose these centromeric regions. Repetitive DNA, such as centromeric satellite DNA, is expected to expand in the genome and evolve in concert by a series of mechanisms, including unequal crossing-over, gene conversion, rolling circle replication and reinsertion, and transposon-mediated exchange (see Dover 1982, 1986 and the review by Plohl et al. 2008). The telomeric repeats present in heterochromatic sites should be subject to the same evolutionary forces (see Ruiz-Herrera et al. 2008). In contrast, the absence of het-ITS in the centromere of two chromosome pairs (telocentric chromosome pairs 5 and 6) of *Dendropsophus soaresi* suggests that these centromeres do not evolve in concert with the remaining centromeric regions of the genome. The reason for such differential behavior is intriguing and remains unknown.

Similar to observations of *Dendropsophus soaresi*, large blocks of centromeric/pericentromeric ITSs that were widely distributed throughout the genome were previously found in four other hylid species [i.e. *Aplastodiscus albofrenatus* (Lutz, 1924), *A. arildae* (Cruz & Peixoto, 1987) and *A. eugenioi* (Carvalho-e-Silva & Carvalho-e-Silva, 2005)—Carvalho et al. 2009, Gruber et al. 2012a; *Hypsiboas faber* (Wied-Neuwied, 1821)—Schmid and Steinlein 2016]. In the hylid *Itapotihyla langsdorffii* (Duméril & Bibron, 1841), ITSs were also observed in several centromeres (Gruber et al. 2012b), but in this case the het-ITSs are not as large as those previously mentioned. In addition to the aforementioned hylids, other fifteen hylid species showed ITSs in their karyotypes (Meyne et al. 1990, Wiley et al. 1992, Suárez et al. 2013, Mattos et al. 2014, Bruschi et al. 2014, Schmid and Steinlein 2016), which suggests that the appearance of this type of sequence is recurrent in the Hylidae family. Only the centromeric ITS found in chromosome 3 of *Scarthyia goinorum* (Bokermann, 1962) was clearly interpreted as a remnant of a chromosomal fusion that in that case could respond to the reduced chromosome number observed in this species (Suárez et al. 2013). The insertion of telomeric repeats during the repair of double-strand

breaks in DNA as a phenomenon putatively involved in the origin of ITS in Hylidae remains unexplored.

It is worth noticing that in the sample of metaphases analyzed in this paper, large signals of the telomeric probe were detected at a subterminal non-heterochromatic site of some chromosomes of *Dendropsophus seniculus* (Figure 4a, at the long arm of the right homologous of chromosome 9). This hybridization pattern resembles that pattern interpreted by Wood et al. (2014, 2015) as cytological evidence of the occurrence of t-loops formed between telomere and s-ITS. However, studies designed to search for s-ITSs in hylid karyotypes have not yet been performed, and the prevalence of ITSs in Hylidae remains an intriguing question to be assessed in further studies.

### Association between cytogenetic data and 16S rDNA sequences

The high similarity between the 16S rDNA sequence of *Dendropsophus seniculus* we provided and that previously obtained by Faivovich et al. (2005) enables a reliable association between the cytogenetic data shown here and the analyses hitherto conducted with the previously available sequence, including the studies of Fouquet et al. (2015) and Duellman et al. (2016). On the other hand, the nucleotide sequences obtained here from *Dendropsophus soaresi* and *D. novaisi* were the first report of 16S rDNA sequences for these species.

*Dendropsophus* systematics are in flux and even comprehensive datasets are unable to provide a stable historical hypothesis (Fouquet et al. 2011; Peloso et al. 2016). The association between the cytogenetic dataset and 16S rDNA sequences may be very helpful in future analyses, especially because the species-level taxonomy of *Dendropsophus* has been subject to several changes. A number of *Dendropsophus* species has been described in the last few years (Rivera-Correia and Orrico 2013, Ortega-Andrade and Ron 2013, Orrico et al. 2014, Fouquet et al. 2015, Peloso et al. 2016) as well as species synonymization has been proposed (Guarnizo et al. 2012, Orrico et al. 2013). Therefore, a reliable association between different sets of data is fundamental for further integrative studies.

### Conclusion

All of the karyotypes found in the *Dendropsophus marmoratus* group to date showed five pairs of telocentric chromosomes and were also similar in the location of NORs (except for the *D. nahdereri* karyotype, described by Gruber et al. 2005) and C-banding pattern. Because of this karyotypic conservatism, the het-ITSs present in the majority of the centromeres of the karyotype of *D. soaresi* may not be interpreted as direct remnants of ancestral chromosomal rearrangements.

## Acknowledgments

The authors are thankful to Daniel Pacheco Bruschi for help in producing the cytogenetic data from *Dendropsophus novaisi*. This study was supported by the Brazilian agencies São Paulo Research Foundation (FAPESP) and Fundo de Apoio ao Ensino, à Pesquisa e Extensão (FAEPEX) da Universidade Estadual de Campinas.

## References

- Aksenova AY, Greenwell PW, Dominska M, Shishkin AA, Kim JC, Petes TD, Mirkina SM (2013) Genome rearrangements caused by interstitial telomeric sequences in yeast. *PNAS* 110: 19866–19871. <https://doi.org/10.1073/pnas.1319313110>
- Azzalin CM, Nergadze SG, Giulotto E (2001) Human intrachromosomal telomeric-like repeats: sequence organization and mechanisms of origin. *Chromosoma* 110: 75–82. <https://doi.org/10.1007/s004120100135>
- Bogart JP (1973) Evolution of anuran karyotypes. In: Vial JL (Ed.) *Evolutionary Biology of the Anurans: Contemporary Research on Major Problems*. University of Missouri Press, Columbia, 337–349.
- Bolzán AD (2012) Chromosomal aberrations involving telomeres and interstitial telomeric sequences. *Mutagenesis* 27: 1–15. <https://doi.org/10.1093/mutage/ger052>
- Bruschi DP, Rivera M, Lima AP, Zúñiga AB, Recco-Pimentel SM (2014) Interstitial Telomeric Sequences (ITS) and major rDNA mapping reveal insights into the karyotypical evolution of Neotropical leaf frogs species (*Phyllomedusa*, Hylidae, Anura). *Molecular Cytogenetics* 7: 22. <https://doi.org/10.1186/1755-8166-7-22>
- Catrolí GF, Kasahara S (2009) Cytogenetic data on species of the family Hylidae (Amphibia, Anura): results and perspectives. *Publ. UEPG. Ciências Biológicas e da Saúde* 15: 67–86.
- Carvalho KA, Garcia PCA, Recco-Pimentel SM (2009) NOR Dispersion, Telomeric Sequence Detection in Centromeric Regions and Meiotic Multivalent Configurations in Species of the *Aplastodiscus albofrenatus* Group (Anura, Hylidae). *Cytogenetic Genome Research* 126: 359–367. <https://doi.org/10.1159/000264179>
- Dover GA (1982) Molecular drive: a cohesive mode of species evolution. *Nature* 299: 111–117. <https://doi.org/10.1038/299111a0>
- Dover GA (1986) Molecular drive in multigene families: how biological novelties arise, spread and are assimilated. *Trends Genet* 122: 159–165. [https://doi.org/10.1016/0168-9525\(86\)90211-8](https://doi.org/10.1016/0168-9525(86)90211-8)
- Duellman WE, Marion AB, Hedges SB (2016) Phylogenetics, classification, and biogeography of the treefrogs (Amphibia: Anura: Arboranae). *Zootaxa* 4101: 1–109. <https://doi.org/10.11646/zootaxa.4104.1.1>
- Faivovich J, Haddad CFB, Garcia PCA, Frost DR, Campbell JA, Wheeler WC (2005) Systematic review of the frog family Hylidae, with special reference to Hylinae: phylogenetic analysis and taxonomic revision. *Bulletin of the American Museum of Natural History* 294: 1–240. [https://doi.org/10.1206/0003-0090\(2005\)294\[0001:SROTFF\]2.0.CO;2](https://doi.org/10.1206/0003-0090(2005)294[0001:SROTFF]2.0.CO;2)

- Farré M, Ponsà M, Bosch M (2009) Interstitial telomeric sequences (ITSs) are not located at the exact evolutionary breakpoints in primates. *Cytogenetic and Genome Research* 124: 128–131. <https://doi.org/10.1159/000207517>
- Feller A, Hedges SB (1998) Molecular evidence for the early history of living amphibians. *Molecular Phylogenetics and Evolution* 9: 509–516. <https://doi.org/10.1006/mpev.1998.0500>
- Fouquet A, Noonan B, Blanc M, Orrico VGD (2011) Phylogenetic position of *Dendropsophus gaucheri* (Lescure and Marty 2000) highlights the need for an in-depth investigation of the phylogenetic relationships of *Dendropsophus* (Anura: Hylidae). *Zootaxa* 3035: 59–67. <https://doi.org/10.11646/zootaxa.4052.1.2>
- Fouquet A, Orrico VGD, Ernst R, Blanc M, Martinez Q, Vacher JP, Rodrigues MT, Ouboter PE, Jairam R, Ron S (2015) A new *Dendropsophus* Fitzinger, 1843 (Anura: Hylidae) of the *parviceps* group from the lowlands of the Guiana Shield. *Zootaxa* 4052: 39–64.
- Frost DR (2016) Amphibian species of the world: an online reference. Version 6.0 (1<sup>st</sup> July, 2016). Electronic database accessible at <http://research.amnh.org/herpetology/amphibia/index.html>. American Museum of Natural History, New York, USA.
- Guarnizo CE, Escallón C, Cannatella DC, Amésquita A (2012) Congruence between acoustic traits and genealogical history reveals a new species of *Dendropsophus* (Anura: Hylidae) in the high Andes of Colombia. *Herpetologica* 68: 523–540. <https://doi.org/10.1655/HERPETOLOGICA-D-10-00038>
- Gruber SL, Haddad CFB, Kasahara S (2005) Evaluating the karyotypic diversity in species of *Hyla* (Anura; Hylidae) with 2n=30 chromosomes based on the analysis of ten species. *Folia Biologica (Praha)* 51: 68–75.
- Gruber SL, Zina J, Narimatsu H, Haddad CFB, Kasahara S (2012a) Comparative karyotype analysis and chromosome evolution in the genus *Aplastodiscus* (Cophomantini, Hylinae, Hylidae). *BMC Genetics* 13: 28. <https://doi.org/10.1186/1471-2156-13-28>
- Gruber SL, Haddad CFB, Kasahara S (2012b) Karyotype analysis of seven species of the tribe Lophiohylini (Hylinae, Hylidae, Anura), with conventional and molecular cytogenetic techniques. *Comparative Cytogenetics* 6: 409–423. <https://doi.org/10.3897/compcytogen.v6i4.3945>
- Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95–98.
- He L, Liu J, Torres GA, Zhang H, Jiang J, Xie C (2013) Interstitial telomeric repeats are enriched in the centromeres of chromosomes in *Solanum* species. *Chromosome Research* 21: 5–13. <https://doi.org/10.1007/s10577-012-9332-x>
- Hedges SB (1994) Molecular evidence for the origin of birds. *Proceedings of the National Academy of Sciences* 91: 2621–2624. <https://doi.org/10.1073/pnas.91.7.2621>
- Howell WM, Black DA (1980) Controlled silver staining of nucleolar organizer regions with a protective colloidal developer a 1-step method. *Experientia* 36: 1014–1015. <https://doi.org/10.1007/BF01953855>
- Kaiser H, Mais C, Bolános F, Steinlein C, Feichtinger W, Schmid M (1996) Chromosomal, investigation of three Costa Rican frogs from the 30-chromosome radiation of *Hyla* with the description of a unique geographic variation in nucleolus organizer regions. *Genetica* 98: 95–102. <https://doi.org/10.1007/BF00120223>

- King M, Rofe R (1976) Karyotypic variation in the Australian gekko *Phyllodactylus marmoratus* (Gray) (Gekkonidae: Reptilia). *Chromosoma* 54: 75–87. <https://doi.org/10.1007/BF00331835>
- Lee C, Sasi R, Lin CC (1993) Interstitial localization of telomeric DNA sequences in the Indian muntjac chromosomes: further evidence for tandem chromosome fusions in the karyotypic evolution of the Asian muntjacs. *Cytogenetics and Cell Genetics* 63: 156–159. <https://doi.org/10.1159/000133525>
- Mattos TL, Coelho AC, Schneider CH, Telles DOC, Menin M, Gross MC (2014) Karyotypic diversity in seven Amazonian anurans in the genus *Hypsiboas* (family Hylidae). *BMC Genetics* 15: 43. <https://doi.org/10.1186/1471-2156-15-43>
- Medeiros LR, Rossa-Feres DC, Recco-Pimentel SM (2003) Chromosomal differentiation of *Hyla nana* and *Hyla sanborni* (Anura, Hylidae) with a description of NOR polymorphism in *H. nana*. *Journal of Heredity* 94: 149–154. <https://doi.org/10.1093/jhered/esg019>
- Medeiros LC, Lourenço LB, Rossa-Feres DC, Lima AP, Andrade GV, Giaretta AA, Egito GTBT, Recco-Pimentel SM (2013) Comparative cytogenetic analysis of some species of the *Dendropsophus microcephalus* group (Anura, Hylidae) in the light of phylogenetic inferences. *BMC Genetics* 14: 59. <https://doi.org/10.1186/1471-2156-14-59>
- Meyne J, Baker RJ, Hobart HH, Hsu TC, Ryder OA, Ward OG, Wiley JE, Wurster-Hill DH, Yates TL, Moyzis RK (1990) Distribution of non-telomeric sites of the (TTAGGG)<sub>n</sub> telomeric sequence in vertebrate chromosomes. *Chromosoma* 99: 3–10. <https://doi.org/10.1007/BF01737283>
- Nanda I, Schrama D, Feichtinger W, Haaf T, Scharl M, Schmid M (2002) Distribution of telomeric (TTAGGG)<sub>n</sub> sequences in avian chromosomes. *Chromosoma* 111: 215–227. <https://doi.org/10.1007/s00412-002-0206-4>
- Nergadze SG, Rocchi M, Azzalin CM, Mondello C, Giulotto E (2004) Insertion of telomeric repeats at intrachromosomal break sites during primate evolution. *Genome Research* 14: 1704–1710. <https://doi.org/10.1101/gr.2778904>
- Nergadze SG, Santagostino MA, Salzano A, Mondello C, Giulotto E (2007) Contribution of telomerase RNA retrotranscription to DNA double-strand break repair during mammalian genome evolution. *Genome Biology* 8: R260. <https://doi.org/10.1186/gb-2007-8-12-r260>
- Ocalewicz K, Furgala-Selezniow G, Szmyt M, Lisboa R, Kucinski M, Lejk AM, Jankun M (2013) Pericentromeric location of the telomeric DNA sequences on the European grayling chromosomes. *Genetica* 141: 409–416. <https://doi.org/10.1007/s10709-013-9740-7>
- Oliveira IS, Noletto RB, Oliveira AKC, Toledo LF, Cestari MM (2016) Comparative cytogenetic analysis of four species of *Dendropsophus* (Hylinae) from the Brazilian Atlantic forest. *Journal of Genetics* 95: 349–355. <https://doi.org/10.1007/s12041-016-0645-y>
- Orrico VGD, Duellman WE, Souza MB, Haddad CFB (2013) The taxonomic status of *Dendropsophus allenorum* and *D. timbeba* (Anura: Hylidae). *Journal of Herpetology* 46: 615–618. <https://doi.org/10.1670/12-208>
- Orrico VGD, Peloso PLV, Sturaro MJ, Silva-Filho HF, Neckel-Oliveira S, Gordo M, Faivovich J, Haddad CFB (2014) A new “Bat-Voiced” species of *Dendropsophus* Fitzinger, 1843 (Anura, Hylidae) from the Amazon Basin, Brazil. *Zootaxa* 3881: 341–361. <https://doi.org/10.11646/zootaxa.3881.4.3>

- Ortega-Andrade HM, Ron SR (2013) A new species of small tree frog, genus *Dendropsophus* (Anura: Hylidae) from the eastern Amazon lowlands of Ecuador. *Zootaxa* 3652: 163–178. <https://doi.org/10.11646/zootaxa.3652.1.6>
- Paço A, Chaves R, Vieira-da-Silva A, Adegas F (2013) The involvement of repetitive sequences in the remodelling of karyotypes: The *Phodopus* genomes (Rodentia, Cricetidae). *Micron* 46: 27–34. <https://doi.org/10.1016/j.micron.2012.11.010>
- Palumbi SR, Martin A, Romano S, McMillan WO, Stice L, Grabowski G (1991) The simple fool's guide to PCR, Version 2.0. Privately published document compiled by Palumbi SR. Special Publication of Department of Zoology, University of Hawaii, Honolulu, USA, 44 pp.
- Peloso PLV, Orrico VGD, Haddad CFB, Lima-Filho GR, Sturaro MJ (2016) A new species of clown tree frog *Dendropsophus leucophyllatus* species group, from Amazonia (Anura, Hylidae). *South American Journal of Herpetology* 11: 66–80. <https://doi.org/10.2994/SAJH-D-16-00003.1>
- Plohl M, Luchetti A, Meštrović N, Mantovani B (2008) Satellite DNAs between selfishness and functionality: Structure, genomics and evolution of tandem repeats in centromeric (hetero) chromatin. *Gene* 409: 72–82. <https://doi.org/10.1016/j.gene.2007.11.013>
- Pyron RA, Wiens JJ (2011) A large-scale phylogeny of Amphibia including over 2800 species, and a revised classification of extant frogs, salamanders, and caecilians. *Molecular Phylogenetics and Evolution* 61: 543–583. <https://doi.org/10.1016/j.ympev.2011.06.012>
- Rivera-Correa M, Orrico VGD (2013) Description and phylogenetic relationships of a new species of treefrog of the *Dendropsophus leucophyllatus* group (Anura: Hylidae) from the Amazon basin of Colombia and with an exceptional color pattern. *Zootaxa* 3686: 447–460. <https://doi.org/10.11646/zootaxa.3686.4.3>
- Ropiquet A, Hassanin A, Pagacova E, Gerbault-Seureau M, Cernohorska H, Kubickova S, Bonillo C, Rubes J, Robinson TJ (2010) A paradox revealed: karyotype evolution in the four-horned antelope occurs by tandem fusion (Mammalia, Bovidae, *Tetracerus quadricornis*). *Chromosome Research* 18: 277–286. <https://doi.org/10.1007/s10577-010-9115-1>
- Rovatsos M, Kratochvíl L, Altmanová M, Pokorná MJ (2015) Interstitial telomeric motifs in squamate reptiles: when the exceptions outnumber the rule. *PLoS ONE* 10: e0134985. <https://doi.org/10.1371/journal.pone.0134985>
- Ruiz-Herrera A, Nergadze SG, Santagostino M, Giulotto E (2008) Telomeric repeats far from the ends: mechanisms of origin and role in evolution. *Cytogenetic and Genome Research* 122: 219–228. <https://doi.org/10.1159/000167807>
- Schmid M, Steinelin C (2016) Chromosome Banding in Amphibia. XXXIV. Intrachromosomal Telomeric DNA Sequences in Anura. *Cytogenetic and Genome Research* 148: 211–226. <https://doi.org/10.1159/000446298>
- Slijepcevic P (1998) Telomeres and mechanisms of Robertsonian fusion. *Chromosoma* 107: 136–140. <https://doi.org/10.1007/s004120050289>
- Suárez P, Cardozo D, Baldo D, Pereyra MO, Faivovich J, Orrico VGD, Catroli GF, Grabielle M, Bernarde PS, Nagamachi CY, Haddad CFB, Pieczarka JC (2013) Chromosome evolution in *Dendropsophini* (Amphibia, Anura, Hylinae). *Cytogenetic and Genome Research* 141: 295–308. <https://doi.org/10.1159/000354997>



- Sumner AT (1972) A simple technique for demonstrating centromeric heterochromatin. *Experimental Cell Research* 75: 304–306. [https://doi.org/10.1016/0014-4827\(72\)90558-7](https://doi.org/10.1016/0014-4827(72)90558-7)
- Tek AL, Jiang J (2004) The centromeric regions of potato chromosomes contain megabase-sized tandem arrays of telomere-similar sequence. *Chromosoma* 113: 77–83. <https://doi.org/10.1007/s00412-004-0297-1>
- Titus TA, Larson A (1996) Molecular phylogenetics of desmognathine salamanders (Caudata: Plethodontidae): a reevaluation of evolution in ecology, life history, and morphology. *Systematic Biology* 45: 229–238. <https://doi.org/10.1093/sysbio/45.4.451>
- Wiley JE, Meyne J, Little ML, Stout JC (1992) Interstitial hybridization sites of the (TTAGGG)<sub>n</sub> telomeric sequence on the chromosomes of some North American hylid frogs. *Cytogenetics and Cell Genetics* 61: 55–57. <https://doi.org/10.1159/000133368>
- Wood AM, Rendtlew Danielsen JM, Lucas CA, Rice EL, Scalzo D, Shimi T, Goldman RD, Smith ED, Le Beau MM, Kosak ST (2014) TRF2 and lamin A/C interact to facilitate the functional organization of chromosome ends. *Nature Communications* 5: 5467. <https://doi.org/10.1038/ncomms6467>
- Wood AM, Laster K, Rice EL, Kosak ST (2015) A beginning of the end: new insights into the functional organization of telomeres. *Nucleus* 6: 172–178. <https://doi.org/10.1080/19491034.2015.1048407>
- Young MJ, O’Meally D, Sarre SD, Georges A, Ezaz T (2013) Molecular cytogenetic map of the central bearded dragon, *Pogona vitticeps* (Squamata: Agamidae). *Chromosome Research* 21: 361–374. <https://doi.org/10.1007/s10577-013-9362-z>

## Supplementary material I

### 16S rDNA sequences

Authors: Lívia S. R. Teixeira, Karin Regina Seger, Cíntia Pelegrineti Targueta, Victor G. Dill Orrico, Luciana Bolsoni Lourenço

Data type: Tif file

Explanation note: Alignment of 16S rDNA fragments obtained from specimens of *Dendropsophus seniculus*, *D. novaisi* and *D. soaresi* that were used in cytogenetic analyses. GenBank accession numbers: KY053469 (*D. seniculus*), KY053470 (*D. novaisi*), and KY053471 (*D. soaresi*).

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.