

Electronic spectroscopy, stimulated emission, and persistent spectral hole burning of cryogenic nitrogen matrices doped with tetrabenzoporphin

S. M. Arabei

Belarusian State Agrarian and Technical University, Minsk, Belarus

J.-P. Galaup^{a)}

Laboratoire Aimé Cotton, CNRS & Université Paris-Sud 11, Orsay, France

J. G. McCaffrey

Department of Chemistry, National University of Ireland, Maynooth, Ireland

N. Shafizadeh and C. Crépin

Institut des Sciences Moléculaires d'Orsay, CNRS & Université Paris-Sud 11, Orsay, France

(Submitted April 20, 2012)

Fiz. Nizk. Temp. **38**, 917–922 (August 2012)

This paper deals with our on-going work on the electronic spectroscopy of tetrapyrrole molecules embedded in cryogenic hosts. Under nanosecond laser excitation of free-base tetrabenzoporphin molecules in a nitrogen matrix at 8 K, increasing laser pulse energy results in essential enhancement of the intensity of one or several vibronic emission lines in the fine-structure fluorescence spectrum. Some characteristics of stimulated radiation are realized in the observed intense monochromatic radiation as a result of a transition from the purely electronic S_1 level to a vibrational sublevel of the S_0 state, corresponding to the excitation of C–C stretching vibrations of tetrapyrrole methane bridges. We also report on persistent spectral hole-burning effects. Hole and antihole formation analysis gives some insight on the photochemical properties of tetrabenzoporphin in a nitrogen matrix. © 2012 American Institute of Physics. [<http://dx.doi.org/10.1063/1.4746794>]

Introduction

Development of solid-state laser-active media based on dyes continues to be one of the important problems of modern high technology. It is stimulated, first of all, by the need to create micron-size lasers. The problem of miniaturization of solid-state elements motivates the study of spectral and kinetic characteristics of spontaneous and induced (stimulated) radiation of thin-film colored materials. Studies of thin 1 μm silica gel slabs doped with dyes of different classes (rhodamines, oxazole derivatives, coumarins, porphyrins, etc.) showing lasing characteristics^{1–6} are described in the literature. It is important to note that in some experiments^{3,5} external optical resonator mirrors were missing, but stimulated emission (SE) was nevertheless detected due to light amplification in a single pass through the layer by just a “traveling wave” (traveling lasing wave).

We have performed spectroscopic studies of free base phthalocyanine (H_2Pc) and its zinc counterpart (ZnPc),⁷ as well as of free base tetrabenzoporphin (H_2TBP)⁸ in matrices of rare gases (Ne, Ar, Kr, Xe) at low temperature (8 K). We found that under the action of pulsed-laser excitation, SE was observed in guest molecules in these matrices and corresponds to transitions between the lowest singlet state S_1 and specific vibrational levels of the ground electronic state S_0 (S_1 (0–0) \rightarrow S_0 (0–1) transition). The interest in porphyrin and phthalocyanine molecules as systems capable of SE is due to their high fluorescence quantum yields, high thermal and photo stabilities, and intense absorption bands in the visible spectrum, which allow them to effectively carry out their excitation and localization of the fluorescence spectra

in the red and near-IR region. We recall that lasing emission from an organic molecule was observed for the first time in 1964 by Sorokin and Lankard:^{9,10} stimulated emission was observed when a solution of chloro-aluminum phthalocyanine dissolved in ethanol was irradiated by a beam from a giant-pulsed ruby laser. The wavelength of this stimulated emission was centered at approximately 755 nm.

The present work is a logical continuation of previously published results,^{7,8} in which the authors discussed the occurrence of the phenomenon of SE in H_2Pc , ZnPc , and H_2TBP molecules isolated in cryogenic matrices of rare gases, identifying a number of problems, so far—unsolved.

For instance, a necessity arises to clarify the role of different parameters on the stimulated emission phenomenon by changing the nature of the host matrix, then changing the electron–phonon interaction. It is still unclear why the intensification of light emission only occurs in a very limited number of vibronic transitions. Therefore, it appears necessary to establish the critical values for achieving population inversion of levels, taking into account the “distribution” of fluorescence quantum yield between the vibronic transitions (i.e., the intensity of vibronic lines), the value of the relaxation rate, and the contribution of different normal modes in the vibrational levels of the ground state S_0 .

The purpose of the present paper is to communicate new information on the spectral characteristics of H_2TBP molecules embedded in a low-temperature matrix of molecular nitrogen (N_2). The choice of H_2TBP , the structural formula for which is shown in Fig. 1, is for two main reasons. First, the quasi-line fluorescence spectra of H_2TBP in *n*-octane




FIG. 1. Absorption spectra of H₂TBP in pyridine at 293 K (1) and in a nitrogen matrix at 8 K (2). The inset shows the structure of H₂TBP.

Shpol'skii matrix at 4.2 K reveal lines assigned to individual vibronic transitions with exceptionally narrow spectral widths. For example, the components of the 0–0-doublet, recorded in the fluorescence spectrum of H₂TBP in a single-crystal matrix of *n*-octane at 4.2 K, have half-widths of 1.5 and 1.8 cm⁻¹, respectively.¹¹ Second, the role and the high speed of photochemical NH-tautomerism in H₂TBP,¹² which is the cause of efficient photoinduced interconversion of impurity centers in Shpol'skii matrices,¹³ needs to be studied further.

Experimental

In order to obtain low-temperature matrices of nitrogen doped with H₂TBP, a special device was designed and built to be attached to the cryostat. All the details have been fully described previously.⁸ The temperature of the heater, which could reach up to 300 °C, was adjusted for the H₂TBP powder. Sublimed porphyrin molecules were carried by the flow of nitrogen, forming a gaseous mixture of N₂ + H₂TBP, which was deposited on a sapphire window in a closed-cycle helium displax cryostat at ~20 K to form a thin polycrystal film with a typical thickness of about 250 μm. Gas flow rates of 10 mmol/h allowed good matrix isolation of tetrapyrrole molecules as monomers, as monitored by the absorption spectrum of the prepared sample, which also allowed to estimate the resultant concentration of the dye. Fluorescence, stimulated emission, and fluorescence excitation spectra were produced with a pulsed dye laser at 10 Hz using DCM, Rh610, or Rh6G pumped by a frequency-doubled Nd³⁺:YAG laser or an excimer XeCl laser. Fluorescence was recorded through an HRS 600 monochromator from Horiba-Jobin-Yvon using a high-speed iCCD-camera (ANDOR, model DH-720). Most of the measurements were made at about 8 K.

Results and discussion

Absorption

The absorption spectrum of H₂TBP in pyridine at room temperature is shown by the dotted trace in Fig. 1 (curve 1).

According to Refs. 11, 14–16, the band at 663.4 nm belongs to the S₁ ← S₀ transition (Q_x band), while the bands at 432.5 and 417.0 nm are assigned to the S₃ ← S₀ and S₄ ← S₀ transitions (B_x and B_y bands), respectively. The 580–620 nm region is where the Q_y band is located, and whose absorption spectrum shows the presence of three peaks. Careful analysis of the data, taking into account polarization

of the quasi-line spectra, showed that the contour of this band results from the contributions of two different types of transitions: the high-frequency vibronic transitions of the lowest S₁ electronic state, and the pure electronic S₂ ← S₀ transition. Contemplation of the nature of the violation of mirror symmetry in frequencies, as well as in the activity of individual normal modes in the S₁ and S₀ states of H₂TBP, led to the conclusion that the purely electronic band assigned to the S₂ ← S₀ transition corresponds to the central maximum of the complex recorded contour.^{14,15} Accordingly, an estimate of the energy difference $\Delta E_{S_2S_1} = E_{S_2} - E_{S_1}$ for H₂TBP in pyridine gives the value of ~1370 cm⁻¹.^{11,14,16} The same value, as well as similar band shape, were also observed for H₂TBP in DMSO. A noticeable change in the shape of the absorption band in the 580–620 nm region is observed in solvents such as *n*-octane ($\Delta E_{S_2S_1} \approx 1445$ cm⁻¹),¹⁶ ethanol ($\Delta E_{S_2S_1} \approx 1480$ cm⁻¹),¹⁷ and in a mixture of tetrahydrofuran and ether,¹⁸ where the most intense feature is in the 580–620 nm range—the short-wavelength component of the complex structure.

New information is obtained on the spectral characteristics of H₂TBP molecules, whose inhomogeneous broadening is partially removed in their low-temperature absorption spectra in a solid N₂ matrix (Fig. 1, curve 2). As can be seen, significant differences exist in the absorption spectrum of H₂TBP/N₂ and the spectrum in pyridine. Compared with a pyridine matrix, the ~10 nm blue-shift of absorption bands at 653.4 nm (=λ₀₀ of S₁ ← S₀) is indicative of a weakening in the interaction between the molecules and the surrounding N₂ matrix. Also, the half-width of the spectral bands is up to 3.5 times smaller. For example, the band of the purely electronic S₁ ← S₀ transition becomes narrower, changing from ~265 cm⁻¹ in pyridine down to ~75 cm⁻¹ in N₂. However, an accurate localization of the S₂ ← S₀ transition of H₂TBP in N₂ is not well defined, in spite of a significant intensity of the absorption bands in the 570–600 nm region. Although this cannot be determined with great accuracy, the similarity between the absorption spectra of H₂TBP in N₂ and in *n*-octane matrix is noticeable, as both these spectra have an intense band in the short wavelength range 580–620 nm. The spectral separation between this intense band and the band assigned to the S₁ ← S₀ electronic transition is equal to ~1750 cm⁻¹ in the *n*-octane matrix and ~1850 cm⁻¹ in the N₂ matrix. This observation allows us to infer that the exact $\Delta E_{S_2S_1}$ for H₂TBP/N₂ should be slightly greater than that in *n*-octane Shpol'skii matrix, for which $\Delta E_{S_2S_1} \approx 1445$ cm⁻¹. A recent investigation of phthalocyanine in inert gas matrices¹⁹ has similarly concluded that $\Delta E_{S_2S_1}$ was slightly larger for H₂Pc/N₂ than for H₂Pc trapped in the *n*-decane Shpol'skii matrix.

Stimulated emission and its excitation

Upon excitation in the Q_y absorption band at λ_{exc} = 582.7 nm with the pulsed dye laser (P_{exc} ≈ 10 mW, pulse length = 10 ns), the fluorescence spectrum of H₂TBP in N₂ at 8 K has the form shown in Fig. 2 (curve 1). An intense band assigned to the pure electronic S₁ → S₀ transition occurs at 653.8 nm, followed by low-intensity bands corresponding to vibronic transitions covering the 660–740 nm region. The resulting spectrum resembles the quasi-line




FIG. 2. Fluorescence and stimulated emission spectra of H₂TBP molecules in a nitrogen matrix at 8 K ($\lambda_{exc} = 582.7$ nm, $P_{laser} \approx 10$ (1), 30 (2) and 100 (3) mW).

fluorescence spectrum of H₂TBP in *n*-octane at 4.2 K,¹⁶ where the most intense vibronic band is a band corresponding to a low-frequency normal mode at ~ 220 cm⁻¹ (Fig. 2, see the band at 663.0 nm). As the power of pulsed laser excitation is increased from $P_{laser} \approx 30$ mW to 100 mW, two intense lines (730.7 and 731.0 nm) from the weak vibronic band at 731 nm of the normal fluorescence spectrum increase drastically (Fig. 2, compare curves 2 and 3). Attention is drawn to the sharp narrowing of these lines (down to ~ 5 cm⁻¹, which is the resolution limit of our spectrometer) and to their dependence on the relative intensities of laser excitation power. These lines are assigned to SE. Changing the laser excitation wavelength in the region 580–585 nm does not lead to the appearance of other new SE lines in the recorded spectrum. There is only a slight difference in the relative intensities of the lines at 730.7 and 731.0 nm. Similarly to the observations reported in Refs. 7 and 8, we can conclude that emission at these wavelengths is strongly enhanced when the exciting laser intensity exceeds a certain threshold value. This means that an inversion of population is achieved between the lowest electronic state S_1 and some vibrational sublevels of the ground state S_0 . When such a situation arises, stimulated emission can occur and overpass any spontaneous fluorescence. In the case of H₂TBP/N₂, the analysis of the spectral data shows that SE takes place on a vibronic transition that involves a vibrational sublevel with energy of about 1615 cm⁻¹ from the ground S_0 state. This vibrational sublevel corresponds to the excitation of valence vibrations involving the C _{α} C _{m} methine bridges (see Fig. 1) with B_{1g} symmetry.^{11,20} It is important to note that SE is observed under pulsed laser excitation, with whose duration it does not exceed the lifetime of the S_1 state (for example, in the case of H₂TBP in an ethanol-ether mixture, $\tau_S = 11$ ns (Ref. 21)), and at a repetition rate of 10 Hz, i.e., 100 ms between successive pulses. Such a situation precludes an effective population of the triplet state, well known for its role as “traps” for the active sites, which should be the case under steady-state photoexcitation. It is then difficult to achieve conditions for population inversion.

In order to clarify the conditions under which SE occurs in H₂TBP/N₂ we studied the excitation spectra of stimulated emission. Figure 3(a) shows such spectra for the Q_y band region at 8 K (curve 1, monitoring emission at $\lambda_{obs} = 730.7$ nm, and curve 2 for $\lambda_{obs} = 731.0$ nm), which is compared with the absorption spectrum of H₂TBP/N₂ (curve 3) in the same




FIG. 3. Comparison of the excitation spectra of stimulated emission (1 and 2) and the absorption spectrum (3) of H₂TBP molecules in a nitrogen matrix at 8 K for $S_2 \leftarrow S_0$ ($\lambda_{obs} = 730.7$ (1) and 731.0 (2) nm) (a) and $S_1 \leftarrow S_0$ ($\lambda_{obs} = 730.7$ (1) and 731.1 (2) nm) (b).

region. With the detection of emission in each of the two observed SE components it can be seen that the contours of the bands in the excitation spectra of SE are almost identical, and their spectral shift relative to each other is negligible. However, the maxima of the bands in the excitation spectra of SE do not coincide with the maximum absorption spectrum in this area, observed at 582.9 nm. This can be considered as a consequence of the population inversion, induced at the wavelengths 580–585 nm and excitation powers of up to 100 mW, which was used for the two types of impurity centers. It should be noted that SE intensity at 730.7 nm is higher than SE intensity at 731.0 nm. This is true for almost all excitation wavelengths. An explanation of the observed features of the excitation spectra of SE for each type of impurity centers requires a detailed analysis of the electronic and electronic-vibrational levels of the Q_x and Q_y states involved in the active transitions.^{11,16}

Stimulated emission around 731 nm is also observed under pulsed laser excitation in the long wavelength absorption Q_x band of H₂TBP/N₂ (Fig. 4). However, unlike in the




FIG. 4. Fluorescence and stimulated emission spectra of H₂TBP molecules in a nitrogen matrix at 8 K ($\lambda_{exc} = 652.7$ (a) and 654.0 (b) nm, $P_{laser} \approx 10$ (1), 30 (2) and 100 (3) mW).

case of excitation in the Q_y band, in this situation there is a dependence of the number of SE components on the change in λ_{exc} within the Q_x absorption band (Fig. 3(b)), the spectral position of the laser exciting radiation is shown by the arrows and vertical dotted lines).

At $\lambda_{\text{exc}} = 652.7$ nm and under laser excitation power gradually increasing up to $P_{\text{laser}} \approx 100$ mW, four SE components appear in the emission spectrum (Fig. 4(a)), while at $\lambda_{\text{exc}} = 654.0$ nm only one SE component is intensified (Fig. 4(b)). An explanation of this behavior observed in the spectral composition of the SE is provided by an analysis of the excitation spectra of the SE in the Q_x band absorption (Fig. 3(b)). As shown in Fig. 3(b), the broad Q_x absorption band of $\text{H}_2\text{TBP}/\text{N}_2$ at 8 K (curve 3) is transformed into a structured band in the SE excitation spectrum.

For example, it can be noticed that for $\lambda_{\text{obs}} = 731.1$ nm, the SE emission line component is on the long-wavelength side of the excitation spectrum of the SE (curve 2); there is a narrow intense line at 653.6 nm and a broad and structured band on its short-wavelength side. Clearly, such a structure in the SE excitation spectrum is formed by the excitation (absorption) of the narrow zero-phonon line (ZPL) corresponding to the $S_1 \leftarrow S_0$ (0,0) transition and to its broad phonon side band (PSB). A similar observation can be made regarding the shorter-wavelength component of the SE spectrum recorded at $\lambda_{\text{obs}} = 730.7$ nm (curve 1), while a slight blue shift can also be noticed. The observed SE excitation spectra may explain the dependence on λ_{exc} of the structured components. The excitation on the short-wavelength side within the Q_x band induces a simultaneous photoactivation of “red” (via their PSB) and “blue” (through their ZPL) kinds of impurity centers, whereas the excitation on the long-wavelength side affects only the “red” impurity centers through their ZPL. Indeed, in the first case, four types of impurity centers are identified, exhibiting SE at 731.1, 730.7, 730.3 and 730.0 nm, respectively (Fig. 4(a)), while in the second case, only one SE component at 731.1 nm can be induced (Fig. 4(b)). Notice that in the resulting SE excitation spectra shown in Fig. 3(b), SE is observed upon excitation in the Q_x (0–0) band of the purely electronic transition, and also in the vibronic Q_x (0–1) band at ~ 644 nm.

The obtained results demonstrate that under relatively high-power pulsed laser excitation in the Q_x and Q_y band of $\text{H}_2\text{TBP}/\text{N}_2$, compared with the normal fluorescence excitation spectrum, in the SE excitation spectra a sharp narrowing of lines is observed, the latter “exhibiting” thereby a greater selectivity of individual types of impurity centers in the matrix under study. These high-resolution SE spectra will allow more detailed studies of the spectral properties of certain types of impurity centers in a given matrix. Understanding the nature of the multiple components of SE may be useful for identifying the main types of impurity centers of pigments in cryogenic matrices. Moreover, such studies have a methodological significance and are of great importance as a method for selective low-temperature molecular spectroscopy.

Hole-burning

In this work we also carried out investigations of the photochemical properties of the $\text{H}_2\text{TBP}/\text{N}_2$ system via a method involving the photoburning of stable spectral holes

(persistent spectral hole burning—PSHB) in the Q_x absorption band. The phenomenon of spectral hole-burning at low temperature is quite well known and has been studied in detail for the case of free base porphyrins (see Ref. 22 for a review). It is due to the intramolecular NH-tautomerism, which in porphyrins implies only the displacement of the two central protons from one pair of opposite nitrogen atoms to the other (see structure in Fig. 1).²³

The experimental results described previously have shown that the Q_x (0–0) band of $\text{H}_2\text{TBP}/\text{N}_2$ is structureless, i.e., inhomogeneously broadened due to the presence in the matrix of a large number of impurity centers and to the superposition of their different individual spectral bands. This inhomogeneous broadening can be eliminated through exposure to monochromatic laser light. Figure 5 presents the effect of such laser irradiation on the Q_x band (Fig. 5(a)). The difference spectrum, obtained by subtracting the burnt spectrum from the unburnt one, is also shown (Fig. 5(b)). PSHB experiments then show us that burning at $\lambda_{\text{burn}} = 652.7$ nm (Fig. 5(a)) forms a stable spectral hole corresponding to the zero-phonon line (zero-phonon hole), which can be saturated with increasing burning time. Typical laser exposure time for the observation of the spectral changes was chosen to be about 10 min, and laser excitation power for the burning was ~ 100 mW. This burning time was sufficient to achieve saturation of the zero-phonon hole, which then resulted in its broadening up to $\sim 10 \text{ cm}^{-1}$. As can be seen from Fig. 5, where the effect of laser excitation on the short-wavelength side of the ZPL is observed, there is a marked increase in absorption intensity due to the formation of a photoproduct (giving rise to an “antihole”) on the high-energy side from λ_{burn} . The formation of an “antihole” during photoburning is associated with the formation of NH-tautomers of H_2TBP . This asymmetrical spectral distribution of the “antihole” due to the phototransformation of




FIG. 5. (a) Q_x (0–0) absorption band of H_2TBP molecules in a nitrogen matrix at 8 K before (1) and after burning (2), and (b) the difference of the two spectra showing a persistent spectral hole ($\lambda_{\text{burn}} = 652.7$ nm, $P_{\text{burn}} \approx 100$ mW, $t_{\text{burn}} \approx 10$ min).

H₂TBP symmetric molecules (point group symmetry D_{2h}) is a consequence of the existence of a rather intense pseudo-phonon wing at long wavelengths resulting from the burning of impurity centers, which absorb laser radiation in the region of their phonon wing, i.e., centers with $\lambda_{ZPL} > \lambda_{burn}$. Naturally, at $\lambda_{exc} = 652.7$ nm there would be more centers excited through their extensive phonon band than through their ZPL. As result, the long wavelength (with respect to the selected λ_{burn}) photoproduct absorption and the intense pseudo-phonon wing at $\lambda > \lambda_{burn}$ almost cancel out each other. It is essential to notice that similar spectral changes were also observed for the Q_x (0–1) band located at ~ 643.6 nm, which means, for the band that corresponds to the electron-vibrational transition.

As it turned out in the case of H₂TBP/N₂, the intersystem conversion in impurity centers has a rather low efficiency, which is the reason for the almost complete lack of influence of the photochemical transformation on the efficiency of the SE phenomenon.

Conclusion

The main results obtained in the present work on absorption, fluorescence (stimulated emission), and persistent spectral hole-burning of matrix-isolated molecules of H₂TBP in solid nitrogen can be summarized as follows: It was found that the exposure to pulsed laser excitation in the Q_x and Q_y absorption bands of H₂TBP molecules, trapped in the low-temperature nitrogen matrix, induces the occurrence of stimulated emission, together with normal spontaneous emission of fluorescent H₂TBP molecules. This stimulated emission is evidenced by the strong enhancement of the intensity of one or more vibronic fluorescence lines, and it occurs when the intensity of the exciting laser radiation exceeds a certain threshold. The observation of such phenomenon means the achievement of a population inversion between S_1 (0–0) ground level and some vibronic sublevel of the ground electronic state. The vibronic transition involved in the process of stimulated emission corresponds to a vibrational level of S_0 state with an energy of ~ 1615 cm⁻¹, which is assigned to the stretching vibrations of $C_\alpha C_m$ methine bridges with B_{1g} symmetry.

The observation of stimulated emission was possible using pulsed laser excitation with nanosecond duration, not exceeding the lifetime of the lowest singlet S_1 state of H₂TBP, and a 10 Hz low repetition rate. In such a situation, the intersystem singlet-triplet conversion mechanism populating the lowest triplet state (a “trap” of the excited molecules) will only play a minor role, which is an important condition for the creation of population inversion in the system of singlet levels during the laser pulse.

The recorded spectra of the excitation of stimulated emission for individual impurity centers exhibit enhanced selectivity, which creates conditions for a more detailed study of the spectral properties of the molecules forming these impurity centers in the matrix.

The obtained results determine the scientific significance of these investigations. For instance, a practical significance of these results is our ability to identify opportunities for applications in the development of laboratory models of new

thin-film materials based on low-temperature matrices colored with tetrapyrrole compounds for creating microlasers and light transformer devices (in the bluegreen region) from the monochromatic radiation in the red or the near infrared region. The results obtained can be used to expand the search for new microlaser media based on organic dyes.

This work was supported by BRFFR (projects F09F-001 and F11F-001) and CNRS France (Project Nos. 23 181 and 24 692), and the Ulysses France-Ireland exchange (PHC program Ulysses No. 25709VA).

^{a)}Email: jean-pierre.galaup@lac.u-psud.fr

¹R. Reisfeld, E. Yariv, and H. Vinti, *Opt. Mater.* **8**, 31 (1997).

²E. Yariv and R. Reisfeld, *Opt. Mater.* **13**, 49 (1999).

³R. T. Kuznetsova, Yu. A. Manekina, E. N. Tel'minov, G. V. Mayer, S. M. Arabei, T. A. Pavich, and K. N. Solovyov, *Opt. Atmos. Okeana* **19**, 653 (2006) [*Atmos. Oceanic Opt.* **19**, 586 (2006)].

⁴R. T. Kuznetsova, G. V. Maier, Yu. A. Manekina, V. A. Svetlichnyi, E. N. Tel'minov, S. M. Arabei, T. A. Pavich, and K. N. Solovyov, *Opt. Spektrosk.* **102**, 241 (2007) [*Opt. Spectrosc.* **102**, 208 (2007)].

⁵R. T. Kuznetsova, G. V. Maier, Yu. A. Manekina, E. N. Tel'minov, S. M. Arabei, T. A. Pavich, and K. N. Solovyov, *Kvantovaya Élektron.* **37**, 760 (2007) [*Quantum Electron.* **37**, 760 (2007)].

⁶R. T. Kuznetsova, G. V. Mayer, Yu. A. Manekina, E. N. Tel'minov, S. M. Arabei, T. A. Pavich, and K. N. Solovyov, *Opt. Spektrosk.* **104**, 223 (2008) [*Opt. Spectrosc.* **104**, 190 (2008)].

⁷N. Dozova, C. Murray, J. G. McCaffrey, N. Shafizadeh, and C. Crépin, *Phys. Chem. Chem. Phys.* **10**, 2167 (2008).

⁸C. Crépin, N. Shafizadeh, W. Chin, J.-P. Galaup, J. G. McCaffrey, and S. M. Arabei, *Fiz. Nizk. Temp.* **36**, 563 (2010) [*Low Temp. Phys.* **36**, 451 (2010)].

⁹P. P. Sorokin and J. R. Lankard, *IBM J. Res. Dev.* **10**, 162 (1966).

¹⁰P. P. Sorokin, J. R. Lankard, E. C. Hammond, and V. L. Moruzzi, *IBM J. Res. Dev.* **11**, 130 (1967).

¹¹S. M. Arabei, K. N. Solovyov, and Yu. I. Tatulchenkov, *Opt. Spektrosk.* **73**, 686 (1992) [*Opt. Spectrosc.* **73**, 406 (1992)].

¹²I. E. Zaleski, V. N. Kotlov, A. N. Sevchenko, K. N. Solovyov, and S. F. Shkirman, *Dokl. Akad. Nauk SSSR* **207**, 1314 (1972) [*Proc. USSR Acad. Sci. Sov. Phys. Dokl.* **17**, 1183 (1973)].

¹³K. N. Solovyov, I. E. Zaleski, V. N. Kotlov, and S. F. Shkirman, *Pisma Zh. Eksp. Teor. Fiz.* **17**, 463 (1973) [*JETP Lett.* **17**, 332 (1973)].

¹⁴K. N. Solovyov, S. F. Shkirman, and T. F. Kachura, *Izv. Akad. Nauk SSSR Ser. Fiz.* **27**, 767 (1963) [*Bull. Acad. Sci. USSR, Phys. Ser.* **27**, 763 (1963)].

¹⁵A. N. Sevchenko, K. N. Solovyov, S. F. Shkirman, and T. F. Kachura, *Dokl. Akad. Nauk SSSR* **161**, 1313 (1965) [*Proc. USSR Acad. Sci. Sov. Phys. Dokl.* **10**, 349 (1965)].

¹⁶S. M. Arabei, *Zh. Prikl. Spektrosk.* **57**, 56 (1992) [*J. Appl. Spectrosc.* **57**, 572 (1992)].

¹⁷B. Ehrenberg and F. M. Johnson, *Spectrochim. Acta, A* **46**, 1521 (1990).

¹⁸V. G. Maslov, *Opt. Spektrosk.* **50**, 1091 (1981) [*Opt. Spectrosc.* **50**, 599 (1981)].

¹⁹C. Murray, N. Dozova, J. G. McCaffrey, N. Shafizadeh, W. Chin, M. Broquier, and C. Crépin, *Phys. Chem. Chem. Phys.* **13**, 17543 (2011).

²⁰S. F. Shkirman, L. L. Gladkov, V. K. Konstantinova, and K. N. Solovyov, *Zh. Prikl. Spektrosk.* **66**, 375 (1999) [*J. Appl. Spectrosc.* **66**, 400 (1999)].

²¹V. A. Kuzmitsky, K. N. Solovyov, V. N. Knyukshto, I. K. Shushkevich, V. N. Kopranenkov, and A. M. Vorotnikov, *Teor. Eksp. Khim.* **19**, 655 (1983) [*Theor. Exp. Chem.* **19**, 604 (1984)].

²²J.-P. Galaup, “Spectral selective studies of molecular doped solids and applications,” in *Advances in Multi-Photon Processes and Spectroscopy*, edited by S. H. Lin, A. A. Villaeys, and Y. Fujimura (World Scientific, 2004), Vol. 16, p. 73.

²³V. A. Kuzmitsky, A. N. Sevchenko, and K. N. Solovyov, *Dokl. Akad. Nauk SSSR* **239**, 308 (1978) [*Proc. USSR Acad. Sci. Sov. Phys. Dokl.* **239**, 194 (1978)].

This article was published in English in the original Russian journal. Reproduced here with stylistic changes by AIP.