

HOMOTOPY CLASSES WITH SMALL JACOBIANS

STEFAN BECHTLUFT-SACHS

Abstract

If the infimum of the conformal k -Jacobian on the homotopy class of a map between compact Riemannian manifolds vanishes then the map factors rationally through the k -skeleton of the target manifold.

AMS Subject Classification: 58E15 (53C43 57D99)

1. Introduction

It follows from the Sobolev inequality that a map $f: M^m \rightarrow X$ between connected compact Riemannian manifolds M and X , $m = \dim(M)$, is nullhomotopic if its differential df has sufficiently small L^r -norm for some $r > m$. In fact, the diameter of the image of f is bounded by $\text{diam}(\text{im}(f)) \leq C \|df\|_r$ with some constant C depending only on M , X and $r > m$. In the conformal case $r = m$ this simple argument fails. A map f with arbitrary small $\|df\|_m$ can have arbitrarily large image. But by a theorem of White [10] there is a constant $\epsilon > 0$ depending on the geometries of M and X such that f is nullhomotopic if $\|df\|_m < \epsilon$.

We consider the analogous question for the Jacobian in place of the L^r -norm. For $k \in \mathbb{N}$ and $r \in \mathbb{R}^+$ these are the functionals

$$(1) \quad J_k^r: \mathcal{C}^\infty(M, X) \rightarrow \mathbb{R}_0^+, \quad J_k^r(f) = \int_M \phi(df)$$

where

$$\phi(df) = \underbrace{|df \wedge \dots \wedge df|}_{k}^{r/k} = \sigma(df^*df)^{r/2k}$$

and $\sigma_k(df^*df)$ denotes the k th elementary symmetric polynomial in the eigenvalues of df^*df . If $r = m = \dim(M)$ this functional is invariant under conformal changes of the metric on M .

In more general framework, for functionals $E: \mathcal{C}^\infty(M^m, X) \rightarrow \mathbb{R}_0^+$ we are interested in the information on the homotopy class of f detected by the infimum

$$\tilde{E}: [M^m, X] \rightarrow \mathbb{R}_0^+, \quad \tilde{E}(f) := \inf\{E(g) | g: M^m \rightarrow X, g \simeq f\},$$

in particular in the consequences of $\tilde{E}(f) = 0$.

We write $E_1 \gg E_2$ if $\lim_\nu E_1(f_\nu) = 0$ implies $\lim_\nu E_2(f_\nu) = 0$ for any sequence $(f_\nu)_\nu$ in $\mathcal{C}^\infty(M^m, X)$. Among the Jacobians the Hölder inequality gives estimates

$$(2) \quad J_1^r \gg J_2^r \gg \dots J_l^r \gg \dots \gg J_m^r$$

and $J_l^{r_1} \gg J_l^{r_2}$ if $r_1 \geq r_2$. With respect to \gg , the Jacobian $J_1^r(f)$ is equivalent to the L^r -norm of the differential and $J_m^m \gg \text{vol}(\text{im } f)$. If f is homotopic to a

map $\tilde{f}: M \rightarrow X^{l-1} \subset X$ into the $(l-1)$ -skeleton X^{l-1} of a triangulation of X we obviously have $\tilde{J}_l^r(f) = 0$ for any r . The converse is known to hold in the extreme cases $l = 1$ and $l = m$ of (2) if $r \geq m$. It follows from a theorem of Pluzhnikov, [7], and White, [10], that \tilde{J}_l^r depends only on the restriction of f to the $[r]$ -skeleton of a triangulation of M .

Let $f: M^m \rightarrow X$ be a map with $\tilde{J}_l^m(f) = 0$. In [2] it is shown that f behaves homologically like a map into the $(l-1)$ -skeleton X^{l-1} of a triangulation of X , i.e. induces 0 in homology of degree at least l . It is also shown there that f does not need to be homotopic to a map into X^{l-1} . Counterexamples produced in [2] arise from torsion elements in the higher homotopy groups of spheres. This suggests that f factors rationally. We prove:

Theorem 3. *Let X^{l-1} be the $(l-1)$ -skeleton of a triangulation of X and assume that $\pi_1(X^{l-1}) = 0$. Let $f: M^m \rightarrow X$ be a map with $\tilde{J}_l^m(f) = 0$. Then the rationalization $f_{\mathbb{Q}}: M \rightarrow X_{\mathbb{Q}}$ is homotopic to a map $f_{\mathbb{Q}}: M \rightarrow X_{\mathbb{Q}}^{l-1}$ into the rationalization of the $(l-1)$ -skeleton of X .*

Remarks

- (1) Theorem 3 extends a result of Rivière in [8] who showed that the Hopf invariant of a map $f: S^{4k-1} \rightarrow S^{2k}$ is estimated by \tilde{J}_{2k}^{4k-1} . For maps between spheres the rational homotopy type is controlled by the Hopf invariant.
- (2) For the Jacobians $J_l^r(f)$ with $l \geq 2$ and arbitrary large r one easily constructs surjective maps $M^m \rightarrow X^l$ with $\tilde{J}_l^m(f) = 0$. Thus a simple argument based on a Sobolev-type inequality is not available in this case.

2. Factorization in Rational Homotopy

The proof of Theorem 3 is a computation in suitable relative Sullivan algebras, along the lines of [5], [6] where the number of homotopy classes of maps f was estimated by bounds on the dilatation. As before X^{l-1} denotes the $(l-1)$ -skeleton of X . We denote by $X_{\mathbb{Q}}, X_{\mathbb{Q}}^{l-1}$ the rationalisations of X and X^{l-1} respectively. Thus we have maps $X \rightarrow X_{\mathbb{Q}}$ and $X^{l-1} \rightarrow X_{\mathbb{Q}}^{l-1}$ inducing isomorphisms $H^*(X, \mathbb{Q}) \cong H^*(X_{\mathbb{Q}}, \mathbb{Z})$ and $H^*(X^{l-1}, \mathbb{Q}) \cong H^*(X_{\mathbb{Q}}^{l-1}, \mathbb{Z})$. We assume that X^{l-1} is simply connected and $l \geq 2$. Then X is also simply connected and the above rationalisations are unique up to homotopy.

Let $\Omega(M), \Omega(X)$ and $\Omega(X^k)$ denote the respective algebras of differential forms. By the functorial properties of Sullivan algebras the rationalization $f_{\mathbb{Q}}: M \rightarrow X_{\mathbb{Q}}$ is homotopic to a map $F: M \rightarrow X_{\mathbb{Q}}^{l-1} \subset X_{\mathbb{Q}}$ if there is a relative Sullivan algebra $\mathcal{S} := \Omega(X) \otimes_d \Lambda V \simeq \Omega(X^{l-1})$ and an extension $F^*: \mathcal{S} \rightarrow \Omega(M)$ of $f^*: \Omega(X) \rightarrow \Omega(M)$, see [3], [4]. We will first construct a suitable Sullivan algebra \mathcal{S} and then use the estimate on the Jacobian of f to define F^* .

2.1. Construction of \mathcal{S}

We abbreviate $\mathcal{X} := \Omega(X)$, $\mathcal{Y} := \Omega(X^{l-1})$ and let $j: \mathcal{X} \rightarrow \mathcal{Y}$ be the morphism of commutative cochain algebras obtained by restriction. Up to homotopy we want to replace j by a morphism \tilde{j} into a relative Sullivan algebra \mathcal{S} homotopy equivalent to \mathcal{Y} such that the triangle

$$\begin{array}{ccc} \mathcal{X} & \xrightarrow{\tilde{j}} & \mathcal{S} = \mathcal{X} \otimes_d \Lambda V & \xrightarrow[m \simeq]{} & \mathcal{Y} \\ & \searrow & \text{---} & \nearrow & \\ & & j & & \end{array}$$

commutes up to homotopy. A relative Sullivan algebra ([4]) is a commutative cochain algebra $\mathcal{S} = \mathcal{X} \otimes_d \Lambda V$ such that there are graded vector spaces V_i , $i \in \mathbb{N}$,

$$V := \bigoplus_{i>0} V_i = \bigcup_{i>0} V(i), \quad V(i) := V(i-1) \oplus V_i, \quad V(-1) := 0$$

and homomorphisms

$$d_i: V_i \rightarrow \mathcal{S}(i-1) := \mathcal{X} \otimes_d \Lambda V(i-1)$$

extending to a differential $d: \mathcal{S} \rightarrow \mathcal{S}$ which is nilpotent in the sense that

$$dV(i) \subset \mathcal{S}(i-1) := \mathcal{X} \otimes_d \Lambda V(i-1) \text{ where } \mathcal{S}(-1) := \mathcal{X}.$$

The algebras $\mathcal{S}(q)$ are constructed together with morphisms $m_q: \mathcal{S}(q) \rightarrow \mathcal{Y}$ which are quasiisomorphisms in degrees increasing with q . To begin, let $V(0) \xrightarrow{\iota} \ker d \subset \mathcal{Y}$ be a vectorspace of cycles in \mathcal{Y} such that $V(0) \oplus \text{im } H_j$ generates $H(\mathcal{Y})$ as an algebra. Define

$$\mathcal{S}(0) := \mathcal{X} \otimes \Lambda V(0) \text{ with } d|_{V(0)} = 0$$

and

$$m_0: \mathcal{X} \otimes \Lambda V(0) \xrightarrow{j \otimes \Lambda \iota} \mathcal{Y}.$$

By construction m_0 induces an epimorphism $Hm_0: H(\mathcal{X}) \otimes \Lambda V(0) \rightarrow H(\mathcal{Y})$ in homology.

Proceeding by induction in q , assume $V(q)$ and

$$\mathcal{S}(q) := \mathcal{X} \otimes_d \Lambda V(q) \xrightarrow{m_q} \mathcal{Y}$$

have already been constructed such that Hm_q is surjective. Let $V_{q+1} \subset \mathcal{S}(q)$ with $V_{q+1} \cong \ker Hm_q \subset H(\mathcal{S}(q))$ be a vector space of representing cycles where the degree on V_{q+1} is set to be the degree inherited from of $\mathcal{S}(q)$ diminished by 1. The differential $d: V_{q+1} \rightarrow \mathcal{S}(q)$ is defined to be the inclusion $V_{q+1} \hookrightarrow \mathcal{S}(q)$. The map $m_{q+1}: V_{q+1} \rightarrow \mathcal{Y}$ is a lift of $m_q \circ d$ over $d^{\mathcal{Y}}$:

$$(1) \quad \begin{array}{ccccc} \ker Hm_q & \hookrightarrow & H\mathcal{S}(q) & \xrightarrow{Hm_q} & H\mathcal{Y} \\ \cong \uparrow & & \downarrow & & \downarrow \\ V_{q+1} & \xrightarrow{d_{q+1}} & \mathcal{S}(q) & \xrightarrow{m_q} & \mathcal{Y} \\ & \searrow \text{---} & & \nearrow & \uparrow d^{\mathcal{Y}} \\ & & & & \mathcal{Y} \\ & & & & \uparrow m_{q+1} \text{---} \end{array}$$

Extending m_{q+1} to $\Lambda V(q+1) = \Lambda(V_{q+1} \oplus V(q))$ we obtain $m_{q+1}: \mathcal{S}(q+1) := \mathcal{X} \otimes_d V(q+1) \rightarrow \mathcal{Y}$ which again induces an epimorphism in homology.

2.2. Extension of f

Let $f^*: \mathcal{X} = \Omega(X) \rightarrow \mathcal{M} = \Omega(M)$ and $j^*: \mathcal{X} = \Omega(X) \rightarrow \mathcal{Y} = \Omega(X^{l-1})$ be the morphisms of commutative cochain algebras induced by $f: M \rightarrow X$ and the inclusion $j: X^{l-1} \hookrightarrow X$ respectively. The homomorphism induced by j in cohomology is an isomorphism in degrees $< l-1$, injective in degree $l-1$ and 0 in degrees $> l-1$. Hence we may choose $V(0) \subset H^{l-1}(\mathcal{Y})$ such that

$$H(\mathcal{Y}) = V(0) \oplus \text{im } H j .$$

The morphism m_0 constructed as before then induces an isomorphism Hm_0 in degrees $\leq l-1$. Denote by $\mathfrak{i}(q)$ the ideal generated by $V(0)$ in $\mathcal{S}(q)$. In the diagram (1) we may split

$$V_{q+1} = V'_{q+1} \oplus d_{q+1}^{-1} \mathfrak{i}(q)$$

where V'_{q+1} lies in degrees $> l-1$.

We will inductively extend f^* to maps $f_q^*: \mathcal{S}(q) \rightarrow \mathcal{M}$ such that the f_q^* vanish on $\mathfrak{i}(q)$. To this end set

$$f_{-1}^* := f: \mathcal{S}(-1) = \mathcal{X} \rightarrow \mathcal{M}$$

and assume that we already have constructed the extension

$$f_q^*: \mathcal{S}(q) \rightarrow \mathcal{M}$$

satisfying estimates

$$(2) \quad \|f_q^* \omega\|_{n/r} \leq \epsilon \|\omega\|_{n/r}$$

for all $\omega \in \mathcal{S}(q)$ of degree r , $n \geq r > l-1$. Changing f by a homotopy we can have (2) with arbitrarily small value of ϵ .

For any r -cycle σ in M we find a homologous r -cycle σ' such that $\int_{\sigma'} |f_q^* \omega| < C_1 \int_M |f_q^* \omega|$ where C_1 does not depend on f_q^* (see [10], Proposition 3.1 for instance, or [2], proof of Theorem 3.2). If $d\omega = 0$ the Hölder inequality gives an estimate

$$\left| \int_{\sigma} f_q^* \omega \right| = \left| \int_{\sigma'} f_q^* \omega \right| \leq C_1 \|f_q^* \omega\|_1 \leq C_1 C_2 \|f_q^* \omega\|_{n/r} \leq C_1 C_2 \epsilon \|\omega\|_{n/r}$$

with C_2 independent of f . In particular $f_q^* \omega$ is exact if $d\omega = 0$.

Thus $f_q^*(dV'_{q+1}) \subset d\mathcal{M}$. Let $\{\omega_j\}_j$ be a basis for V'_{q+1} . We define f_{q+1}^* on V_{q+1} by lifting f_q^* . More precisely, choose for each ω_j some $\alpha_j \in \mathcal{M}$ satisfying $d\alpha_j = f_q^*(d\omega_j)$ and the estimate (4) of the following Lemma 3. Define $f_{q+1}^*|_{\mathfrak{i}(q)} := 0$ and $f_{q+1}^*(\omega_j) := \alpha_j$.

Lemma 3. *Let M be a compact n -dimensional Riemannian manifold and denote by $\|\cdot\|_p$ the L^p -norm of differential forms given by the Riemannian metric. There is a constant $C \in \mathbb{R}$ depending on M (but not on β) such that for each exact r -form $\beta \in \Omega^r(M)$, $\beta \in d\Omega^{r-1}(M)$ there is $\alpha \in \Omega^{r-1}(M)$ with*

$$(4) \quad \beta = d\alpha \text{ and } \|\alpha\|_{n/(r-1)} \leq C \|\beta\|_{n/r} .$$

Proof: From the Hodge decomposition

$$\Omega(M) = \ker \Delta \oplus \operatorname{im} d \oplus \operatorname{im} d^*$$

the Laplacian Δ is invertible on $\operatorname{im} d \oplus \operatorname{im} d^*$. Let $\alpha := d^* \Delta^{-1} \beta$. Clearly $d\alpha = \beta$. Extending the operator Δ^{-1} on $\operatorname{im} d \oplus \operatorname{im} d^*$ by 0 to all of $\Omega^r(M)$ yields a bounded operator $L^p = W^{0,p} \rightarrow W^{2,p}$ into the Sobolev space $W^{2,p}$, [9]. Also $d^*: W^{2,p} \rightarrow W^{1,p}$ is bounded, [1]. From the Sobolev-embedding $\iota: W^{1,p} \subset L^{np/(n-p)}$ we infer that

$$\|\alpha\|_{np/(n-p)} \leq C \|\beta\|_p$$

where $C := \|\iota d^* \Delta^{-1}\|_{\text{op}}$ is the operator norm. With $p = n/r$, $np/(n-p) = n/(r-1)$ the assertion follows. •

References

- [1] Adams, Robert A. *Sobolev spaces* Pure and Applied Mathematics, Vol. 65. Academic Press, New York-London, 1975.
- [2] Bechtluft-Sachs, S.; *Infima of Universal Energy Functionals on Homotopy Classes*, to appear in *Mathematische Nachrichten*.
- [3] Griffiths, Phillip A.; Morgan, John W. *Rational homotopy theory and differential forms* Progress in Mathematics, 16. Birkhäuser, 1981.
- [4] Félix, Yves; Halperin, Stephen; Thomas, Jean-Claude *Rational homotopy theory* Graduate Texts in Mathematics, 205. Springer-Verlag, 2001.
- [5] Gromov, M. *Metric structures for Riemannian and non-Riemannian spaces*, Progress in Mathematics 152, Birkhäuser, 1999.
- [6] Gromov, M. *Homotopical effects of dilatation* J. Differential Geom. 13 (1978), no. 3, 303–310.
- [7] Pluzhnikov, A. I. *Topological aspects of the problem of minimizing the Dirichlet functional*, Proc. Steklov Inst. Math. 1993, no. 3 (193), 167–171.
- [8] Rivière, T. *Minimizing fibrations and p-harmonic maps in homotopy classes from S^3 into S^2* , Comm. Anal. Geom. 6 (1998), no. 3, 427–483.
- [9] Taylor, Michael E. *Partial differential equations III. Nonlinear equations* Applied Mathematical Sciences, 117. Springer-Verlag, New York, 1997.
- [10] White, B. *Infima of energy functionals in homotopy classes of mappings*, J. Differential Geom. 23 (1986), no. 2, 127–142.