



# The effect of the nitrification inhibitor dicyandiamide (DCD) on herbage production when applied at different times and rates in the autumn and winter

P.J. O'Connor<sup>a,b</sup>, D. Hennessy<sup>a,\*</sup>, C. Brophy<sup>c</sup>, M. O'Donovan<sup>a</sup>, M.B. Lynch<sup>b</sup>

<sup>a</sup> Teagasc, Animal & Grassland Research and Innovation Centre, Moorepark, Fermoy, Co. Cork, Ireland

<sup>b</sup> UCD School of Agriculture and Food Science, University College Dublin, Belfield, Dublin 4, Ireland

<sup>c</sup> Department of Mathematics & Statistics, National University of Ireland Maynooth, Maynooth, Co. Kildare, Ireland

## ARTICLE INFO

### Article history:

Received 15 June 2011

Received in revised form 15 February 2012

Accepted 19 February 2012

### Keywords:

Herbage production  
Nitrification inhibitor  
Dicyandiamide  
Urine  
Soil mineral N  
N uptake

## ABSTRACT

The high rate of urine excreted during animal grazing in late autumn provides a source of nitrogen (N) to the growing sward and also provides the potential for losses of N over the winter months. This study was established to evaluate the potential of applying a nitrification inhibitor, dicyandiamide (DCD), to urine patches to increase N use efficiency in grassland. Four simulated grazing plot experiments were undertaken across two experimental sites, one a free-draining acid brown earth (Experiments 1 and 3) and the other a moderate to heavy brown earth soil (Experiments 2 and 4). Experiments 1 and 2 received no fertiliser N application, and Experiments 3 and 4 received a split application of 350 kg N fertiliser ha<sup>-1</sup> year<sup>-1</sup>. The effect of applying the nitrification inhibitor dicyandiamide (DCD) at 5 or 10 kg DCD ha<sup>-1</sup> in autumn and winter to plots receiving synthetic urine or zero urine on spring and annual herbage production was examined in all experiments. The application of DCD did not increase spring herbage production in any of the experiments. Over the two years, the application of 5 or 10 kg DCD ha<sup>-1</sup> increased annual herbage production in Experiment 1 when applied to October and November deposited urine patches. Urine application increased herbage production in spring and annually in Experiments 1 and 2, and increased herbage crude protein content and herbage N uptake in all experiments. The application of urine increased soil ammonium and TON content in the 0–100 mm horizon at both sites. The application of 10 kg DCD ha<sup>-1</sup> reduced surplus N in Experiment 1 when applied to October and November deposited urine. Overall the effects of DCD on herbage production, surplus N and other parameters in this study were not consistent.

© 2012 Elsevier B.V. All rights reserved.

## 1. Introduction

Increasing the proportion of grazed grass in the diet of the dairy cow, particularly in early spring, reduces milk production costs and can increase the profitability of grass based milk production systems in Ireland and other temperate climates (Shalloo et al., 2004; Dillon et al., 2005; Kennedy et al., 2005). Nitrogen (N) availability is one of the key factors driving grass growth. Increasing N availability in spring through fertiliser or slurry application can result in increased grass growth and therefore herbage availability for grazing. Urine and dung are also sources of N in grazed swards, although their deposition is localised. Nitrogen concentration under urine patches is very high, equivalent to a fertiliser N application rate of up to 1000 kg N ha<sup>-1</sup> (Whitehead, 1995). The majority of this N is in excess of sward requirements and is often lost by nitrate (NO<sub>3</sub><sup>-</sup>)

leaching through the soil profile or nitrous oxide (N<sub>2</sub>O) emissions, particularly over winter when grass growth rates are low.

Nitrification inhibitors are being investigated in many countries as a strategy to mitigate NO<sub>3</sub><sup>-</sup> leaching, denitrification and N<sub>2</sub>O emissions under urine patches (Serna et al., 1995). They therefore have the potential to increase N availability in the soil for grass growth, thereby increasing the N use efficiency of grazed swards, as well as reducing N losses to the environment (O'Connell et al., 2004). Dicyandiamide (DCD; C<sub>2</sub>H<sub>4</sub>N<sub>4</sub>) is one such nitrification inhibitor. It is a white crystalline nitrogenous powder naturally broken down in the soil, with no traces of residue remaining beyond the cropping year (Amberger, 1989). Dicyandiamide slows the conversion of ammonium (NH<sub>4</sub><sup>+</sup>) to NO<sub>3</sub><sup>-</sup> in the soil by interfering with the cytochrome oxidase in the respiratory electron transport system of *Nitrosomonas* bacteria, which are responsible for the first step of the nitrification process (Serna et al., 1995). Reductions in NO<sub>3</sub><sup>-</sup> leaching and N<sub>2</sub>O emissions following the application of DCD have been reported by many authors including Moir et al. (2007) and Dennis et al. (2008). Di and Cameron (2002) reported reductions in annual NO<sub>3</sub><sup>-</sup> leaching of 59% from urine

\* Corresponding author. Tel.: +353 025 42297; fax: +353 025 4234.

E-mail address: [deirdre.hennessy@teagasc.ie](mailto:deirdre.hennessy@teagasc.ie) (D. Hennessy).

patches (lysimeter study) following DCD application; in addition Di and Cameron (2005) reported a 68% reduction in  $\text{NO}_3^-$  leaching from dairy cow urine N when DCD was applied in autumn (Di and Cameron, 2005). Selbie et al. (2011) reported that DCD ( $10 \text{ kg ha}^{-1}$ ) reduced  $\text{NO}_3^-$  leaching by 45% and  $\text{N}_2\text{O}$  emissions by 70% on dairy cow urine ( $1000 \text{ kg N ha}^{-1}$ ) treatments on Irish soils. Richards et al. (2008) also reported that DCD application on urine patches reduced  $\text{NO}_3^-$  leaching, especially on Irish soils.

In addition to the reductions in environmental N losses, Di and Cameron (2002) also observed an 18% increase in herbage production following DCD application to spring deposited urine patches, and an average increase in herbage production of 49% following DCD application to autumn deposited urine patches. The same authors observed that the application of DCD to autumn deposited urine patches, followed by a second application in spring, increased herbage production by 33% annually (Di and Cameron, 2005). Zaman and Blennerhassett (2010) reported that the application of DCD to spring deposited urine increased herbage production by an average of 12%.

An N balance can be used to describe the potential for N loss to the environment; it gives an indication of the quantity of N that may be lost through leaching, denitrification and volatilization or immobilization into soil organic N. Research has identified the importance of some individual loss processes, such as N losses by ammonia volatilization in pastures grazed by dairy cows (Bussink and Oenema, 1996),  $\text{NO}_3^-$ -N leaching (Scholefield et al., 1993),  $\text{N}_2\text{O}$  emissions and N removal by immobilization (Ledgard et al., 1999).

The efficacy of DCD is influenced by several factors, including soil and environmental factors. Temperature is the most influential environmental factor; an increase in temperature can have a negative effect on the persistence of DCD in the soil, reducing the time frame in which it can provide effective nitrification inhibition. The half life of DCD at  $6^\circ\text{C}$  is 100 days (Williamson et al., 1996), and 18–25 days at  $20^\circ\text{C}$  (Di and Cameron, 2004). As a consequence, DCD should be applied in cool conditions such as late autumn, winter and early spring in temperate climates to maximise its potential effectiveness in inhibiting nitrification in the soil (Kelliher et al., 2008). Therefore, the hypothesis of this experiment was that the application of DCD in autumn and winter will increase N availability for grass growth in spring and, therefore, increase spring herbage production.

The objective of this study was to establish if herbage production was increased following the application of DCD to autumn and winter deposited urine patches, and to determine the appropriate rate and time of application of DCD on grass swards receiving zero or  $350 \text{ kg N ha}^{-1} \text{ year}^{-1}$ .

## 2. Materials and methods

### 2.1. Soil type and pasture

Four experiments were undertaken using simulated grazing plots at the Animal and Grassland Research and Innovation Centre, Teagasc, Moorepark, Fermoy, Co. Cork, Ireland, on two contrasting soil types. The soils were a free-draining acid brown earth of sandy loam to loam in texture with a pH of 6.02 and bulk density of  $1.00 \text{ g cm}^{-3}$  at Moorepark Research Farm, hereafter referred to as MPK ( $50^\circ 07' \text{N}$ ,  $08^\circ 16' \text{W}$ ) in Experiments 1 and 3 and a moderate to heavy brown earth with evidence of an iron pan with a pH of 5.52 and bulk density of  $0.83 \text{ g cm}^{-3}$  at Ballydague Research Farm, hereafter referred to as BD ( $52^\circ 12' \text{N}$ ,  $08^\circ 13' \text{W}$ ) in Experiments 2 and 4. Ballydague Research Farm is approximately 8 km from MPK. Swards were predominately perennial ryegrass (*Lolium perenne* L.) and were previously rotationally grazed by dairy cows at MPK and by dairy heifers at BD. In Experiments 1 and 2 no fertiliser N was

applied, and in Experiments 3 and 4 split applications of N fertiliser (total  $350 \text{ kg N ha}^{-1} \text{ year}^{-1}$ ) were applied between mid-January and mid-September (as specified in the Irish Governments Nitrates Action Plan (S.I. 378, 2006)) as urea (46% N) from February to April ( $120 \text{ kg N ha}^{-1} \text{ year}^{-1}$ ), and calcium ammonium nitrate (CAN; 27% N) from May to September ( $230 \text{ kg N ha}^{-1} \text{ year}^{-1}$ ).

### 2.2. Experimental design

Apart from the N fertiliser application strategy (zero N in Experiments 1 and 2, and  $350 \text{ kg N ha}^{-1}$  in Experiments 3 and 4), all experiments had the same design. Each experiment had three replications (blocks), and within each replication five factors (details below) were manipulated resulting in 28 treatments in total. The 28 treatments were each applied to one plot within each block; plots were  $5 \text{ m} \times 1 \text{ m}$  at BD and  $5 \text{ m} \times 1.5 \text{ m}$  at MPK. Experiments were established in September 2008 and completed in November 2010. A cleaning cut was undertaken in November 2008 and the recording of experimental data began in February 2009. Each plot was harvested ten times in year 1 (2009) and eight times in year 2 (2010).

### 2.3. Treatments

Within each experiment, the five factors manipulated were urine rate, date of application of urine, DCD rate, date of first application of DCD and date of second application of DCD (Table 1). The two urine application rates were 0 and  $1000 \text{ kg N ha}^{-1}$  (OU or U, respectively) applied on one of three occasions in autumn – late September, October or November. Synthetic urine was used (urea and water mix) so that a known quantity of N was applied. Synthetic urine was deposited using 10 L watering cans with rose caps (cap with small openings). Dicyandiamide was applied at rates of 0, 5 and  $10 \text{ kg ha}^{-1}$  to all designated plots within 24 h of urine application in either late September, October or November. Of the plots that received a first application of  $\text{DCD} > 0 \text{ kg ha}^{-1}$ , half received a second application of the same rate approximately 90 days later. The DCD was applied as a fine particle suspension (FPS) using a walk behind sprayer (Kestrel Spray-Master Sprayer, R&J Hay, St Johnston, Cavanacaw, Co. Donegal, Ireland).

Plots receiving DCD at rates of 0, 5 and  $10 \text{ kg ha}^{-1}$  will hereafter be referred to as 0, 5 and 10. Plots receiving a single application of urine and DCD in late September, October or November will hereafter be referred to as S, O or N, and those receiving a second application of DCD 90 days later will hereafter be referred to as S+90, O+90 and N+90. Plots that received zero urine and zero DCD are called control and hereafter be referred to as C. In total there are 28 unique combinations of factor levels or treatments (including control) listed in Table 1. Treatments applied in year 1 were applied between September 2008 and March 2009 and treatments applied in year 2 were applied between September 2009 and March 2010.

### 2.4. Measurements

#### 2.4.1. Herbage production

Herbage was mechanically harvested every four weeks from February to November 2009 and in 2010 using an Agria auto-scythe mower (Agria Werke GmbH, Bittelbronnerstr 42, Moeckmuehl 74219, Germany) at MPK and a Honda rotary blade lawnmower (Honda HRH 536 HX Pro Hydrostatic 4-wheel mower, Honda, Swepsonville, NC, USA) at BD. The Honda lawnmower was used at BD due to the heavy soil at the site (Hennessy et al., 2008). All fresh samples were weighed and a sub-sample (100 g) was dried at  $40^\circ\text{C}$  for 48 h to determine dry matter (DM) content. Dry matter yield was calculated by multiplying DM% of the subsample by

**Table 1**

Experimental treatments 1–28; for each of the four experiments, each treatment appeared once in each of three blocks.

| Treatment # | Treatment code | Urine (kg N ha <sup>-1</sup> ) | Date of urine application <sup>a</sup> | DCD rate (kg ha <sup>-1</sup> ) | Date first DCD application <sup>a,b</sup> | Date of second DCD application (days post first application) <sup>a</sup> |
|-------------|----------------|--------------------------------|--|---------------------------------|---|---|
| 1           | C              | 0                              | 0                                      | 0                               | 0   | 0   |
| 2           | S0U5           | 0                              | 0                                      | 5                               | September                                 | 0   |
| 3           | S0U10          | 0                              | 0                                      | 10                              | September                                 | 0   |
| 4           | S+900U5        | 0                              | 0                                      | 5                               | September                                 | +90   |
| 5           | S+900U10       | 0                              | 0                                      | 10                              | September                                 | +90   |
| 6           | O0U5           | 0                              | 0                                      | 5                               | October                                   | 0   |
| 7           | O0U10          | 0                              | 0                                      | 10                              | October                                   | 0   |
| 8           | O+900U5        | 0                              | 0                                      | 5                               | October                                   | +90   |
| 9           | O+900U10       | 0                              | 0                                      | 10                              | October                                   | +90   |
| 10          | N0U5           | 0                              | 0                                      | 5                               | November                                  | 0   |
| 11          | N0U10          | 0                              | 0                                      | 10                              | November                                  | 0   |
| 12          | N+900U5        | 0                              | 0                                      | 5                               | November                                  | +90   |
| 13          | N+900U10       | 0                              | 0                                      | 10                              | November                                  | +90   |
| 14          | SU0            | 1000                           | September                              | 0                               | 0   | 0   |
| 15          | SU5            | 1000                           | September                              | 5                               | September                                 | 0   |
| 16          | SU10           | 1000                           | September                              | 10                              | September                                 | 0   |
| 17          | S+90U5         | 1000                           | September                              | 5                               | September                                 | +90   |
| 18          | S+90U10        | 1000                           | September                              | 10                              | September                                 | +90   |
| 19          | OU0            | 1000                           | October                                | 0                               | 0   | 0   |
| 20          | OU5            | 1000                           | October                                | 5                               | October                                   | 0   |
| 21          | OU10           | 1000                           | October                                | 10                              | October                                   | 0   |
| 22          | O+90U5         | 1000                           | October                                | 5                               | October                                   | +90   |
| 23          | O+90U10        | 1000                           | October                                | 10                              | October                                   | +90   |
| 24          | NU0            | 1000                           | November                               | 0                               | 0   | 0   |
| 25          | NU5            | 1000                           | November                               | 5                               | November                                  | 0   |
| 26          | NU10           | 1000                           | November                               | 10                              | November                                  | 0   |
| 27          | N+90U5         | 1000                           | November                               | 5                               | November                                  | +90   |
| 28          | N+90U10        | 1000                           | November                               | 10                              | November                                  | +90   |

<sup>a</sup> In each of the three columns related to date of application 0 indicates that there was no application.<sup>b</sup> DCD was applied within 24 h of urine application.

the fresh weight (kg<sup>-1</sup>) recorded and converted to hectares (Ansah et al., 2010). The responses of annual herbage production (DM was summed over all harvest dates within each year) and spring (February–April, inclusive) herbage production (DM was summed over spring harvest dates within each year) were computed for each plot for each year of the experiments.

#### 2.4.2. Herbage crude protein content and herbage N uptake

The dried herbage was subsequently milled through a 1 mm screen (Tecator Cyclotec 1093 Mill) for chemical analysis. Herbage crude protein (CP) content was determined using near infra-red spectroscopy (NIRS) analysis (NIRS, Model 6500, FOSS-NIR System, 3400 Hillerød, Denmark) and the equation developed by Burns et al. (2010). Herbage N uptake was computed using the following equations:

Herbage N content

$$= \frac{\text{herbage mass (kg DM ha}^{-1}) \times \text{herbage CP content (g kg}^{-1})}{6.25 \times 1000}$$

Herbage N uptake = treatment herbage N content

– control herbage N content

Spring herbage crude protein (averaged over spring harvest dates within each year) and spring herbage N uptakes (summed over spring harvest dates within each year) were computed for each treatment for the two years of the experiments.

#### 2.4.3. Soil ammonium and total oxidised nitrogen content

Soil sampling took place in year 2 on day 7, 14, 28, 56 and 84 post treatment application to determine NH<sub>4</sub><sup>+</sup>-N and total oxidised nitrogen (TON) content. Total oxidised nitrogen is NO<sub>3</sub><sup>-</sup> plus nitrite (NO<sub>2</sub><sup>-</sup>). Two soil cores (0–100 mm depth and 22.5 mm

diameter) were removed from each plot. Soil samples from each plot were composited and then sieved through a 2 mm screen to remove debris. Subsamples of 100 g were dried at 105 °C for 24 h to determine DM content. Another subsample of 30 g was extracted with 2 M KCl (Merck KGaA, Frankfurter StraBe 250, 64293 Darmstadt, Germany) solution by shaking for 2 h using the methodology described by Zaman et al. (1999). The extractant was filtered and frozen until analysis to determine NH<sub>4</sub><sup>+</sup>-N and TON content. The NH<sub>4</sub><sup>+</sup>-N and TON content in the extractant were measured using an Aquakem 600 Discrete Analyser (Thermo Scientific, Ratastie 2, P.O. Box 100, FI-01621 Vantaa, Finland). Soil bulk density (g m<sup>3</sup>) was calculated using methodology described by Hao et al. (2008). Soil NH<sub>4</sub><sup>+</sup>-N and TON content (kg N ha<sup>-1</sup>) were calculated by multiplying NH<sub>4</sub><sup>+</sup>-N (mg kg<sup>-1</sup>) or TON (mg kg<sup>-1</sup>) by the bulk density. Soil NH<sub>4</sub><sup>+</sup>-N and TON were recorded on each plot repeatedly on five sampling occasions in year 2 of the experiments only.

#### 2.4.4. Nitrogen balance

An N balance was calculated for the urine treatments with and without DCD, and the zero urine treatments, again with and without DCD. An N balance was also calculated for a hypothetical farm system in which 2.77% of the land area would receive urine patches treated with DCD; 2.77% represents the land area on which urine was deposited in a single grazing month. Nitrogen surplus was calculated by subtracting N outputs from N inputs.

#### 2.5. Nitrogen inputs

The N inputs consisted of fertilizer (0 or 350 kg N ha<sup>-1</sup>) and livestock urine deposition (0 or 208.3 kg N ha<sup>-1</sup>). Annual urine deposition per hectare was calculated using a stocking rate (SR) of 2.5 LU ha<sup>-1</sup>, urine patch coverage of 20.83% of pasture area per year, and urine deposition of 1000 kg N ha<sup>-1</sup> at each urination



**Table 3**The effect of DCD applied following urine application on herbage production (kg DM ha<sup>-1</sup>), crude protein (g kg<sup>-1</sup> DM) and nitrogen uptake (kg N ha<sup>-1</sup>) in Experiment 1.

| Measurement period                                       | Herbage production (kg DM ha <sup>-1</sup> ) |        | Crude protein (g kg ha <sup>-1</sup> ) | Nitrogen Uptake (kg ha <sup>-1</sup> ) | Surplus N (kg ha <sup>-1</sup> ) |
|--|--|--------|--|--|----------------------------------|
|  | Spring                                       | Annual | Spring                                 | Spring                                 | Annual                           |
| Year   |  |        |  |  |                                  |
| Year 1   | 3027   | 8067   | 246                                    | 115                                    | 22                               |
| Year 2   | 2617   | 8078   | 234                                    | 89                                     | 45                               |
| s.e.m.   | 60   | 137    | 1.9                                    | 2.6                                    | 16.1                             |
| Significance   | ***  | NS     | ***                                    | ***                                    | NS                               |
| Urine  |  |        |  |  |                                  |
| Zero urine   | 1688   | 6341   | 211                                    | 51                                     | -33                              |
| Urine  | 3200   | 8649   | 232                                    | 119                                    | 89                               |
| s.e.m.   | 97   | 222    | 2.6                                    | 3.9                                    | 8.2                              |
| Significance   | ***  | ***    | ***                                    | ***                                    | ***                              |
| DCD rate (kg ha <sup>-1</sup> )                          |  |        |  |  |                                  |
| 0  | 2714   | 7076   | 228                                    | 93                                     | 129                              |
| 5  | 2838   | 8200   | 243                                    | 104                                    | 28                               |
| 10   | 2914   | 8840   | 251                                    | 110                                    | 15                               |
| s.e.m.   | 79   | 184    | 2.2                                    | 3.2                                    | 19.5                             |
| Significance   | NS   | **     | ***                                    | **                                     | ***                              |
| Urine application date                                   |  |        |  |  |                                  |
| September  | 3175   | 7991   | 234                                    | 114                                    | 114                              |
| October  | 3125   | 8538   | 252                                    | 119                                    | 91                               |
| November   | 3300   | 9420   | 265                                    | 125                                    | 55                               |
| s.e.m.   | 98   | 225    | 2.7                                    | 3.9                                    | 16.6                             |
| Significance   | NS   | ***    | ***                                    | NS                                     | *                                |
| Urine application date × DCD rate (kg ha <sup>-1</sup> ) |  |        |  |  |                                  |
| September 0  | 3112   | 7397   | 223                                    | 106                                    | 137                              |
| September 5  | 2843   | 6943   | 235                                    | 104                                    | 116                              |
| September 10   | 3225   | 7796   | 239                                    | 113                                    | 101                              |
| October 0  | 3107   | 7997   | 238                                    | 114                                    | 139                              |
| October 5  | 3244   | 8858   | 253                                    | 122                                    | 95                               |
| October 10   | 3275   | 9960   | 271                                    | 127                                    | 63                               |
| November 0   | 3306   | 8578   | 241                                    | 121                                    | 111                              |
| November 5   | 3288   | 9811   | 266                                    | 133                                    | 55                               |
| November 10  | 3402   | 10503  | 286                                    | 135                                    | 28                               |
| s.e.m.   | 240  | 381    | 5.0                                    | 6.7                                    | 20.4                             |
| Significance   | NS   | *      | ***                                    | NS                                     | **                               |

NS, not significant.

\*  $P < 0.05$ .\*\*  $P < 0.01$ .\*\*\*  $P < 0.001$ .

production in Experiment 1 by +25%, compared to the OU0 treatment, and treatment NU5 and NU10 significantly increased annual herbage production compared to the NU0 treatment in Experiment 1 by +14 and +22%, respectively (Table 3).

### 3.2.2. Herbage crude protein content and nitrogen uptake

Spring herbage CP content was significantly ( $P < 0.001$ ) greater in Experiment 1 (+5%) (Table 3) and Experiment 2 (+5%) in Year 1 compared with Year 2 (Table 4).

Spring herbage CP content was significantly ( $P < 0.001$ ) greater following urine application in Experiment 1 (+10%) (Table 3) and in Experiment 2 (+18%) (Table 4).

There was a significant ( $P < 0.001$ ) effect of urine application date on spring herbage CP content in both Experiments (Tables 3 and 4). Applying urine in October and November significantly ( $P < 0.001$ ) increased spring herbage CP content compared to September application in Experiment 1 (Table 3). Urine application in September significantly ( $P < 0.01$ ) increased spring herbage CP content by +7 and +4%, respectively, compared to October application in Experiment 2 (Table 4).

Applying 5 or 10 kg DCD ha<sup>-1</sup> significantly ( $P < 0.001$ ) increased spring herbage CP content by +7 and +10%, respectively, compared to applying zero DCD in Experiment 1 (Table 3). There was no significant effect of DCD first application date or DCD second application date on spring herbage CP content in either Experiment 1 or 2.

There was a significant ( $P < 0.01$ ) interaction between urine application date and DCD rate on spring herbage CP content in Experiment 1 (Table 3). Treatment OU5 and OU10 had significantly ( $P < 0.05$ ) greater spring herbage CP content compared to the OU0 treatment, and treatment NU5 and NU10 had significantly ( $P < 0.05$ ) greater spring herbage CP content compared to the NU0 treatment (Table 3).

Herbage N uptake was significantly greater in Year 1 compared to Year 2 (Tables 3 and 4).

Herbage N uptake was significantly ( $P < 0.001$ ) greater on U treatments compared to OU treatments in Experiment 1 (+133%) and in Experiment 2 (+44%) (Tables 3 and 4). September urine application significantly ( $P < 0.01$ ) increased herbage N uptake compared to all other application dates in Experiment 2 (Table 4). There was a significant ( $P < 0.001$ ) effect of DCD rate on herbage N uptake in Experiment 1 but not in Experiment 2 (Tables 3 and 4). Applying

**Table 4**  
The effect of DCD applied following urine application on herbage production (kg DM ha<sup>-1</sup>), crude protein (g kg<sup>-1</sup> DM) and nitrogen uptake (kg N ha<sup>-1</sup>) in Experiment 2.

| Measurement period                                       | Herbage production (kg DM ha <sup>-1</sup> ) |        | Crude protein (g kg ha <sup>-1</sup> ) | Nitrogen Uptake (kg ha <sup>-1</sup> ) | Surplus N (kg ha <sup>-1</sup> ) |
|--|--|--------|--|--|----------------------------------|
|  | Spring                                       | Annual | Spring                                 | Spring                                 | Annual                           |
| Year   |  |        |  |  |                                  |
| Year 1   | 1685   | 5177   | 272                                    | 72                                     | 11                               |
| Year 2   | 1268   | 7360   | 258                                    | 47                                     | 13                               |
| s.e.m.   | 35   | 96     | 1.9                                    | 1.7                                    | 18.2                             |
| Significance   | ***  | ***    | ***                                    | ***                                    | NS                               |
| Urine  |  |        |  |  |                                  |
| Zero urine   | 1274   | 5548   | 234                                    | 45                                     | -51                              |
| Urine  | 1543   | 6508   | 275                                    | 65                                     | 67                               |
| s.e.m.   | 66   | 163    | 3.2                                    | 2.9                                    | 12                               |
| Significance   | ***  | ***    | ***                                    | ***                                    | ***                              |
| DCD rate (kg ha <sup>-1</sup> )                          |  |        |  |  |                                  |
| 0  | 1515   | 6388   | 265                                    | 60                                     | 62                               |
| 5  | 1457   | 6155   | 266                                    | 60                                     | 5                                |
| 10   | 1457   | 6262   | 264                                    | 59                                     | 2                                |
| s.e.m.   | 54   | 132    | 2.6                                    | 2.4                                    | 23.4                             |
| Significance   | NS   | NS     | NS                                     | NS                                     | NS                               |
| Urine application date                                   |  |        |  |  |                                  |
| September  | 1623   | 7013   | 284                                    | 73                                     | 30                               |
| October  | 1486   | 6014   | 265                                    | 63                                     | 115                              |
| November   | 1521   | 6497   | 276                                    | 58                                     | 55                               |
| s.e.m.   | 66   | 163    | 3.2                                    | 2.9                                    | 23.4                             |
| Significance   | NS   | ***    | ***                                    | **                                     | **                               |
| Urine application date × DCD rate (kg ha <sup>-1</sup> ) |  |        |  |  |                                  |
| September 0  | 1707   | 5951   | 279                                    | 76                                     | 51                               |
| September 5  | 1553   | 5258   | 265                                    | 72                                     | 25                               |
| September 10   | 1446   | 5273   | 270                                    | 72                                     | 25                               |
| October 0  | 1576   | 7060   | 288                                    | 72                                     | 105                              |
| October 5  | 1467   | 5942   | 267                                    | 63                                     | 114                              |
| October 10   | 1529   | 6340   | 276                                    | 61                                     | 121                              |
| November 0   | 1586   | 7029   | 286                                    | 72                                     | 93                               |
| November 5   | 1439   | 5844   | 262                                    | 60                                     | 66                               |
| November 10  | 1589   | 6878   | 282                                    | 62                                     | 26                               |
| s.e.m.   | 162  | 397    | 7.7                                    | 5.0                                    | 46.3                             |
| Significance   | NS   | NS     | *                                      | NS                                     | NS                               |

NS, not significant.

\*  $P < 0.05$ .

\*\*  $P < 0.01$ .

\*\*\*  $P < 0.001$ .

DCD at a rate of 5 or 10 kg ha<sup>-1</sup> significantly ( $P < 0.001$ ) increased herbage N uptake by 11 and 17 kg N ha<sup>-1</sup>, respectively, compared to applying zero DCD (Table 3). There was no significant effect of DCD first application date or DCD second application date on herbage N uptake in either Experiment 1 or 2.

There was no significant interaction between urine application date and DCD rate on herbage N uptake in either Experiment 1 or 2 (Tables 3 and 4).

### 3.2.3. Soil ammonium and total oxidised nitrogen content

Urine application had a significant effect ( $P < 0.001$ ) on soil NH<sub>4</sub><sup>+</sup>-N content compared to 0U treatments in Experiments 1 and 2 (Fig. 1A and B). Soil NH<sub>4</sub><sup>+</sup>-N content declined from day 7 to day 28 in Experiment 1 (Fig. 1A), and from day 7 to 84 in Experiment 2 (Fig. 1B) following urine application. There was no significant effect of DCD rate on NH<sub>4</sub><sup>+</sup>-N content in either Experiment 1 or 2 (data not shown).

Urine application had a significant effect ( $P < 0.001$ ) on soil TON content compared to 0U treatments in Experiments 1 and 2 (Fig. 1A and B). There was a significant ( $P < 0.001$ ) increase in soil TON content from sampling day 7 to 84 following urine application in


Experiment 1 (Fig. 1A) and to day 84 in Experiment 2 (Fig. 1B). There was a significant ( $P < 0.05$ ) effect of DCD rate on soil TON content in Experiment 1 (data not shown); treatment U5 had significantly lower soil TON content on sampling day 7 compared to the treatment U0.

### 3.2.4. Nitrogen balance

Surplus N was significantly ( $P < 0.001$ ) lower on treatments receiving zero urine compared to those receiving urine in Experiments 1 and 2 (Tables 3 and 4). September applied urine significantly ( $P < 0.05$ ) increased surplus N compared to November applied urine in Experiment 1 (Table 3). October applied urine significantly ( $P < 0.01$ ) increased surplus N compared to September applied urine in Experiment 2 (Table 4).

The application of 5 and 10 kg DCD ha<sup>-1</sup> significantly ( $P < 0.001$ ) reduced N surplus in Experiment 1 (Table 3); there was no effect in Experiment 2.

There was a significant ( $P < 0.01$ ) effect of the interaction between urine application date and DCD rate on N surplus in Experiment 1 only (Table 3). The N surplus on treatment OU10 was



**Fig. 1.** The effects of urine or zero urine applied at N equivalent rates of 1000 or 0 kg N ha<sup>-1</sup> on NH<sub>4</sub><sup>+</sup>-N and TON content in the upper layers (0–100 mm) of the soil profile in Experiment 1 (A) and in Experiment 2 (B).

significantly lower than treatment OU0, and was significantly lower on treatment NU10 compared to treatment NU0 (Table 3).

### 3.3. Experiments 3 and 4

#### 3.3.1. Herbage production

Spring herbage production was significantly ( $P < 0.001$ ) reduced in Year 2 compared to Year 1 in Experiments 3 and 4 (Tables 5 and 6). In Experiment 3 annual herbage production was significantly ( $P < 0.001$ ) reduced in Year 2 compared to Year 1 (Table 5), while in Experiment 4 annual herbage production was significantly ( $P < 0.001$ ) greater in Year 2 than in Year 1 (Table 6).

Urine application had no significant effect on spring herbage production in Experiment 3 (Table 5). Annual herbage production was significantly ( $P < 0.05$ ) increased (+5%) following urine application compared to zero urine application in Experiment 3 (Table 5). In Experiment 4, spring herbage production was significantly ( $P < 0.001$ ) reduced following urine application, but there was no significant effect on annual herbage production (Table 6).

In Experiment 3 September urine application significantly ( $P < 0.01$ ) increased spring herbage production compared to November urine application (Table 5).


There was no significant effect of DCD rate, DCD first application date or DCD second application date on spring or annual herbage production in either Experiment 3 or 4 (Tables 5 and 6).

#### 3.3.2. Herbage crude protein content and nitrogen uptake

Spring herbage CP content was significantly ( $P < 0.001$ ) greater in Year 1 (+5%) compared to Year 2 in Experiment 4 (Table 6).

Urine application significantly ( $P < 0.001$ ) increased spring herbage CP content in Experiments 3 and 4 (Tables 5 and 6) compared to the OU treatment. October and November applied urine significantly ( $P < 0.01$ ) increased spring herbage CP content compared to September applied urine in Experiment 3 (Table 5).

Applying DCD at a rate of 5 or 10 kg ha<sup>-1</sup> significantly ( $P < 0.001$ ) increased spring herbage CP content compared to applying zero



**Fig. 2.** The effects of urine or zero urine applied at N equivalent rates of 1000 or 0 kg N ha<sup>-1</sup> on NH<sub>4</sub><sup>+</sup>-N and TON content in the upper layers (0–100 mm) of the soil profile in Experiment 3 (A) and in Experiment 4 (B).

DCD in Experiments 3 and 4 (Tables 5 and 6). There was no significant effect of DCD first application date or DCD second application date on spring herbage CP content in either Experiment 3 or 4.

There was a significant interaction ( $P < 0.001$ ) between urine application date and DCD rate on spring herbage CP content in Experiment 3 (Table 5); treatment NU0 had a significantly ( $P < 0.01$ ) lower spring herbage CP content than NU10.

Herbage N uptake was significantly ( $P < 0.001$ ) reduced in Year 2 (–55%) compared to Year 1 in Experiment 3 (Table 5). In Experiment 4, herbage N uptake was significantly ( $P < 0.001$ ) greater in Year 2 compared to Year 1 (Table 6).

Urine application significantly ( $P < 0.001$ ) increased herbage N uptake in both Experiments 3 and 4 compared to OU treatments (Tables 5 and 6). There was no significant effect of urine application date on herbage N uptake in either Experiment 3 or 4.

Applying DCD at a rate of 5 or 10 kg ha<sup>-1</sup> significantly ( $P < 0.001$ ) increased herbage N uptake by +19 and +22%, respectively, compared to applying zero DCD in Experiment 4 (Table 6). There was no significant effect of DCD first application date or DCD second application date on herbage N uptake in either Experiment 3 or 4.

Treatments SU5 and SU10 significantly ( $P < 0.01$ ) increased herbage N uptake in Experiment 4 by +27 and 29%, respectively, compared to SU0 treatment (Table 6).

#### 3.3.3. Soil ammonium and total oxidised nitrogen content

Urine application significantly ( $P < 0.001$ ) increased soil NH<sub>4</sub><sup>+</sup>-N content compared to the OU treatments up to day 14 in Experiment 3 and up to day 56 in Experiment 4 (Fig. 2A and B). The NH<sub>4</sub><sup>+</sup>-N content declined from day 7 to day 84 post urine application. There was no significant effect of DCD rate on soil NH<sub>4</sub><sup>+</sup>-N content in both Experiments 3 and 4 (data not shown).

Urine application significantly ( $P < 0.001$ ) increased soil TON content compared to OU treatments across all sampling days in Experiments 3 and 4 (Fig. 2A and B). There was no significant effect

**Table 5**  
The effect of DCD applied following urine application on herbage production (kg DM ha<sup>-1</sup>), crude protein (g kg<sup>-1</sup> DM) and nitrogen uptake (kg N ha<sup>-1</sup>) in Experiment 3.

| Measurement period                                       | Herbage production (kg DM ha <sup>-1</sup> ) |        | Crude protein (g kg ha <sup>-1</sup> ) | Nitrogen Uptake (kg ha <sup>-1</sup> ) | Surplus N (kg ha <sup>-1</sup> ) |
|--|--|--------|--|--|----------------------------------|
|  | Spring                                       | Annual | Spring                                 | Spring                                 | Annual                           |
| Year   |  |        |  |  |                                  |
| Year 1   | 4437   | 13467  | 293                                    | 204                                    | 396                              |
| Year 2   | 2122   | 9682   | 290                                    | 92                                     | 438                              |
| s.e.m.   | 59   | 138    | 1.7                                    | 2.5                                    | 22.6                             |
| Significance   | ***  | ***    | NS                                     | ***                                    | NS                               |
| Urine  |  |        |  |  |                                  |
| Zero urine   | 3142   | 11170  | 279                                    | 134                                    | 307                              |
| Urine  | 3325   | 11710  | 296                                    | 153                                    | 496                              |
| s.e.m.   | 103  | 241    | 2.8                                    | 4.2                                    | 5.9                              |
| Significance   | NS   | *      | ***                                    | ***                                    | ***                              |
| DCD rate (kg ha <sup>-1</sup> )                          |  |        |  |  |                                  |
| 0  | 3345   | 11588  | 279                                    | 145                                    | 498                              |
| 5  | 3256   | 11637  | 296                                    | 150                                    | 413                              |
| 10   | 3238   | 11499  | 300                                    | 149                                    | 401                              |
| s.e.m.   | 84   | 199    | 2.3                                    | 3.4                                    | 30                               |
| Significance   | NS   | NS     | ***                                    | NS                                     | NS                               |
| Urine application date                                   |  |        |  |  | 0.08                             |
| September  | 3565   | 11583  | 289                                    | 156                                    | 499                              |
| October  | 3301   | 11648  | 297                                    | 152                                    | 494                              |
| November   | 3108   | 11898  | 302                                    | 150                                    | 493                              |
| s.e.m.   | 103  | 243    | 2.8                                    | 4.2                                    | 12.5                             |
| Significance   | **   | NS     | **                                     | NS                                     | NS                               |
| Urine application date × DCD rate (kg ha <sup>-1</sup> ) |  |        |  |  |                                  |
| September 0  | 3781   | 11940  | 279                                    | 157                                    | 494                              |
| September 5  | 3123   | 10841  | 278                                    | 152                                    | 504                              |
| September 10   | 3440   | 12480  | 288                                    | 159                                    | 498                              |
| October 0  | 3350   | 11239  | 295                                    | 137                                    | 524                              |
| October 5  | 3478   | 12060  | 302                                    | 160                                    | 488                              |
| October 10   | 3008   | 11937  | 302                                    | 159                                    | 485                              |
| November 0   | 3565   | 11569  | 292                                    | 156                                    | 477                              |
| November 5   | 3304   | 12043  | 311                                    | 150                                    | 497                              |
| November 10  | 2877   | 11277  | 317                                    | 144                                    | 496                              |
| s.e.m.   | 245  | 671    | 6.8                                    | 7.2                                    | 24.9                             |
| Significance   | NS   | NS     | *                                      | NS                                     | NS                               |

NS, not significant.

\*  $P < 0.05$ .

\*\*  $P < 0.01$ .

\*\*\*  $P < 0.001$ .

of DCD rate on soil TON content in Experiments 3 and 4 (data not shown).

### 3.3.4. Nitrogen balance

Urine application significantly ( $P < 0.001$ ) increased surplus N compared to zero urine application by 62 and 59% in Experiments 3 and 4, respectively (Tables 5 and 6).

The application of DCD at a rate of 5 and 10 kg ha<sup>-1</sup> significantly ( $P < 0.05$ ) reduced N surplus by 20 and 21%, respectively, in Experiment 4 (Table 6).

## 4. Discussion

### 4.1. Herbage production, herbage crude protein content and herbage N uptake

Whitehead and Bristow (1990) reported that the recovery of urinary N by the soil/plant component of the sward is subject to a number of factors including rainfall amount, which affects plant growth and leaching, and variation in temperature, which contributes to evaporation and to plant growth. In Experiment 1 (zero

N fertiliser) average spring and annual herbage production was increased at both sites when urine was applied, similar to Qiu et al. (2010), due to high N content in urine (Haynes and Williams, 1993). In Experiment 3 (350 kg N ha<sup>-1</sup> year<sup>-1</sup>) there was a positive effect of urine application on annual herbage production. While some of the additional N applied in the urine would have been used for herbage growth, it is likely that most of the N was surplus to the requirements of the sward. Frame (1992) and Van Burg et al. (1981) suggest that N supplies greater than 350–400 kg N ha<sup>-1</sup> year<sup>-1</sup> are surplus to the requirements of the growing plant. Lantinga et al. (1987) reported that grass is more susceptible to urine scorch at N fertilisation levels similar to the rate applied in Experiments 3 and 4. Middelkoop and Deenan (1990) observed that urine had a negative effect on herbage production in the treated areas due to scorching; this effect was strongest at 400 kg N ha<sup>-1</sup> and increased with increasing N concentration. The application of urine to N fertiliser treated swards may have inhibited herbage production as a result of some scorching in Experiments 3 and 4.

The application of DCD at 5 or 10 kg ha<sup>-1</sup> significantly increased annual herbage production in Experiment 1; similar to Di and Cameron (2002, 2005). In a whole paddock scenario receiving zero



**Table 6**The effect of DCD applied following urine application on herbage production (kg DM ha<sup>-1</sup>), crude protein (g kg<sup>-1</sup> DM) and nitrogen uptake (kg N ha<sup>-1</sup>) in Experiment 4.

| Measurement period                                       | Herbage production (kg DM ha <sup>-1</sup> ) |        | Crude protein (g kg ha <sup>-1</sup> ) | Nitrogen Uptake (kg ha <sup>-1</sup> ) | Surplus N (kg ha <sup>-1</sup> ) |
|--|--|--------|--|--|----------------------------------|
|  | Spring                                       | Annual | Spring                                 | Spring                                 | Annual                           |
| Year   |  |        |  |  |                                  |
| Year 1   | 2038   | 7254   | 299                                    | 21                                     | 429                              |
| Year 2   | 1146   | 8507   | 285                                    | 52                                     | 440                              |
| s.e.m.   | 35   | 117    | 1.4                                    | 1.6                                    | 24.0                             |
| Significance   | ***  | ***    | ***                                    | ***                                    | NS                               |
| Urine  |  |        |  |  |                                  |
| Zero urine   | 1800   | 8083   | 279                                    | 25                                     | 327                              |
| Urine  | 1592   | 7880   | 296                                    | 40                                     | 521                              |
| s.e.m.   | 60   | 167    | 1.9                                    | 1.7                                    | 7.4                              |
| Significance   | ***  | NS     | ***                                    | ***                                    | ***                              |
| DCD rate (kg ha <sup>-1</sup> )                          |  |        |  |  |                                  |
| 0  | 1544   | 7776   | 288                                    | 32                                     | 530                              |
| 5  | 1560   | 7894   | 294                                    | 38                                     | 426                              |
| 10   | 1672   | 7971   | 294                                    | 39                                     | 420                              |
| s.e.m.   | 50   | 147    | 1.5                                    | 1.6                                    | 31.2                             |
| Significance   | NS   | NS     | *                                      | **                                     | *                                |
| Urine application date                                   |  |        |  |  |                                  |
| September  | 1470   | 7643   | 294                                    | 41                                     | 520                              |
| October  | 1459   | 7678   | 296                                    | 40                                     | 525                              |
| November   | 1639   | 8119   | 299                                    | 39                                     | 517                              |
| s.e.m.   | 61   | 176    | 1.9                                    | 1.9                                    | 7                                |
| Significance   | NS   | NS     | NS                                     | NS                                     | NS                               |
| Urine application date × DCD rate (kg ha <sup>-1</sup> ) |  |        |  |  |                                  |
| September 0  | 1489   | 7271   | 289                                    | 34                                     | 536                              |
| September 5  | 1337   | 7435   | 292                                    | 43                                     | 517                              |
| September 10   | 1630   | 8134   | 298                                    | 44                                     | 516                              |
| October 0  | 1342   | 7615   | 293                                    | 37                                     | 538                              |
| October 5  | 1504   | 7804   | 299                                    | 44                                     | 519                              |
| October 10   | 1512   | 7092   | 300                                    | 40                                     | 525                              |
| November 0   | 1578   | 8043   | 299                                    | 37                                     | 517                              |
| November 5   | 1537   | 7794   | 296                                    | 38                                     | 520                              |
| November 10  | 1774   | 8131   | 300                                    | 43                                     | 514                              |
| s.e.m.   | 151  | 410    | 4.7                                    | 3.0                                    | 13.4                             |
| Significance   | NS   | NS     | *                                      | NS                                     | NS                               |

NS, not significant.

\*  $P < 0.05$ .\*\*  $P < 0.01$ .\*\*\*  $P < 0.001$ .

N fertiliser, growing 8000 kg DM ha<sup>-1</sup> and grazed by dairy cows the addition of DCD at either rate would increase annual herbage production by between 200 and 400 kg DM ha<sup>-1</sup>, assuming that 25% of the paddock receives urine deposition per year plus 5 or 10 kg DCD ha<sup>-1</sup> (Haynes and Williams, 1993). The application of DCD did not increase herbage production in Experiment 2, 3 or 4, similar to Cookson and Cornforth (2002), Menneer et al. (2008) and Monaghan et al. (2009), and therefore the hypothesis that the application of DCD at different rates and times in autumn and winter would increase herbage production in Experiment 2, 3 and 4 is rejected. Previous authors report that DCD should be applied in cool temperatures and periods of low rainfall to maximise its effectiveness (Vallejo et al., 2005). The rainfall experienced following DCD application in this experiment may have leached the DCD beyond the rooting zone, therefore reducing its effectiveness. High soil and air temperatures recorded in October and November may have reduced the half life of the DCD, therefore reducing its capacity to slow the conversion of NH<sub>4</sub><sup>+</sup> to TON. As mentioned temperature is one of the most influential environmental factors affecting the effectiveness of DCD (Williamson et al., 1996). An increase in temperature can have a negative effect on the persistence of DCD in

the soil, reducing the time frame in which it can provide effective nitrification reduction.

Developing a DCD application strategy to contribute to increased herbage production and help to mitigate environmental emissions associated with increased agricultural production in Ireland would most likely require consideration of factors other than soil temperature. It may be worth considering DCD application at a time when rainfall is lower than in autumn/winter so that the DCD is less likely to be leached beyond the rooting zone. In such instances the DCD will reduce nitrification in a soil zone in which herbage could benefit from the availability of additional N for plant growth. The soil sampling undertaken in Experiments 3 and 4 indicates that DCD had almost no effect on soil mineral N content in the 0–100 mm soil horizon. However, as DCD has been shown to be effective at reducing N<sub>2</sub>O emissions and NO<sub>3</sub><sup>-</sup> leaching in Irish soils (Selbie et al., 2011; Richards et al., 2008) it is quite possible that the DCD is inhibiting nitrification beyond a soil depth of 100 mm.

During this study urine deposition increased spring herbage CP content in all experiments similar to other authors (Ledgard et al., 1982; Williams and Haynes, 1994). Similar to Moir et al. (2007), the application of DCD had an effect on spring herbage CP content in

Experiments 1 and 3 at MPK as a result of increased N uptake. The application of DCD may have provided additional N in the rooting zone for plant uptake; but due to low temperatures in spring this did not result in increased grass growth but rather resulted in the accumulation of N in the plant foliage.

Throughout this study urine application increased herbage N uptake compared to zero urine application, similar to Cuttle and Bourne (1993). The increase in herbage N uptake may be due to the uptake of urea into the plant by roots; this mode of N uptake is small compared to the uptake of  $\text{NO}_3^-$ -N and  $\text{NH}_4^+$ -N. However, foliar adsorption of urea can occur at a high rate and can contribute substantially to plant N accumulation and not result in herbage production (Camberato, 2001).

Also, N uptake is reported to increase with increasing rates of N applied, including rates of N above which herbage production is maximised (Lkhagvasuren, 2007). Nitrogen uptake is also reported to occur in periods of low grass growth (Blombäck and Eckersten, 1997), also resulting in increased CP content but not herbage production. The ability of perennial ryegrasses to increase N uptake below the top 100 mm of the soil profile are limited as a small proportion of the roots occur below the top 100 mm of the soil profile (Garwood, 1967) and also 90% of the root activity of perennial ryegrass occurs in the top 125 mm of the soil profile (Syers et al., 1984), thereby, the capability of perennial ryegrasses to utilise the available N for herbage production below the top 100 mm of the soil profile is greatly reduced.

#### 4.2. Soil ammonium and total oxidised nitrogen content

Urine application increased soil  $\text{NH}_4^+$ -N and TON content compared to zero urine application. In urine treated soils, urea is rapidly hydrolysed to  $\text{NH}_4^+$ -N and reaches maximum values soon after urine application (Zaman et al., 2009). Nitrification then oxidises the  $\text{NH}_4^+$ -N to nitrite and the nitrite is further oxidised to  $\text{NO}_3^-$  (Estavillo et al., 2002). In the experiments reported here, soil  $\text{NH}_4^+$ -N content declined with time following urine application. The period following urine application was one of low sward N demand due to low rates of grass growth and as a result N was likely to be lost by leaching through the soil profile or through immobilization (Woodmaness et al., 1981). At MPK the elevated  $\text{NH}_4^+$ -N content following urine application declined and was similar to the zero urine treatments by day 84 and so no additional N was available for spring grass growth. At BD, there was considerably more soil  $\text{NH}_4^+$ -N present in the rooting zone (0–100 mm) at sampling day 84 on the urine treatments. The rapid movement of N to below the rooting zone at MPK can be attributed to the free draining soil characteristics at this site, while at BD the drainage capabilities are more restricted, resulting in reduced leaching potential (Decau et al., 2003; Richards et al., 2008).

Moir et al. (2007) reported that the application of 10 kg DCD  $\text{ha}^{-1}$  to urine patches inhibited urine sourced  $\text{NH}_4^+$ -N in the soil for several weeks over the autumn/winter period, however, in Experiments 1 and 2 reported here DCD provided inhibition of urine sourced  $\text{NH}_4^+$ -N for a short period of 14 days. The soil sampling undertaken in the four experiments indicate that DCD had almost no effect on soil mineral N content to a soil depth of 100 mm suggesting that the DCD may have leached below this soil depth.

#### 4.3. Nitrogen balance

Research has shown that N losses from a growing sward increase exponentially with an increase in N inputs (Scholefield et al., 1993), this was observed in the experiments reported here when urine and fertiliser were applied to a growing sward. As it is not possible to quantify the total quantity of surplus N in this study as the only N output measured was herbage N uptake, the N balances reported

are a crude estimate of N surplus. As DCD can reduce  $\text{NO}_3^-$  leaching and  $\text{N}_2\text{O}$  emissions, as measured by other authors, it will have a positive effect on a farm N balance. However, DCD is most effective on urine patches and in cool temperatures, and so the overall effect in reducing N surplus will be proportional to the area of the farm affected by urine patches, particularly in autumn. For example, using an annual farm-gate N balance scenario where DCD was applied in September following urine deposition on a farm applying 250 kg inorganic N  $\text{ha}^{-1}$  year<sup>-1</sup> compared to a farm applying zero DCD, the N surplus would be reduced by 3 kg N based on the effect of DCD on herbage production.

## 5. Conclusions

Urine increased spring and annual herbage production when applied to swards receiving zero fertiliser; however when urine was applied to swards receiving fertiliser, spring herbage production was not significantly increased. Urine application also increased N uptake and spring herbage CP content. DCD applied at 5 and 10 kg  $\text{ha}^{-1}$  increased annual herbage production at MPK by 200 to 400 kg DM  $\text{ha}^{-1}$ . Dicyandiamide increased spring herbage CP content at MPK in Experiments 1 and 3. The addition of DCD also increased herbage N uptake at MPK in Experiment 1 and at BD in Experiment 4. In Experiments 1 and 3, soil  $\text{NH}_4^+$ -N content on the urine treatments declined over the sampling period, while soil TON content was significantly increased on the urine treatments compared to the zero urine treatments in both experiments. Dicyandiamide applied to October and November deposited urine reduced N surplus in Experiments 1 and 4. Overall the effects of DCD on herbage production, surplus N and other parameters in this study were not consistent.

## Acknowledgements

The authors wish to thank Messrs. M. Feeney, J. McCarthy and F. Flynn for their technical assistance. Gratitude is also expressed to the Moorepark farm staff for their assistance with measurements taken during this study. The first author (P.J. O'Connor) was in receipt of a Teagasc Walsh Fellowship.

## References

- Amberger, A., 1989. Research on dicyandiamide as a nitrification inhibitor and future outlook. *Commun. Soil Sci. Plant Anal.* 20, 1933–1955.
- Ansah, T., Osafo, E.L.K., Hansen, H.H., 2010. Herbage yield and chemical composition of four varieties of Napier (*Pennisetum purpureum*) grass harvested at three different days after planting. *Agric. Biol. J. N. Am.* 1, 923–929.
- Blombäck, K., Eckersten, H., 1997. Simulated growth and nitrogen dynamics of a perennial rye grass. *Agric. Forest Meteorol.* 88, 37–45.
- Burns, B.A., Gilliland, T.J., McGilloway, D.A., O'Donovan, M., Lewis, E., Blount, N., O'Kiely, P., 2010. Using NIRS to predict composition characteristics of *Lolium perenne* L. cultivars. In: *Advances in Animal Biosciences. Food, Feed, Energy and Fibre from the Land—a Vision for 2020*. British Soc. Anim. Sci., Belfast, UK, 321 pp.
- Bussink, D.W., Oenema, O., 1996. Ammonia volatilization from dairy farming systems in temperate areas: a review. In: *Scandinavian-Association-of-Agricultural-Scientists Seminar on Ammonia Emissions from Agriculture*, Uppsala, Sweden, pp. 19–33.
- Camberato, J.J., 2001. Nitrogen in Soil and Fertilizers, vol. 8. SC Turf-grass Foundation News, pp. 6–10.
- Cookson, W.R., Cornforth, I.S., 2002. Dicyandiamide slows nitrification in dairy cattle urine patches: effects on soil solution composition, soil pH and pasture yield. *Soil Biol. Biochem.* 34, 1461–1465.
- Cuttle, S.P., Bourne, P.C., 1993. Uptake and leaching of nitrogen from artificial urine applied to grassland on different dates during the growing season. *Plant Soil* 150, 77–86.
- Decau, M.L., Simon, J.C., Jacquet, A., 2003. Fate of urine nitrogen in three soils throughout a grazing season. *J. Environ. Qual.* 32, 1405–1413.
- Dennis, S.J., Richards, K., Cameron, K.C., Di, H.J., Moir, J.L., Fay, D., Staples, V., Sills, P., 2008. Dicyandiamide (DCD) nitrification inhibitor reduces nitrous oxide emissions from soils. In: *Proceedings of Agricultural Research Forum*. The Tullamore Court Hotel, Tullamore, Co., Offaly, 6 pp.

- Di, H.J., Cameron, K.C., 2002. The use of a nitrification inhibitor, dicyandiamide (DCD), to decrease nitrate leaching and nitrous oxide emissions in simulated grazed and irrigated grassland. *Soil Use Manage.* 18, 395–403.
- Di, H.J., Cameron, K.C., 2004. Effects of temperature and application rate of a nitrification inhibitor, dicyandiamide (DCD), on nitrification rate and microbial biomass in a grazed pasture soil. *Aust. J. Soil Res.* 42, 927–932.
- Di, H.J., Cameron, K.C., 2005. Reducing environmental impacts of agriculture by using a fine particle suspension nitrification inhibitor to decrease nitrate leaching from grazed pastures. *Agric. Ecosyst. Environ.* 109, 202–212.
- Dillon, P., Roche, J.R., Shalloo, L., Horan, B., 2005. Optimising financial return from grazing in temperate pastures. In: *Utilisation of grazed grass in temperate animal systems, Proceedings of a Satellite Workshop of the XXth International Grassland Congress*. Wageningen Academic Publishers, Cork, Ireland, pp. 131–147.
- Estavillo, J.M., Merino, P., Pinto, M., Yamulki, S., Gebauer, G., Sapek, A., Corré, W., 2002. Short term effect of ploughing a permanent pasture on N<sub>2</sub>O production from nitrification and denitrification. *Plant Soil* 239, 253–265.
- Frame, J., 1992. *Improved Grassland Management*. Farming Press Books, Ipswich, United Kingdom.
- Garwood, E.A., 1967. Some effects of soil water conditions and soil temperature on the roots of grasses. *Grass Forage Sci.* 22, 176–181.
- Hao, X., Ball, B.C., Culley, J.L.B., Carter, M.R., Parkin, G.W., 2008. Soil density and porosity. In: *Carter, M.R., Gregorich, E.G. (Eds.), Soil Sampling and Methods of Analysis*. Taylor and Francis Group, Boca Raton, FL, USA, pp. 743–759.
- Haynes, R.J., Williams, P.H., 1993. Nutrient cycling and soil fertility in the grazed pasture ecosystem. *Adv. Agron.* 49, 119–199.
- Hennessy, D., O'Donovan, M., French, P., Laidlaw, A.S., 2008. Manipulation of herbage production by altering the pattern of applying nitrogen fertilizer. *Grass Forage Sci.* 63, 152–166.
- Kelliher, F.M., Clough, T.J., Clark, H., Rys, G., Sedcole, J.R., 2008. The temperature dependence of dicyandiamide (DCD) degradation in soils: a data synthesis. *Soil Biol. Biochem.* 40, 1878–1882.
- Kennedy, E., O'Donovan, M., Murphy, J.P., Delaby, L., O'Mara, F., 2005. Effects of grass pasture and concentrate-based feeding systems for spring-calving dairy cows in early spring on performance during lactation. *Grass Forage Sci.* 60, 310–318.
- Lantinga, E.A., Keuning, J.A., Groenwold, J., Deenen, P.J.A.G., 1987. Distribution of excreted nitrogen by grazing cattle and its effects on sward quality, herbage production and utilization. In: *Animal Manure on Grassland and Fodder Crops. Fertilizer or Waste? Proceedings of an International Symposium of the European Grassland Federation* Wageningen. Martinus Nijhoff Publishers, Dordrecht, The Netherlands.
- Ledgard, S.F., Steele, K.W., Saunders, W.H.M., 1982. Effects of cow urine and its major constituents on pasture properties. *N. Z. J. Agric. Res.* 25, 61–68.
- Ledgard, S.F., Penno, J.W., Sprosen, M.S., 1999. Nitrogen inputs and losses from clover/grass pastures grazed by dairy cows, as affected by nitrogen fertilizer application. *J. Agric. Sci.* 132, 215–225.
- Lkhagvasuren, B., 2007. *Plant and Soil Responses to Fertilization of Grasslands in Saskatchewan, Canada and Selenge, Mongolia*. Department of Soil Science, University of Saskatchewan, Saskatoon, pp. 1–100.
- Menneer, J.C., Ledgard, S.F., Sprosen, M., 2008. Soil N process inhibitors alter nitrogen leaching dynamics in a pumice soil. *Aust. J. Soil Res.* 46, 323–331.
- Middelkoop, N., Deenan, P.J.A.G., 1990. The local influence of cattle dung and urine and its interactions with fertilizer nitrogen on herbage dry matter production. In: *Soil–Grassland–Animals Relationships, vol. 2, Proceedings of the 13th General meeting of the European Grassland Federation, Banska Bystrica, Czechoslovakia*, pp. 67–70.
- Moir, J.L., Cameron, K.C., Di, H.J., 2007. Effects of the nitrification inhibitor dicyandiamide on soil mineral N, pasture yield, nutrient uptake and pasture quality in a grazed pasture system. *Soil Use Manage.* 23, 111–120.
- Monaghan, R.M., Smith, L.C., Ledgard, S.F., 2009. The effectiveness of a granular formulation of dicyandiamide (DCD) in limiting nitrate leaching from a grazed dairy pasture. *N. Z. J. Agric. Res.* 52, 145–159.
- O'Connell, K., Humphreys, J., Watson, C.J., 2004. Quantification of nitrogen sources for grassland. In: *Winter Scientific Meeting, Fertiliser Association of Ireland, Dublin*, pp. 15–28.
- O'Donovan, M., Kennedy, E., 2007. Maximising dairy cow performance from grazed grass. In: *Buckley, F. (Ed.), Moorepark 07 Irish Dairying—Winning on a World Stage*. Teagasc, Dairy Production Research Centre, Moorepark, Fermoy, Co., Cork, Ireland, pp. 29–35.
- Qiu, W., Di, H.J., Cameron, K.C., 2010. Nitrous oxide emissions from animal urine as affected by season and a nitrification inhibitor dicyandiamide. *J. Soils Sediment.* 10, 1229–1235.
- Richards, K., Dennis, S.J., Cameron, K.C., Di, H.J., Moir, J.L., Fay, D., Stark, C.H., Staples, V., Sills, P., 2008. The potential role of the nitrification inhibitor DCD for reducing nitrate leaching from grazed grassland. In: *Proceedings of Agricultural Research Forum, Tullamore, Co., Offaly*, 1 pp.
- Ryan, M., Brophy, C., Connolly, J., McNamara, K., Carton, O.T., 2006. Monitoring of nitrogen leaching on a dairy farm during four drainage seasons. *Ir. J. Agric. Food Res.* 45, 115–134.
- SAS, 2003. *Statistical Analysis System User's Guide, Version 9.1.3*. SAS Institute Inc., Cary, NC.
- Scholefield, D., Tyson, K.C., Garwood, E.A., Armstrong, A.C., Hawkins, J., Stone, A.C., 1993. Nitrate leaching from grazed grassland lysimeters: effects of fertilizer input, field drainage, age of sward and patterns of weather. *J. Soil Sci.* 44, 601–613.
- Selbie, D.R., Lanigan, G., Di, H.J., Moir, J.L., Cameron, K.C., Richards, K.G., 2011. Improving nitrogen efficiency using a nitrification inhibitor on urine-affected soil—a grassland lysimeter study. In: *Proceedings of Agricultural Research Forum, Tullamore, Co., Offaly*, 2 pp.
- Serna, M., Legaz, F., Primo-Millo, E., 1995. Improvement of the N fertilizer efficiency with dicyandiamide (DCD) in citrus trees. *Nutr. Cycl. Agroecosyst.* 43, 137–142.
- Shalloo, L., Dillon, P., O'Loughlin, J., Rath, M., Wallace, M., 2004. Comparison of a pasture-based system of milk production on a high rainfall, heavy-clay soil with that on a lower rainfall, free-draining soil. *Grass Forage Sci.* 59, 157–168.
- Syers, J.K., Ryden, J.C., Garwood, E.A., 1984. Assessment of root activity of perennial ryegrass and white clover measured using <sup>32</sup>phosphorus as influenced by method of isotope placement, irrigation and method of defoliation. *J. Sci. Food Agric.* 35, 959–969.
- Vallejo, A., Garcia-Torres, L., Diez, J.A., Arce, A., Lopez-Fernandez, S., 2005. Comparison of N losses (NO<sub>3</sub><sup>-</sup>, N<sub>2</sub>O, NO) from surface applied, injected or amended (DCD) pig slurry of an irrigated soil in a Mediterranean climate. *Plant Soil* 272, 313–325.
- Van Burg, P.F.J., Prins, W.H., den Boer, D.J., Sluiman, W.J., 1981. Nitrogen and intensification of livestock farming in EEC countries. In: *Proceedings of Fertiliser Society*, vol. 199, London, pp. 1–78.
- Whitehead, D.C., 1995. Grassland nitrogen. In: *O'Connell, K., Humphreys, J., Watson, C.J. (Eds.), Quantification of Nitrogen Sources for Grassland*, vol. 40, Proceedings of Fertiliser Association, Ireland, pp. 15–29.
- Whitehead, D.C., Bristow, A.W., 1990. Transformation of nitrogen following the application of 15N – labelled cattle urine to an established sward. *J. Appl. Ecol.* 27, 667–678.
- Williams, P.H., Haynes, R.J., 1994. Comparison of initial wetting pattern, nutrient concentrations in soil solution and the fate of 15N-labelled urine in sheep and cattle urine patch areas of pasture soil. *Plant Soil* 162, 49–59.
- Williamson, J.C., Menneer, J.C., Torrens, R.S., 1996. Impact of dicyandiamide on the internal nitrogen cycle of a volcanic, silt loam soil receiving effluent. *Appl. Soil Ecol.* 4, 39–48.
- Woodmaness, R.G., Vallis, I., Mott, J.J., 1981. Grassland nitrogen. In: *Clark, F.E., Ross-wall, T. (Eds.), Terrestrial Nitrogen Cycles, Processes, Ecosystem Strategies and Management Impacts*, Ecological bulletins No. 33.
- Zaman, M., Blennerhassett, J.D., 2010. Effects of the different rates of urease and nitrification inhibitors on gaseous emissions of ammonia and nitrous oxide, nitrate leaching and pasture production from urine patches in an intensive grazed pasture system. *Agric. Ecosyst. Environ.* 136, 236–246.
- Zaman, M., Di, H.J., Cameron, K.C., Framptom, C.M., 1999. Gross nitrogen mineralization and nitrification rates and their relationships to enzyme activities and the soil microbial biomass in soils treated with dairy shed effluent and ammonium fertilizer at different water potentials. *Biol. Fertil. Soils* 29, 178–186.
- Zaman, M., Saggat, S., Blennerhassett, J.D., Singh, J., 2009. Effect of urease and nitrification inhibitors on N transformation, gaseous emissions of ammonia and nitrous oxide, pasture yield and N uptake in grazed pasture system. *Soil Biol. Biochem.* 41, 1270–1280.