



Phytochemical and antibacterial potentials of *Tecoma stans* and *Costus afer*

Ekpe Ini P^{1*} and Udosen EO²

Abstract

Phytochemical composition and antibacterial potential of ethanolic extract of leaves/roots of *Tecoma stans* and leaves of *Costus afer* evaluated were collected at Malabo Campus, University of Calabar, Calabar and from Eman-Uruan local government area, Akwa Ibom, Nigeria respectively. Fresh leaves of both plants and roots of *Tecoma stan C* in the refrigerator.

The result showed that the leaves/roots of *T. stans* and leaves of *C. afer* contain Alkaloids, Flavonoids, Saponins and Glycosides. The MIC of leaves of *T. stans* showed 5.21mg/ml for *S. aureus*, 10.4mg/ml for *E.coli*, 83.3mg/ml for *Proteus speies*. The roots of *T. stans* showed 75mg/ml for *S.aureus*. The MIC for *C. afer* showed 5.2mg/ml for *S. aureus* and 312.5mg/ml for *M. organii*. The results indicate that both plants can be used to achieve significant inhibitory effects as antimicrobial agents for treatment of bacterial infections in the absence of orthodox medication.

Keywords: Antibacterial; natural products; nutrient broth; incubation; inhibition

Introduction

Natural products are commonly understood to refer to herbs, herbal concoctions, and dietary supplements, alternative medicine be it Chinese, African, Philippines, Indian origin [1]. They are of prebiotic origin or they may originate from plants, animal or microbial sources

The cost of drug discovery and drug development continues to increase at astronomical rates and despite the successes the interest in natural products has increased and waned in popularity with various pharmaceutical companies. Natural products have contributed immensely to the development of the pharmaceutical industry and research, and have been widely applied in human medicine with significant promise to target resurgent and emerging diseases [2]. Some pharmaceutical companies have however shown considerable interest in drugs derived from natural sources as they are “greener sources”, safe and more dependable compared to synthetic drugs. The research into the use of medicinal plants derived from natural sources alone in the field of medicine covers a broad spectrum of activities which include

analgesics, cardioprotectants, antidiabetic, antiviral, antibacterial agents etc [3]. Natural products have been the source of inspiration for numerous pharmaceutical agents utilized for human health; however, the development of many natural product leads is often hindered due to the low quantities of material that can be harvested from plants and microbial sources [4]. Indeed without natural products, medicine would be lacking in therapeutic tools in several important clinical areas such as cardiovascular disease, immunoinflammatory disease, neurodegenerative disease etc. [5, 6]

In the last decade pathogenic microbial infectious agents have exhibited resistance to chemotherapeutics and antibiotic treatments triggering a renewed interest in antibacterial agents from natural sources which are effective against pathogenic bacteria [7]. Nature has provided varied resources for humans over the years including tools for the first attempts at therapeutic intervention [1]. Many medicinal plants have been screened extensively for their antimicrobial potential worldwide [8, 9].

Most plants possess limitless ability to synthesize aromatic substances have been reported to have antimicrobial activity [10].

Tecoma stans (family, Bignoniaceae) commonly known as yellow elder, yellow trumpet is a semi ever green ornamental

*Correspondence: iekpe@yahoo.com

¹Department of Biochemistry, College of Natural and Applied Sciences, Veritas University (The Catholic University of Nigeria), Abuja

Full list of author information is available at the end of the article.

Received: 07 Sep 2018, Accepted: 09 Dec 2018



shrub or tree [11]. It is a native of southern Texas, Latin America, Bahamas, and Trinidad. It has naturalised in much of tropical and subtropical Africa, Asia and Australia

Tecoma stans is an ornamental plant. It is planted and managed to enhance beauty of green belts and natural forests for purpose of providing recreation [11]. The leaves of *T. Stans* contain many active substances such as tannins, alkaloids; saponins etc have been shown to effectively reduce the symptoms of *diabetes mellitus* [12]. The plant is traditionally used in parts of Mexico for control of diabetes, digestive and urinary disorders [13]. It has been reported that tecomine an alkaloid present in *T. Stans* is responsible for the hypoglycaemic action [12]. *Costus. afer* Ker-Gawl. (Costaceae) is among 150 species of stout, perennial and rhizomatous herbs of the genus [14]. It can be found in the forest belt of Senegal, South Africa, Guinea, Niger, Sierra Leone and Nigeria [14, 15]. The plant is commonly called bush cane, irekeomode (Yoruba-Western part of Nigerian) and opete (Igbo-Eastern part of Nigeria) and mbriem (Ibibio/ Efik - Southern part of Nigeria). *C. afer* is a useful medicinal plant that is highly valued for its anti-diabetic, anti-inflammatory and anti-anthritic properties in South-East and South-West Nigeria. An infusion of the dried aerial parts is used to treat hypertension; the juice is taken orally for the treatment of cough and stomach-ache while the boiled tender leaves are used as soothing fomentation for rheumatic pains [16].

The pharmacological activities of many plants including *T. Stans* is derived from the presence of its phytochemical constituents such as alkaloids, saponins, flavonoids, tannins, glycosides, antioxidants, flavones, isoflavones, catechins, anthocyanidins, isothiocyanates, carotenoids, allyl sulfides, polyphenols [17].

This study was undertaken to demonstrate phytochemical and antibacterial potentials of *Tecoma stans* and *Costus afer* Kgwál cultivated in the southern part of Nigeria to determine their therapeutic potentials.

Materials and Method

Plant Material

Fresh leaves and roots of *Tecoma stans* were collected at Malabo campus, University of Calabar, Calabar. *Costus afer* leaves were collected from Eman Uruan, Uruan local government area of Akwa Ibom State, Nigeria. They were identified by Mr Frank I. Apojoye of Botany Department, University of Calabar and voucher samples were deposited at the herbarium of Botany Department, University of Calabar for reference. The leaves and roots were washed and shade dried for seven and ten days respectively. They were crushed into fine powder. The dried pow-

der of both plant parts were separately soaked in 80% ethanol for 72 hours with occasional agitation after which they were filtered through chess material and the Whatman NO.1 filter paper to obtain a homogenous filtrate. The filtrate was concentrated in vacuum at low temperature 37°-40° C to about one tenth of the original volume. The concentrates were allowed in an open water bath 40° C to evaporate the concentrate to complete dryness. A semi solid brown and dark green colored concentrates were obtained for roots and leaves respectively. They were stored in clean capped bottles in a refrigerator for further use.

Bacterial Isolates

Five (5) bacterial isolates (*Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Klebsilla pneumoniae*, *P.speceis*, *Shigella sonnei*, *Morganella morgani*, and *Maraxella catarrhlis* were obtained from the University of Calabar Teaching Hospital, Calabar. They were maintained on nutrient agar at 4° C in the refrigerator.

Screening for Antibacterial Activity

The agar-well diffusion technique as described by Irobi et al, [18] was employed. Bacterial inocula were prepared by giving each pure culture in nutrient broth for 18-24 hours at 37° C. Dilutions of the nutrient broth culture were prepared to obtain 0.5 Narfarland turbidity standards. 100ul of the nutrient broth culture was spread plated on the Muller Hinton agar plate. After spread plating 5mm diameter cork borer was used to make wells on the Muller Hinton agar plates. 1gm of the leaf and root extracts of *Tecoma stans* were dissolved in 2ml of dimethylsulphurdioxide (DMSO) respectively to achieve a concentration of 500mg/ml of extract. 100ul of the dissolved extracts were pipette in the agar wells. 100ul DMSO was also pipette into a separate well to serve as control. 10ug disc of gentamicin was also included to serve as antibiotic control. All plates were incubated at 37° C for 24 hours. Following incubation the diameter of zones of inhibition were measured and recorded in millimetres. The average of three independent determinations was recorded

Results and discussion

Discusion

The results of the antibacterial activity of the ethanol extract of the root of *T. stans* (Table 2) shows inhibitory activity over all selected bacterial organisms at zones of inhibition against 10ug/ml of the standard gentamicin.

The leaf extract (*T. Stans*) 416.7mg/ml exhibited highest inhibition 14.5mm for *S.aureus* against 16.5mm produced by 10ug/ml of the standard.

Table 1 Phytochemical Composition of Ethanolic Extract of *Tecoma stans* (Leaves/Root) and *Costus afer* K gwal

Chemical Constituents	Leaves <i>T.stans</i>	Root <i>T.stans</i>	Leaves <i>Costus afer</i>
Alkaloids	+	+++	–
Glycosides	++	+	+++
lavonoids	+++	++	+++
Saponins	++	+++	+

+ Slight presence, ++ Presence, +++ Strong presence, _ not detected

Table 2 Antibacterial activity of ethanolic extract of leaves of *Tecoma stans* on Different bacteria

Concentration (mg/ml)	Zone of Inhibition (mm)										Gentamycin (10µg/ml)
	2.6	5.2	10.4	20.8	41.7	83.3	25.0	187.5	312.5	416.7	
Organism											
<i>S.aureus</i>	0.00	1.50	4.00	5.00	7.00	9.00	10.00	12.00	12.50	14.50	16.5
<i>E.coli</i>	0.00	0.00	1.00	3.00	5.00	8.00	9.00	11.00	12.00	12.50	18.0
<i>P.aeruginosa</i>	0.00	0.00	0.00	0.00	3.00	6.00	8.00	8.50	8.50	9.00	17.5
<i>Proteus speies</i>	0.00	0.00	0.00	0.00	0.00	1.00	3.00	4.00	5.00	5.00	0.00
<i>K.pneumoniae</i>	0.00	0.00	0.00	0.00	0.50	2.00	4.00	4.00	4.00	5.00	15.00
<i>Moraxella catarrhis</i>	0.00	0.00	2.40	2.60	3.00	3.00	4.20	4.20	8.50	10.20	18.0
<i>Morganella morganii</i>	0.00	0.00	0.00	0.00	2.50	2.50	2.70	3.00	4.10	5.40	0.00
<i>Shigella sonnei</i>	0.00	0.00	0.00	0.00	0.00	0.00	1.50	3.00	4.50	7.40	22.40

Table 3 Antibacterial activity of ethanolic extract of root of *Tecoma stans* on Different bacteria

Concentration (mg/ml)	Zone of Inhibition (mm)										Gentamycin (10µg/ml)
	2.6	5.2	10.4	20.8	41.7	83.3	125.0	187.5	312.5	416.7	
Organism											
<i>S.aureus</i>	0.00	0.00	0.50	2.00	4.00	5.00	5.50	5.50	6.00	6.00	16.5
<i>E.coli</i>	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	18.0
<i>P.aeruginosa</i>	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	17.5
<i>Proteus speies</i>	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<i>K.pneumoniae</i>	0.00	0.00	1.00	4.00	6.00	6.00	7.00	7.00	7.50	8.00	15.00
<i>Moraxella catarrhis</i>	0.00	0.00	2.40	2.60	3.00	3.00	4.20	8.00	8.50	10.0	18.0
<i>Morganella morganii</i>	0.00	0.00	0.00	0.00	2.50	2.70	3.00	4.00	4.10	5.40	0.00
<i>Shigella sonnei</i>	0.00	0.00	0.00	0.00	0.00	1.50	3.00	4.00	4.50	7.40	22.40

Table 4 Antibacterial activity of ethanolic extract of leaves of *Costus afer* Ker gwal on Different bacteria

Concentration (mg/ml)	Zone of Inhibition (mm)										Gentamycin (10µg/ml)
	2.6	5.2	10.4	20.8	41.7	83.3	125.0	187.5	312.5	416.7	
Organism											
<i>S. aureus</i>	0.00	4.00	4.20	4.50	5.80	6.20	9.00	10.00	10.70	11.40	16.5
<i>E. coli</i>	0.00	0.00	0.00	0.00	0.00	4.70	5.40	5.50	5.80	6.80	18.0
<i>P. aeruginosa</i>	0.00	0.00	4.00	4.20	5.30	5.40	6.20	7.00	8.00	9.50	17.5
<i>Proteus speies</i>	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<i>K. pneumoniae</i>	0.00	0.00	0.00	0.00	0.00	0.00	3.40	4.00	4.00	5.00	15.00
<i>Moraxella catarrhis</i>	0.00	0.00	0.00	2.00	4.00	4.20	4.50	5.00	6.50	7.00	18.0
<i>Morganella morganii</i>	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	4.80	4.80	0.00
<i>Shigella sonnei</i>	0.00	0.00	0.00	0.00	3.50	3.50	4.00	4.50	4.50	7.20	22.40

The minimum inhibitory concentrations (MICs) of the leaf (*T. Stans*) extract for *S.aureus* was 5.21mg/ml indicating that a small dosage of the leaf extract will eliminate *S.aureus* while 10.4mg/ml was recorded for *E. coli*, 41.7mg/ml for *P. aeruginosa* and *K. pneumoniae* (Table 2). The leaf extract recorded an MIC of 83.3mg/ml for *Proteus speies* and highest of 416.7 mg/ml at 5mm compared to gentamycin which recorded no inhibitory activity for *Proteus speies*.

The ethanol extract of the root of *T. Stans* exhibited moderate to significant antibacterial activity against five (5) out of eight (8) tested bacterial organisms as compared to the standard gentamycin (16.5mm) Table 3. Other plant extracts e.g. *Delonix elata* and *Marrubium vulgare* have shown to be ineffective against *P. aeruginosa* [10] [19]. The maximum zone of inhibition (23 mm) was observed by Ramesh et al in 225 mg/mL concentration against *Pseudomonas aeruginosa* and the minimum zone of inhibition (16 mm) was observed in 75mg/mL concen-

tration against *Staphylococcus epidermidis* using methanolic extract of roots of *Tecoma stans*.

Table 4 show the result of antibacterial activity of *Costus afer* at 20.8mg/ml an indication the plant extract is a potential for respiratory tract infection (RTI).

Both extracts present broad spectrum antibacterial activity against the organisms tested. This suggests the extract of these plants exhibited a broad spectrum activity since it was active against both the Gram-positive and the Gram-negative organisms tested. The antimicrobial activity shown by this extract on the test organisms may be either due to the presence of alkaloids, flavonoids, saponins and glycosides (Table 1) [20]. [21] reports strong presence of flavonoids and saponins in ethanolic extract of *T.stans* leaves. The potentials for developing antimicrobials from plants seem rewarding as it will lead to the development of a phytomedicine against microbes. The ethanol extract of *T. stans* in combination with *C.afer* extract can be used to achieve significant inhibitory activity effects (polyherbal therapy) as antimicrobial agents in the treatment of bacterial infections.

Author details

¹Department of Biochemistry, College of Natural and Applied Sciences., Veritas University (The Catholic University of Nigeria), Abuja. ²Department of Biochemistry, Faculty of Basic Medical Sciences, University of Calabar, Cross River State, Nigeria.

References

- [1] Newman DJ, Cragg, Snader GM. Natural products as sources of new drugs over 30years from 1981-2010. *Journal Natural Products*. 2012;75:311–355.
- [2] Immune Rebalancing The Future of Immunosuppression Academic. Academic Press; 2016. Eduardo Penton-Arias I, David DH in Immune Rebalancing.
- [3] Dahanukar SA, Kulkarni RA, Rege NN. Pharmacology of medicinal plants and natural products. *Ind J Pharmacology*. 2000;32:81–118.
- [4] Christopher MR, Jeffrey JB, DK. David HS Natural Products Structural Diversity-I Secondary Metabolites: Organization and Biosynthesis in *Comprehensive Natural Products II*; 2010.
- [5] Harvey AL. Strategies for discovering drugs from previously unexplored natural products. *Drug Discov Today*. 2000;5(7):294–300.
- [6] Spainhour CB. Natural products. *Pharmaceutical Sciences Encyclopedia: Drug Discovery, Development, and Manufacturing*. 2010;p. 1–62.
- [7] Soulsby EJ. Resistance to antimicrobials in humans and animals. British Medical Journal Publishing Group; 2005.
- [8] Adedapo AA, Mogbojuri OM. Emikpe BO Safety evaluations of the aqueous extract of the leaves of *Moringa oleifera*. *Journal of Medicinal Plants Research*. 2009;3(8):586–591.
- [9] Basumatary S, Das AK, Raaman N, Sarma GD, Baalan L, Bora R. Phytochemical screening and antimicrobial activity of leaf and fruit extracts of *Hodgsonia heteroclita* (Roxb.) Hook. f. & Thomson. *J Int Acad Res Multidisciplinary*. 2015;3:358–66.
- [10] Pavithra PS, Janani VS, Charumathi KH, Indumathy R, Sirisha P, Verma RS. Antibacterial activity of plants used in Indian herbal medicine; 2010.
- [11] Nidhi M, Jain GC, Geeta P. Effect of *Tecoma stans* leaves on the reproductive system of male albino rats. *International Journal of Pharmacology*. 2010;6(20):152–156.
- [12] Aguilar-Santamaria L, Ramirez G, Nicasio P, Alegria-Rayes C, Herrera-Arellano A. Antidiabetic activities of *Tecoma stans* (L.) Juss. ex Kunth. *Journal of Ethnopharmacology*. 2009;124:284–288.
- [13] Costantinol LA, Barlocco D, Celotti F, El-Abady SA, Baranetti T, Maggi R, et al. Isolation and pharmacological activities of *T.stans* alkaloids. *Pharmaco*. 2003;58(9):781–785.
- [14] Edeoga HO, Okoli BE. Chromosome numbers of *Costus lucanusianus* (Costaceae) in Nigeria. *Folia Geobotanica*. 2000;35:315–318.
- [15] Burkill HM. The useful plants of West Tropical Africa: Rev Dalziels J.M. Royal Botanical Gardens Kew; 1985. 2nd ed. Families A-D.
- [16] Soladoye M, Oyesika O. A Textbook of Medicinal Plants From Nigeria. University of Lagos Press; 2008.
- [17] Harbone JB, Sumere CFV. The Chemistry and Biochemistry of Plant Proteins: Proceedings of the Phytochem. Soc. Symposium; Univ. of Ghent, Belgium, Sept. 1973. Acad. Press; 1975.
- [18] Irobi ON, Moo-Young M, Anderson WA. Antimicrobial activity of annatto (*Bixa orellana*) extract. *International Journal of Pharmacognosy*. 1996;34(2):87–90.
- [19] Masoodi MH, Ahmed B, Zargar IM, Khan SA, Khan S, Singh P. Antibacterial activity of whole plant extract of *Marrubium vulgare*. *African journal of Biotechnology*. 2008;7(2).
- [20] Odoemena CS, Luke MI, Ubon, Dg, Udotung, Ir. *Journal of Research in Bioscience*. 2008;4(1):7–10.
- [21] Govindappa M, Sadananda TS, Channabasava R, Jeevitha MK, Pooja KS, Vinay BR. *Journal of Phytology*. 2011;3(3):2075–6240. *Phytopharmacology*.