

COLLABORATIVE ASTEROID PHOTOMETRY AND LIGHTCURVE ANALYSIS AT OBSERVATORIES OAEGG, OAC, EABA, AND OAS

Fernando Mazzone

Observatorio Astronómico Salvador (MPC I20), Achalay 1469
X5804HMI Río Cuarto, Córdoba, ARGENTINA
Departamento de Matemática
Universidad Nacional de Río Cuarto, Córdoba, ARGENTINA
fmazzone@exa.unrc.edu.ar

Carlos Colazo

Grupo de Astrometría y Fotometría
Observatorio Astronómico Córdoba
Universidad Nacional de Córdoba, (Córdoba) ARGENTINA
Observatorio Astronómico El Gato Gris (MPC I19)
Tanti (Córdoba), ARGENTINA

Federico Mina

Grupo de Astrometría y Fotometría, Observatorio Astronómico
Córdoba, Universidad Nacional de Córdoba
Córdoba, ARGENTINA

Raúl Melia

Grupo de Astrometría y Fotometría, Observatorio Astronómico
Córdoba, Universidad Nacional de Córdoba
Córdoba, ARGENTINA

Julio Spagnotto

Observatorio El Catalejo (MPC I48)
Santa Rosa (La Pampa), ARGENTINA

Alejandro Bernal

Grupo de Astrometría y Fotometría
Observatorio Astronómico Córdoba
Universidad Nacional de Córdoba
Córdoba, ARGENTINA

(Received: 18 September)

Synodic rotation periods and amplitudes are reported for: 1874 Kacivelia, 15.951 ± 0.001 h, 0.21 ± 0.02 mag; 2055 Dvorak, 4.4052 ± 0.0003 h, 0.17 ± 0.04 mag; 2185 Guangdong, 21.089 ± 0.002 h, 0.19 ± 0.02 mag; and 8059 Deliyannis, 6.0041 ± 0.0003 h, 0.39 ± 0.04 mag. The absolute magnitude (H) and/or slope parameter (G) for some of these asteroids are also reported.

This paper presents the collaborative work among a group of amateur astronomers and undergraduate students gathered in two Argentinian associations: Grupo de Astrometría y Fotometría (GAF) and Asociación de Observatorios Argentinos de Cuerpos Menores (AOACM). The observatories and equipment used were:

- Estación Astrofísica de Bosque Alegre (EABA, MPC 821): 1.54-m Newtonian (NT) and Apogee Alta U9 CCD.
- Observatorio Astronómico Córdoba (OAC, MPC 822): 0.35-m Schmidt-Cassegrain (SCT) and SBIG ST7 CCD.
- Observatorio Astronómico El Gato Gris (OAEGG, MPC I19): 0.35-m Schmidt-Cassegrain (SCT) and SBIG ST10 CCD.
- Observatorio Astronómico Salvador (OAS, MPC I20): 0.2-m Schmidt-Newtonian (SNT) and Starlight ST7-XME CCD.

All images were unfiltered, dark, bias and flat-field corrected, and then measured using Astrometrica software (Raab, 2013). We used *Periodos* software (Mazzone, 2012a) for the period analysis.

We find that this software presents some novelties in the mathematical processing of the data. These are discussed in the appendix along with some details regarding our methods.

All targets were selected from the “Potential Lightcurve Targets” web site list on the Collaborative Asteroid Lightcurve Link site (CALL; Warner *et al.*, 2011) as a favorable target for observation and with no previously reported period in the Lightcurve Database (LCDB, Warner *et al.*, 2009).

The lightcurve figures contain the following information: 1) the estimated period and amplitude, 2) a 95% confidence interval regarding the period estimate, 3) RMS of the fitting, 4) estimated amplitude and amplitude error, 5) the Julian time corresponding to 0 rotation phase, and 6) the number of data points. In the reference boxes the columns represent, respectively, the marker, observatory MPC code, session date, session off-set, and number of data points. See the appendix for a description of the off-sets and reduced magnitudes.

8059 Deliyannis. We collected 548 data points in five different sessions. The derived period and amplitude were 6.0041 ± 0.0003 h and 0.39 ± 0.04 mag. There is a lack of data between phase angles 0.63 and 0.7. We estimate the absolute magnitude H to be 11.92 mag. Previously reported values were 11.8 mag (MPO 233564) and 12.0 (MPC 30957).

1874 Kacivelia. We observed this asteroid between phase angles 17° to 2° . We obtained a period of 15.951 ± 0.001 h and amplitude of 0.21 ± 0.02 mag. The MPCORB file gives $H = 11.2$ (MPO 250216). We estimate a value of $H = 11.4$. Given the wide range of phase angles covering our observations, we considered it appropriate to find the slope parameter, G (see the appendix section for details). The MPCORB gives a default value of $G = 0.15$. We found $G = 0.24$ produces a better fit to our data.

2055 Dvorak. Analysis of our data found a period of 4.4052 ± 0.0003 h and amplitude and 0.17 ± 0.04 mag with a large dispersion among the offsets. The calculated absolute magnitude is 12.81. MPO 259350 reports $H = 12.6$ and MPC 17264, $H = 13.5$.

2185 Guangdong. This was a difficult target due to its relatively long rotation period. Unfortunately, the second half of the lightcurve has substantially fewer data than the first half. We derived a period of 21.089 ± 0.002 h and amplitude 0.19 ± 0.02 mag. We computed an absolute magnitude of $H = 11.57$. The MPCORB file gives $H = 11.3$ using $G = 0.15$. We found that $G = 0.33$ produces a smaller root-square norm for off-sets.

Appendix: Data Analysis Strategy

In this section, we describe the method used for the data analysis, which has some differences with the usual methodology in similar work. We have successfully used these techniques before (Ambrosioni *et al.*, 2011; Oey *et al.*, 2012).

We programed a set of MATLAB[®] functions that implemented the calculations described below using functions from *Periodos* and *orbit_calc* (Mazzone, 2012a; Mazzone, 2012b).

Suppose that m_i^j , for $j = 1, \dots, N$ and $i = 1, \dots, M_j$, are the measured magnitudes for the asteroid corresponding to times t_i^j . Here N is the number of different sessions and M_j , $j = 1, \dots, N$, is the number of data points in the session j . By session we mean the data collected by a unique observatory on a single night.

First we perform some corrections on the data. More specifically, times t_i^j were light-time corrected and magnitudes were corrected to unity distance and normalized to the zero phase angle by applying standard formulas (Dymock, 2007). This reduction requires some orbital calculations, which are made by an adaptive collocation method that solves the n-body problem. Sun, planets and Moon were modeled as point masses.

Second we fit the model function

$$f(t_i^j) = \alpha^j + a_0 + \sum_{k=1}^n a_k \cos\left(\frac{2k\pi t_i^j}{T}\right) + b_k \sin\left(\frac{2k\pi t_i^j}{T}\right)$$

to the observed data. More precisely, we look for parameters α^j , a_k , b_k , and T that minimize

$$\sum_{j=1}^N \sum_{i=1}^{M_j} |m_i^j - f(t_i^j)|^2$$

The fitted value of T and a_0 can be interpreted as being the synodic rotation period of the asteroid and the absolute magnitude H , respectively. The parameters α^j depend on the sessions and represent the offsets among sessions. Usually one wants them to be zero. In order to obtain a well-posed problem, we need to introduce an extra condition. We adopted the restriction that the offsets α^j have a zero mean, i.e. $\sum \alpha^j = 0$. We think that this is a plausible assumption, if we consider offsets random normally distributed variables. However this affirmation is false in general. For example, an inaccurate determination of the slope parameter G induces a pattern in the offsets. We think that the value of G such that the offsets squares sum $(\alpha^1)^2 + \dots + (\alpha^N)^2$ are minimized gives a good estimate for the G parameter. In this way we obtained the value of G reported for 1874 Kacivelia and 2185 Guangdong.

We note that our methods incorporate a Fourier algorithm (Harris *et al.*, 1989) and simultaneously adjust the offsets. This is a non-linear curve fitting problem and we use the native `lsqcurvefit` MATLAB® function for solving it.

References

Ambrosioni, C., Colazo, C., and Mazzone, F. (2011). "Period Determination for 1996 Adams and 2699 Kalinin by AOACM." *Minor Planet Bul.* **38**, 102-102.

Dymock, R. (2007). "The H and G magnitude system for asteroids." *Journal of the British Astronomical Association.* **117**, 342-343.

Harris, A.W., Young, J.W., Bowell, E., Martin, L. J., Millis, R.L., Poutanen, M., Scaltriti, F., Zappala, V., Schober, H.J., Debehogne, H., and Zeigler, K. (1989). "Photoelectric Observations of Asteroids 3, 24, 60, 261, and 863" *Icarus* **77**, 171-186.

Mazzone, F.D. (2012a). *Periodos* software, version 1.0. <http://www.astrosurf.com/salvador/Programas.html>

Mazzone, F.D. (2012b). *orbit_calc* software, version 2.0. <http://www.astrosurf.com/salvador/Programas.html>

Oey, J., Colazo, C., Mazzone, F., and Chapman, A. (2012). "Lightcurve Analysis of 918 Itha and 2008 Konstitutsiya." *Minor Planet Bul.* **39**, 1-2.

Raab, H. (2013). *Astrometrica* software, version 4.8.2.405, <http://www.astrometrica.at/>

Warner, B.D., Harris, A.W., and Pravec, P. (2009). "The Asteroid Lightcurve Database." *Icarus* **202**, 134-146.

Warner, B.D., Harris, A.W., and Pravec, P. (2011). Lightcurve Targets. <http://www.minorplanet.info/call.html>



