

# Infinite locally random graphs

Pierre Charbit\* and Alex D. Scott†

## Abstract

Motivated by copying models of the web graph, Bonato and Janssen [3] introduced the following simple construction: given a graph  $G$ , for each vertex  $x$  and each subset  $X$  of its closed neighbourhood, add a new vertex  $y$  whose neighbours are exactly  $X$ . Iterating this construction yields a limit graph  $\uparrow G$ . Bonato and Janssen claimed that the limit graph is independent of  $G$ , and it is known as the *infinite locally random graph*. We show that this picture is incorrect: there are in fact infinitely many isomorphism classes of limit graph, and we give a classification. We also consider the inexhaustibility of these graphs.

## 1 Introduction

The *Rado graph*  $\mathcal{R}$  is the unique graph with countably infinite vertex set such that, for any disjoint pair  $X, Y$  of finite subsets of vertices, there is a vertex  $z$  that is joined to every vertex in  $X$  and no vertex in  $Y$ . If  $0 < p < 1$ , and  $G$  is a random graph in  $\mathcal{G}(\mathbb{N}, p)$ , then with probability 1 we have  $G \cong \mathcal{R}$ . For this reason, the Rado graph is also known as *the infinite random graph* (see [5] for a survey).

The Rado graph can be obtained deterministically by beginning with any finite (or countably infinite) graph  $G$  and iterating the following construction:

[E1] For every finite subset  $X$  of  $V(G)$  add a vertex  $y$  with neighbourhood  $N(y) = X$ .

---

\*Laboratoire Camille Jordan, Université Claude Bernard, Lyon 1, France (email : charbit@univ-lyon1.fr)

†Mathematical Institute, University of Oxford, 24-29 St Giles', Oxford, OX1 3LB, UK (email : scott@maths.ox.ac.uk)

Here  $N(x) = \{y \in V(G) : xy \in E(G)\}$  is the *neighbourhood* of  $x$ ; we also write  $N[x] = N(x) \cup \{x\}$  for the *closed neighbourhood* of  $x$ .

Motivated by copying models of the web graph, Bonato and Janssen [3] (see also [1] and [4]) introduced the following interesting construction. For a finite graph  $G$ , the *pure extension*  $PE(G)$  of  $G$  is obtained from  $G$  by the following construction:

[E2] For every  $x \in V(G)$  and every finite  $X \subseteq N[x]$  add a vertex  $y$  with neighbourhood  $N(y) = X$ .

Iterating this construction, we obtain a limit graph, denoted by  $\uparrow G$ .

Bonato and Janssen ([3], Theorem 3.3) claimed that  $\uparrow G \cong \uparrow H$  for every pair  $G, H$  of finite graphs. The (claimed) unique limit graph, which has become known [1] as the *infinite locally random graph* (see Proposition 1 below for the reason for this name). As we show below, Bonato and Janssen's claim is incorrect. There are in fact infinitely many limit graphs  $G$  (for instance,  $\uparrow C_5, \uparrow C_6, \uparrow C_7, \dots$  are all distinct), and we give a simple criterion that determines when  $\uparrow G \cong \uparrow H$ .

In the next section, we give a few simple properties of limit graphs  $\uparrow G$ ; we prove our classification result in section 3. Finally, in section 4, we prove that for every finite  $G$ ,  $\uparrow G$  is inexhaustible, that is  $(\uparrow G) \setminus x \cong \uparrow G$  for all  $x \in V(\uparrow G)$ . This corrects another result from [3].

## 2 Simple properties of $\uparrow G$

We begin with some notation. We shall refer to the vertices  $y$  that are introduced in [E2] above with neighbourhoods contained in  $N[x]$  as *clones* of  $x$ . Thus a vertex of degree  $d$  in  $G$  has  $2^{d+1}$  clones in  $PE(G)$  (note that we take all subsets of the *closed* neighbourhood  $N[x]$ ), and  $PE(G)$  contains  $|G|$  isolated vertices, each one a clone of a different vertex from  $G$ . As indicated above, iterating construction [E2] gives a sequence of graphs  $G \subseteq PE(G) \subseteq PE^2(G) \subseteq \dots$ , where  $PE^n(G) = PE(PE^{n-1}(G))$ ; we write  $\uparrow G$  for the limit of this sequence. We define the *level*  $L(x)$  of a vertex of  $\uparrow G$  to be the least integer  $k$  such that it is contained in  $PE^k(G)$  (where  $L(x) = 0$  for all  $x \in V(G)$ ), and for a finite subset  $X \subseteq V(\uparrow G)$ , we write  $L(X) = \max_{x \in X} L(x)$ . We also write  $L^{(k)}(\uparrow G)$  for the vertices of level  $k$  in  $\uparrow G$ , and  $L^{(\leq k)}(\uparrow G)$  for the vertices of level  $k$  or less. Note that, by the construction,  $L^{(k)}(\uparrow G)$  is an independent set for every  $k \geq 1$ .

Given a graph  $H$ , a graph  $G$  is *locally  $H$*  if, for every vertex  $x$  of  $G$ , the graph induced by the neighbourhood  $N(x)$  of  $x$  is isomorphic to  $H$ .

Bonato and Janssen note the following property of the construction defined above.

**Proposition 1.** [3] *For every finite graph  $G$ ,  $\uparrow G$  is locally  $\mathcal{R}$*

*Proof.* For every  $x \in V(\uparrow G)$ , and every  $X$  and  $Y$  finite disjoint subsets of  $N(x)$ , we want to find a vertex  $z$  such that  $z$  is adjacent to every vertex in  $X$  and to none in  $Y$ . This is possible by the definition of  $\uparrow G$  by taking a suitable vertex  $z$  of level  $L(X \cup Y) + 1$ .  $\square$

Since  $\mathcal{R}$  is the (unique) infinite random graph, it therefore makes sense to refer to  $\uparrow G$  as an *infinite locally random graph*.

**Corollary 2.** *Let  $G$  be a finite graph. Then  $\uparrow G$  is  $\aleph_0$ -universal (that is,  $\uparrow G$  contains every countable graph  $H$  as an induced subgraph).*

Another easy but important remark concerns the distance between vertices.

**Proposition 3.** *Let  $G$  be a finite graph and  $x$  and  $y$  two vertices of  $PE^k(G)$ , for some integer  $k \geq 0$ . Then the distance between  $x$  and  $y$  is the same in  $PE^k(G)$  and in  $\uparrow G$ .*

*Proof.* It is sufficient to note that the pure extension construction [E2] does not change the distance between vertices.  $\square$

We also note the following simple property.

**Lemma 4.** *Let  $G$  be a finite graph and  $x$  a vertex of  $\uparrow G$ . Let  $X$  be a finite subset of  $N(x)$ . Then there exists a vertex  $y$  with  $L(y) \leq L(X)$  such that  $X \subseteq N[y]$ .*

*Proof.* Let  $x_0$  be a vertex of minimal level with  $X \subseteq N[x_0]$ . If  $L(x_0) \leq L(X)$  then we can take  $y = x_0$ . Otherwise,  $L(x_0) > L(X)$  and so  $x_0 \notin X$ . But  $x_0$  was constructed on level  $L(x_0)$  as the clone of some vertex  $x_1$  with  $L(x_1) < L(x_0)$ . In particular,  $N(x_0) \cap L^{(<L(x_0))}(\uparrow G) \subseteq N[x_1]$  and so  $X \subseteq N[x_1]$ , which contradicts the minimality of  $L(x_0)$ .  $\square$

For  $x \in V(\uparrow G)$ , we write

$$N^-(x) = N(x) \cap L^{(<L(x))}(\uparrow G).$$

Note that  $N^-(x)$  is the set of neighbours assigned to  $x$  at time  $L(x)$ , when  $x$  is first introduced. We say that a subgraph  $G_1$  of  $G$  is *good* if it is an induced subgraph of  $G$  and, for all  $x$  in  $V(G_1)$ ,  $N^-(x) \subseteq V(G_1)$ . Equivalently,  $G_1$  is an induced subgraph such that  $N(y) \cap V(G_1) \subseteq N^-(y)$  for all  $y \in V(G) \setminus V(G_1)$ .

In this context, Lemma 4 gives the following result.

**Lemma 5.** *Let  $G$  be a finite graph and suppose that  $H$  is a good subgraph of  $\uparrow G$ . Then*

$$\forall x \in V(\uparrow G), \exists y \in V(H) \text{ such that } N(x) \cap V(H) \subseteq N[y] \cap V(H)$$

*Proof.* We can assume that  $x \notin V(H)$ . Let  $X = N(x) \cap V(H)$ . Then  $X \subseteq N^-(x)$ , and by Lemma 4 there exists  $y$  of level at most  $L(X)$  with  $X \subseteq N[y]$ . If  $L(y) = L(X)$  then, since the levels are independent sets and  $X \subseteq N[y]$ ,  $y$  must belong to  $X$ , and thus to  $H$ . If  $L(y) < L(X)$ , then  $y$  belongs to  $H$  as  $H$  is a good subgraph of  $\uparrow G$ .  $\square$

### 3 Classification

We now investigate when  $\uparrow G$  and  $\uparrow H$  are isomorphic. In [3], the authors claim that  $\uparrow G \cong \uparrow H$  for any pair of finite graphs  $G$  and  $H$  (this is their Theorem 3.3). Here we disprove this. Their proof seems to fail on page 209 at the end of the first paragraph: the equality  $H_{n+1} - S \cong G_1 \uplus \overline{K_m}$  does not hold because these vertices can be linked by edges. Moreover, it is not clear why this equality would imply  $H - S \cong \uparrow(G_1 \uplus \overline{K_m})$  on the following line, as some vertices in  $H$  can be constructed by cloning elements in  $S$ .

We begin with the following useful consequence of Lemma 5.

**Theorem 6.** *Let  $G$  and  $H$  be finite graphs. Suppose that  $G_1 \supseteq G$  is a good subgraph of  $\uparrow G$  and  $H_1 \supseteq H$  is a good subgraph of  $\uparrow H$ . If  $G_1 \cong H_1$  then  $\uparrow G \cong \uparrow H$*

*Proof.* Let  $\phi : V(G_1) \rightarrow V(H_1)$  be an isomorphism (note that, as  $G_1$  and  $H_1$  are good, they are induced subgraphs of  $\uparrow G$  and  $\uparrow H$ , respectively, so this is an isomorphism between induced subgraphs). Using a classical ‘back

and forth' argument, we extend  $\phi$  one vertex at a time until, in the limit, we obtain an isomorphism between  $\uparrow G$  and  $\uparrow H$ . Let  $x \in V(\uparrow G)$  be a vertex of minimal level with  $x \notin V(G_1)$ . By Lemma 5, there exists  $y \in V(G_1)$  such that  $N(x) \cap V(G_1) \subseteq N[y] \cap V(G_1)$ . Let  $z \notin V(H_1)$  be a clone of  $\phi(y)$  with

$$N^-(z) = N(z) \cap V(H_1) = \phi(N(x) \cap V(G_1)).$$

Such a clone is easily found: let  $k = L(V(H_1))$ , and take the clone of  $\phi(y)$  on level  $k + 1$  with exactly this neighbourhood in  $L^{(\leq k)}(\uparrow H)$ . Then  $V(H_1) \cup \{z\}$  induces a good subgraph of  $\uparrow H$  and, by minimality of  $x$ ,  $V(G_1) \cup \{x\}$  induces a good subgraph of  $\uparrow G$ . We can therefore extend  $\phi$  by setting  $\phi(x) = z$ . Repeating the construction in alternate directions we clearly obtain an isomorphism between  $\uparrow G$  and  $\uparrow H$ .  $\square$

We shall say that a vertex  $x$  of a graph  $G$  is *inessential* if there exists  $y \in V(G)$ ,  $y \neq x$  such that  $N(x) \subseteq N[y]$ . A graph is *essential* if it contains no inessential vertices. Given a graph  $G$ , a sequence of vertices  $x_1, \dots, x_k$  is a *maximal sequence of removals* if  $x_i$  is inessential in  $G \setminus \{x_1, \dots, x_{i-1}\}$  for each  $i$ , and  $G \setminus \{x_1, \dots, x_k\}$  is an essential graph.

We shall show below that every maximal sequence of removals yields the same essential graph (up to isomorphism). However, we first prove a simple lemma. We say that two vertices  $x$  and  $y$  in a graph  $G$  are *equivalent* if  $N(x) = N(y)$  or  $N[x] = N[y]$ . Equivalently,  $N(x) \subseteq N[y]$  and  $N(y) \subseteq N[x]$ . Clearly, if  $x$  and  $y$  are equivalent in  $G$  then  $G \setminus x \cong G \setminus y$ , with the obvious isomorphism given by exchanging  $x$  for  $y$  and leaving the other vertices fixed.

Equivalent vertices play an important role in the removal of inessential vertices.

**Lemma 7.** *Suppose that  $x$  and  $y$  are inessential in a graph  $G$ , but  $x$  is not inessential in  $G \setminus y$ . Then  $x$  and  $y$  are equivalent.*

*Proof.* Note first that since  $x$  and  $y$  are inessential in  $G$ , there are  $x'$  and  $y'$  such that  $N(x) \subseteq N[x']$  and  $N(y) \subseteq N[y']$ . If  $x' \neq y'$  then considering the vertex  $x'$  in  $G \setminus y$  shows that  $x$  is inessential in  $G \setminus y$ , a contradiction. So  $x' = y'$ , and  $N(x) \subseteq N[y]$ .

Now consider  $y'$ . If  $y' \neq x$  then  $N(x) \subseteq N[y] = \{y\} \cup N(y) \subseteq \{y\} \cup N[y']$  implies that  $N(x) \setminus y \subseteq N[y']$ , and so  $y'$  shows that  $x$  is inessential in  $G \setminus y$ , a contradiction. Thus we have  $y' = x$ , and so  $N(y) \subseteq N[x]$ . It follows that  $x$  and  $y$  are equivalent.  $\square$

We now show that maximal sequences of removals define a unique graph up to isomorphism.

**Theorem 8.** *Suppose that  $x_1, \dots, x_k$  and  $y_1, \dots, y_l$  are two maximal sequences of removals in a finite graph  $G$ . Then  $G \setminus \{x_1, \dots, x_k\} \cong G \setminus \{y_1, \dots, y_l\}$ .*

*Proof.* We claim that we can modify the sequence  $\{y_1, \dots, y_l\}$  to obtain the sequence  $\{x_1, \dots, x_k\}$  without changing the isomorphism type of the resulting essential graph  $G \setminus \{y_1, \dots, y_l\}$ .

Suppose first that  $x_1 \notin \{y_1, \dots, y_l\}$ . Then (by maximality)  $x_1$  is inessential in  $G$  but not in  $G \setminus \{y_1, \dots, y_l\}$ . Let  $i$  be maximal such that  $x_1$  is inessential in  $G \setminus \{y_1, \dots, y_i\}$ . Then, by Lemma 7,  $x_1$  and  $y_{i+1}$  are equivalent in  $G \setminus \{y_1, \dots, y_i\}$ , and so we can replace  $y_{i+1}$  by  $x_1$  in the sequence  $y_1, \dots, y_l$ , without effecting the isomorphism type of  $G \setminus \{y_1, \dots, y_l\}$  (the isomorphism is given by exchanging  $x_1$  and  $y_{i+1}$ ). We may therefore assume that  $x_1 \in \{y_1, \dots, y_l\}$ .

We now show that we can modify  $y_1, \dots, y_l$  so that  $y_1 = x_1$ . Suppose that  $x_1 = y_{i+1}$  for some  $i \geq 1$ . If there exists some  $0 \leq j < i - 1$  such that  $x_1$  is inessential in  $G \setminus \{y_1, \dots, y_j\}$  and not in  $G \setminus \{y_1, \dots, y_{j+1}\}$ , Lemma 7 implies that  $x_1$  and  $y_{j+1}$  are equivalent in  $G \setminus \{y_1, \dots, y_j\}$ . Therefore we can exchange them in the sequence. We can repeat this operation as long as such an integer  $j$  exists, and thus we can assume that  $x_1 = y_{i+1}$  is inessential in  $G \setminus \{y_1, \dots, y_j\}$  for all  $j \leq i$ . Now, if  $y_i$  is not inessential in  $G \setminus \{y_1, \dots, y_{i-1}, x_1\}$  then (as it is inessential in  $G \setminus \{y_1, \dots, y_{i-1}\}$ ) Lemma 7 shows that  $x_1$  and  $y_i$  are equivalent in  $G \setminus \{y_1, \dots, y_{i-1}\}$ . It is clear that we may therefore exchange  $y_i$  and  $y_{i+1} = x_1$  in the sequence  $y_1, \dots, y_l$ . Repeating this argument, we move  $x_1$  forward in the sequence  $y_1, \dots, y_l$  until  $x_1 = y_1$ .

Finally, if  $x_1 = y_1$ , we can work instead with the graph  $G \setminus x_1$  and the sequences  $x_2, \dots, x_k$  and  $y_2, \dots, y_l$ , continuing until one (and hence both) of the sequences is exhausted.  $\square$

We shall denote the (isomorphism type of the) subgraph of  $G$  obtained by deleting a maximal sequence of removals  $\downarrow G$ . For instance,  $\downarrow K_n = \downarrow C_4 = K_1$ , but  $\downarrow C_k = C_k$  for all  $k \geq 5$ .

We next show that inessential vertices have no effect on limit graphs.

**Corollary 9.** *Let  $G$  be a finite graph and  $x$  an inessential vertex of  $G$ . Then  $\uparrow G \cong \uparrow(G \setminus x)$*

*Proof.* Let  $H = G \setminus x$ . Since  $x$  is inessential, there exists  $y$  in  $G$  such that  $N(x) \subseteq N[y]$  in  $G$ . In  $\uparrow H$ ,  $y$  has a clone  $x'$  such that  $N^-(x') = N(x) \cap V(G)$ . Clearly  $G_1 = G$  is a good subgraph of  $\uparrow G$  and  $V(H) \cup \{x'\}$  induces a good subgraph  $H_1$  of  $\uparrow H$ . Thus it suffices to apply Theorem 6 to  $G_1$  and  $H_1$ .  $\square$

Corollary 9 implies the following theorem.

**Theorem 10.** *Let  $G$  be a finite graph. Then  $\uparrow G \cong \uparrow(\downarrow G)$*

If  $H$  is an induced subgraph of  $\uparrow G$ , then we define two kinds of transformations on this subgraph, called *reductions*.

- (i) Delete an inessential vertex of  $H$ .
- (ii) For a pair of vertices  $x \in V(H)$  and  $y \notin V(H)$  with  $N(x) \cap V(H) \subseteq N(y) \cap V(H)$ , replace  $H$  by the subgraph of  $\uparrow G$  induced by  $(V(H) \setminus x) \cup \{y\}$ .

**Lemma 11.** *If  $H$  is a finite induced subgraph of  $\uparrow G$ , it is possible to apply a sequence of reductions to transform  $H$  into a subgraph of  $G$ .*

*Proof.* Define the *weight*  $w(H')$  of an induced subgraph of  $\uparrow G$  by

$$w(H') = \sum_{v \in V(H')} L(v).$$

If  $w(H) = 0$  then  $H$  is a subgraph of  $G$ . If  $w(H) > 0$ , then we look for a reduction that decreases the weight or the number of vertices. If  $H$  contains an inessential vertex, then delete it (this can occur at most  $|H| - 1$  times). Otherwise, let  $x \in V(H)$  be a vertex of highest level. Then  $N(x) \cap V(H) = N(x) \cap V(H) \cap L^{(<L(x))}(\uparrow G)$ , as  $L^{(L(x))}(\uparrow G)$  is an independent set. Since  $x$  was built at level  $L(x)$  as the clone of some vertex  $y$  that satisfies  $N(x) \cap V(H) \cap L^{(<L(x))}(\uparrow G) \subseteq N[y] \cap V(H)$  and  $L(y) < L(x)$ , we can replace  $x$  by  $y$ , to obtain  $H'$  with  $w(H') < w(H)$ . Repeating this process, we eventually obtain an induced subgraph of  $\uparrow G$  with weight 0 which, as already noted, is a subgraph of  $G$ .  $\square$

We are now ready to prove our main result.

**Theorem 12.** *Let  $G$  and  $H$  be finite graphs. Then  $\uparrow G \cong \uparrow H \iff \downarrow G \cong \downarrow H$*

*Proof.* By Theorem 10, we may assume that  $G$  and  $H$  do not contain any inessential vertices, that is  $\downarrow G = G$  and  $\downarrow H = H$ . Suppose that  $\uparrow G \cong \uparrow H$ , and fix an isomorphism.

Let  $\{1, 2, \dots, n\}$  be the vertices of  $G$ . We partition the vertices of  $\uparrow G$  into  $n$  classes in the following way. For  $i = 1, \dots, n$ , let  $A_{i,0} = \{i\}$ , and for  $j \geq 1$ , let  $A_{i,j}$  be the vertices of  $\uparrow G$  which are clones of vertices in  $A_{i,j-1}$ . We then define  $A_i = \bigcup_{j=0}^{\infty} A_{i,j}$ . Thus  $A_i$  is the smallest set of vertices containing  $i$  and closed under taking clones. It is easy to see that, for  $i \neq k$ , there is an edge between class  $A_i$  and  $A_k$  if and only if there is an edge between  $i$  and  $k$  (as creating a clone cannot create adjacencies between a new pair of classes). We shall say that *edges between classes respect  $G$* .

Now consider an isomorphic embedding  $\phi$  of  $G$  into  $\uparrow G$ . We say that  $\phi$  is *good* if  $\phi(i) \in A_i$  for every  $i \in V(G)$ . Suppose that  $\phi$  is good and let  $G'$  be the image of  $G$  under  $\phi$ . If we apply a type (ii) reduction to some vertex of  $G'$ , say  $v_i := \phi(i)$ , then it is replaced by a vertex  $x$  such that  $N(x) \cap V(G') \supseteq N(v_i) \cap V(G')$ . Let  $A_j$  be the class containing  $x$ . Since  $\phi$  is good, there is an edge between  $A_j$  and  $A_k$  whenever  $k \in N(i)$ . Since edges between classes respect  $G$ , this implies  $N[j] \supseteq N(i)$ . But since we assumed that  $G$  contains no inessential vertices, this is possible only if  $i = j$ . Indeed,  $N(x) \cap V(G') = N(v_i) \cap V(G')$ , or else we would introduce edges between new pairs of classes. It follows that we obtain a good embedding  $\phi'$  of  $G$  by setting  $\phi'(i) = x$  and  $\phi'(j) = \phi(j)$  otherwise. This remains true for any sequence of reductions starting from a good embedding. In particular, any sequence of reductions starting from  $G$  produces an induced copy of  $G$  (note that reductions of type (i) are not possible at any stage).

By Lemma 11, any induced subgraph of  $\uparrow H$  isomorphic to  $G$  can be reduced to a subgraph of  $H$ . It follows that  $G$  must be isomorphic to a subgraph of  $H$ . Arguing similarly the other way round, we see that  $H$  is isomorphic to a subgraph of  $G$ , and so  $G \cong H$ .  $\square$

Now it is clear that  $\uparrow G$  is not independent of  $G$ : it suffices to consider two circuits of different length (larger than 4). In fact, Theorem 12 immediately gives the following classification of possible limit graphs.

**Corollary 13.** *The isomorphism classes of limit graphs  $\uparrow G$  of finite graphs  $G$  are in bijective correspondence with the class of essential finite graphs.*



## 4 Inexhaustibility

A graph  $G$  is *inexhaustible* if  $G \setminus x \cong G$  for every vertex  $x \in V(G)$ . For instance, the infinite complete graph  $K_\omega$  and its complement are trivially inexhaustible; the Rado graph  $\mathcal{R}$  is also inexhaustible. On the other hand, the infinite two-way path is not inexhaustible, as deleting any vertex increases the number of components. For results on inexhaustible graphs, see Pouzet [7], El-Zahar and Sauer [6] and Bonato and Delić [2].

Bonato and Janssen [3] consider the inexhaustibility of infinite graphs satisfying various properties, and claim a rather general result. Let us define two properties of (infinite) graphs as follows. We say that a graph  $G$  has *Property A* if it satisfies the following condition.

- (A) For every vertex  $x$  of  $G$ , every finite  $X \subseteq N[x]$ , and every finite  $Y \subseteq V(G) \setminus X$ , there is a vertex  $z \notin X \cup Y$  such that  $X \subseteq N(z)$  and  $Y \cap N(z) = \emptyset$ ,

and  $G$  has *Property B* if it satisfies the following.

- (B) For every vertex  $x$  of  $G$ , every finite  $X \subseteq N(x)$ , and every finite  $Y \subseteq V(G) \setminus X$ , there is a vertex  $z \notin X \cup Y$  such that  $X \subseteq N(z)$  and  $Y \cap N(z) = \emptyset$ .

Note that the only difference between (A) and (B) is that (A) is concerned with closed neighbourhoods, while (B) is only concerned with neighbourhoods. Clearly Property A implies Property B; furthermore, for any finite  $G$ , it is clear from the constructive step [E2] that  $\uparrow G$  has Property A (and therefore Property B).

Bonato and Janssen ([3], Theorem 4.1) claim that every graph with Property B is inexhaustible. However, there is a simple counterexample to this assertion: let  $G$  be the Rado graph  $\mathcal{R}$  with an additional isolated vertex  $x$ . Since the Rado graph is connected, and  $G$  is not, it is clear that  $G \setminus x \not\cong G$ . (The proof of Bonato and Janssen in [3] appears to fail with the definition of their sets  $S_i$ .)

In fact, even the stronger Property A does not imply that a graph is inexhaustible. Consider the graph  $G$  defined by starting from the path  $x_1x_2x_3x_4$  of length 3, and alternating the pure extension construction [E2] with the following step.

- [E3] For every pair of vertices  $\{x, y\} \neq \{x_1, x_4\}$ , add a vertex  $z$  with  $N(z) = \{x, y\}$ .

Note that  $x_1$  and  $x_4$  are at distance 3 in the initial graph. The pure extension step [E2] does not change the distance between vertices, while [E3] does not create a path of length 2 from  $x_1$  to  $x_4$ . Thus  $x_1$  and  $x_4$  are at distance 3 in the limit graph. On the other hand, there are infinitely many paths of length 2 between any other pair of vertices. Thus  $G \setminus \{x_1, x_4\} \not\cong G$ , and so  $G$  cannot be inexhaustible (if  $G$  is inexhaustible, then clearly  $G \setminus X \cong G$  for every finite  $X \subseteq V(G)$ ).

On the positive side, we can show that for any finite  $G$ , the limit graph  $\uparrow G$  is actually inexhaustible.

**Theorem 14.** *For every finite graph  $G$ ,  $\uparrow G$  is inexhaustible.*

*Proof.* Let  $v$  be any vertex of  $\uparrow G$ . We shall show that  $\uparrow G \cong (\uparrow G) \setminus v$ . Note that since  $\uparrow G \cong \uparrow PE^{L(v)}(G)$ , we can replace  $G$  by  $PE^{L(v)}(G)$ , and so we may assume that  $v \in V(G)$ .

On the first level above  $G$ ,  $v$  has a clone  $v'$  with  $N(v) \cap G = N(v') \cap G$ . Thus we have an isomorphism between  $G_1 = G$  and  $G_2 = G \setminus v \cup \{v'\}$ . It is clear that  $G_1$  and  $G_2 \cup \{v\}$  are good subgraphs. We will extend this isomorphism by a ‘back and forth’ argument.

Suppose we are given a partial isomorphism  $\phi$  between two subgraphs  $G_1$  and  $G_2$  of  $\uparrow G$ , with the following properties:

1.  $G_1$  and  $G_2 \cup \{v\}$  are good subgraphs of  $\uparrow G$
2.  $V(G) \subseteq V(G_1)$ ,  $V(G) \setminus v \subseteq V(G_2)$  and  $v \notin V(G_2)$
3. There is a vertex  $\tilde{v} \in V(G_2)$  such that  $N(v) \cap V(G_2) \subseteq N(\tilde{v}) \cap V(G_2)$

The vertex  $\tilde{v}$  (in the third property) will change at each step of our construction. We begin by setting  $\tilde{v} = v'$ , and note that our initial  $G_1$  and  $G_2$  satisfy the conditions above.

Let  $x \in V(\uparrow G)$  be a vertex of minimal level with  $x \notin V(G_1)$ . This property implies that  $N^-(x) \subseteq V(G_1)$  and so  $G_1 \cup \{x\}$  is still a good graph. By Lemma 5, there exists  $y \in V(G_1)$  such that  $N(x) \cap V(G_1) \subseteq N[y] \cap V(G_1)$  and we can define  $\phi(x)$  by taking a clone of  $\phi(y)$  of level greater than  $L(V(G_1) \cup V(G_2))$  such that

$$N^-(\phi(x)) = N(\phi(x)) \cap V(G_2) = \phi(N(x) \cap V(G_1)).$$

This extends the isomorphism, implies that  $G_2 \cup \{\phi(x), v\}$  is still a good graph and that the vertex  $\tilde{v}$  still satisfies the desired property.

We now go in the opposite direction. Let  $z$  be a vertex of minimal level with  $z \notin V(G_2) \cup \{v\}$ : we attempt to define  $\phi^{-1}(z)$ .

We distinguish two cases:

- $zv \notin E(\uparrow G)$ , or  $zv \in E(\uparrow G)$  and  $z\tilde{v} \in E(\uparrow G)$ .

As before, we can apply Lemma 5 to get  $y \in V(G_2) \cup \{v\}$  such that  $N(z) \cap V(G_2) \subseteq N[y] \cap V(G_2)$ . If  $y = v$ , we can instead choose  $y = \tilde{v}$ . We can then define  $\phi^{-1}(z)$  as previously to be a suitable clone of  $\phi^{-1}(y)$ .

- $zv \in E(\uparrow G)$  and  $z\tilde{v} \notin E(\uparrow G)$ .

In this case we will have to change  $\tilde{v}$ , because we want the condition  $N(v) \cap V(G_2) \subseteq N(\tilde{v}) \cap V(G_2)$  to hold after adding  $z$  to  $G_2$ . Let  $w$  be a clone of  $v$  such that  $L(w) > L(V(G_1) \cup V(G_2))$  and  $N^-(w) = (N(v) \cap V(G_2)) \cup \{z\}$ . Such a vertex exists, since  $z$  is a neighbour of  $v$ . The only reason why the subgraph induced by  $V(G_2) \cup \{v, w\}$  might not be a good graph is the edge  $zw$ . We therefore extend the isomorphism to  $G_2 \cup \{z, w\}$ . Since  $G_2$  is a good graph, we can use Lemma 5 as before to first extend the isomorphism to  $z$ . Since, by minimality of  $z$ , the subgraph induced by  $V(G_2) \cup \{z, v\}$  is also a good graph, we can use Lemma 5 again to extend the isomorphism to  $w$ . Finally, the definition of  $w$  implies that  $G_2 \cup \{z, w, v\}$  is a good graph, and we can choose the new  $\tilde{v}$  to be  $w$ , as it satisfies the desired property.

Repeating the argument gives, in the limit, an isomorphism between  $\uparrow G$  and  $(\uparrow G) \setminus v$ . □

## References

- [1] A. Bonato, The infinite locally random graph, preprint, 2005
- [2] A. Bonato and D. Delić, On a problem of Cameron's on inexhaustible graphs, *Combinatorica* **24** (2004), 35–51
- [3] A. Bonato and J. Janssen, Infinite limits of copying models of the web graph, *Internet Mathematics* **1** (2003), 193–213
- [4] A. Bonato and J. Janssen, Infinite limits of the duplication model and graph folding, preprint, 2005

- [5] P.J. Cameron, The random graph, *in* The mathematics of Paul Erdős, II, *Algorithms and Combinatorics* **14**, R.L. Graham and J. Nešetřil, eds, Springer, 1997, 333–351
- [6] M. El-Zahar and N.W. Sauer, Ramsey-type properties of relational structures, *Discrete Math.* **94** (1991), 1–10.
- [7] M. Pouzet, Relations impartibles, *Dissertationes Math. (Rozprawy Mat.)* **193** (1981), 43 pp.