

THE
UNIVERSITY
OF RHODE ISLAND

University of Rhode Island
DigitalCommons@URI

Kinesiology Faculty Publications

Kinesiology

2017

A review of early influences on physical activity and sedentary behaviors of preschool-age children in high-income countries

Ana Cristina Lindsay

Mary L. Greaney

Sherrie F. Wallington

Tatiana Mesa

Carlos F. Salas

Follow this and additional works at: https://digitalcommons.uri.edu/kinesiology_facpubs

**The University of Rhode Island Faculty have made this article openly available.
Please let us know how Open Access to this research benefits you.**

This is a pre-publication author manuscript of the final, published article.

Terms of Use

This article is made available under the terms and conditions applicable towards Open Access Policy Articles, as set forth in our [Terms of Use](#).

1 **A review of early influences on physical activity and sedentary behaviors of preschool-**
2 **age children in high-income countries**

3
4 Ana Cristina Lindsay^{1,2}; Mary L. Greaney³; Sherrie F. Wallington⁴; Tatiana Mesa¹; Carlos F.
5 Salas¹

- 6
7 1. University of Massachusetts Boston, Exercise and Health Sciences Department,
8 College of Nursing and Health Sciences, Boston, MA, USA
9 2. Harvard School of Public Health, Department of Nutrition, Boston, MA, USA
10 University of Rhode Island, Health Studies & Department of Kinesiology, Kingston,
11 RI, USA
12 3. University of Rhode Island, Department of Kinesiology
13 4. Lombardi Comprehensive Cancer Center, Georgetown University Medical Center,
14 Washington, DC, USA
15

16 **Authors' email addresses:**

17 **Ana Cristina Lindsay:** Ana.Lindsay@umb.edu
18 **Mary L. Greaney:** mgreaney@uri.edu
19 **Sherrie F. Wallington:** slw49@georgetown.edu
20 **Tatiana Mesa:** Tatiana.Mesa001@umb.edu
21 **Carlos Salas:** Carlos.Salas001@umb.edu
22
23

24 **Address for Correspondence:**

25 Ana Cristina Lindsay, DDS, MPH, DrPH
26 Associate Professor
27 Department of Exercise and Health Sciences
28 University of Massachusetts Boston
29 100 Morrissey Boulevard
30 Boston, MA 02125
31 Phone: 617-287-7579
32 Email: Ana.Lindsay@umb.edu
33
34
35
36
37
38

39 **Abstract**

40 **Purpose:** Promoting physical activity (PA) is a key component of preventing and controlling
41 childhood obesity. Despite well-documented benefits of PA, globally, rates of physical
42 activity among young children have declined over the past decades, and most children are not
43 accruing sufficient physical activity daily. Helping children develop the foundation for PA
44 habits early in life is critical for the promotion of health in childhood and prevention of
45 chronic diseases later in life, and will ultimately promote longer and healthier lives for
46 individuals and the general population. The purpose of this review is to provide a synthesis of
47 current evidence on influences on PA and sedentary behaviors of preschool-age children in
48 high-income countries.

49 **Design and Methods:** A systematic review of three databases was performed. Studies
50 conducted in high-income countries and published from 2000 onwards that addressed
51 influences on physical activity and sedentary behaviors of preschool-age children were
52 identified and reviewed. Additionally, reference lists of identified articles and relevant
53 published reviews were reviewed. Studies that met the following inclusion criteria were
54 considered: (a) sample included preschoolers (age ≤ 5 years); (b) PA and/or sedentary
55 behaviors or factors associated with PA and/or sedentary behaviors was assessed; (c)
56 published in English; (d) used either quantitative or qualitative methods; and (e) conducted in
57 a high-income country. Data were extracted from selected studies to identify influences on PA
58 and sedentary behaviors of preschool-age children and organized using the social-ecological
59 **model** according to multiple levels of influence.

60 **Results:** Results from included studies identify multiple factors that influence PA and
61 sedentary behaviors of young children in high-income countries at the various levels of the

62 social-ecological model including intrapersonal, interpersonal, environmental, organizational,
63 and policy.

64 **Practice Implications:** Given pediatric nurses' role as primary care providers, and their
65 frequent and continued contact with parents and their children throughout childhood through
66 well-child visits, immunization, and minor acute illnesses, they are well positioned to promote
67 and support the development of early healthful PA habits of children starting in early
68 childhood.

69

70 **Keywords:** physical activity, sedentary behaviors, preschool-age children, social-ecological
71 model

72

73

74

75

76

77

78

79

80

81

82

83

84

85 **INTRODUCTION**

86 Several high-income countries including the United States, United Kingdom,
87 Australia, and Canada, have developed physical activity (PA) guidelines specific to preschool-
88 age children recommending that children this age engage in at least 120 to 180 minutes of
89 total PA (including light PA and moderate-to-vigorous PA) daily (Commonwealth of
90 Australia, Department of Health and Ageing, 2010; Department of Health, Physical Activity,
91 Health Improvement and Protection, 2011; NASPE, 2009; Okely, Salmon, Trost, & Hinkley,
92 2008; Tremblay et al., 2012).

93 In the United States, the National Association for Sport and Physical Education
94 (NASPE, 2009) has developed PA guidelines from infancy to the age of 5 years. These
95 guidelines advocate increasing opportunities for preschoolers to engage in a minimum of 120
96 minutes of daily PA in the form of 60 minutes of structured activity and 60 minutes of
97 unstructured or spontaneous active play. **Similarly, the American Academy of Pediatrics**
98 **recommends that clinicians encourage parents to increase physical activities and decrease time**
99 **spent in sedentary activities (e.g., screen-time, time spent indoors, etc.) in a manner**
100 **compatible with the developmental level of a child (AAP, 2006).**

101 Furthermore, high-income countries including the United States, Canada and Australia
102 have created stringent screen-viewing guidelines in response to the increasing rates of
103 sedentary behaviors among preschool-aged children. For example, the American Academy of
104 Pediatrics recommends that screen-viewing be limited to a maximum of two hours per day for
105 children over two years of age (AAP, 2006). Both Canada (Tremblay et al., 2012) and
106 Australia (Commonwealth of Australia, Department of Health and Ageing, 2010) have more
107 stringent screen-viewing guidelines for young children, with both countries recommending

108 that screen-viewing for children 1-4 years of age in Canada and 2-5 years of age in Australia
109 be limited to 1 hour per day.

110 Physical activity is a key component of energy balance consistent with healthy
111 patterns of physical growth and weight status; thus, promoting PA is essential to prevention of
112 childhood obesity (O'Dwyer et al., 2012; Pratt, Epping, & Dietz, 2009). Physically active
113 children have healthier cardiovascular profiles, leaner body frames, and higher peak bone
114 mass compared to physically inactive children (AAP, 2006; Goldfield, Harvey, Grattan, &
115 Adamo, 2012; Hodges, Smith, Tidwell, & Berry, 2013; O'Dwyer, Fairclough, Knowles, &
116 Stratton, 2012). In addition to regulating body weight and improving body composition, PA
117 improves psychological and social wellbeing (AAP, 2006; Beets et al., 2011; Hodges et al.,
118 2013; O'Dwyer et al., 2012).

119 Despite well-documented benefits of PA for children, activity levels of children across
120 the globe have declined over the past decades, and most preschool-age children are not
121 accruing the recommended levels of moderate-vigorous physical activity (MVPA) daily
122 (Montgomery et al., 2004; Pate et al., 2015). Moreover, an increasing number of studies
123 document excessive screen-viewing time including television, computers, smartphones, etc.
124 among preschool-age children (Asplund, Kair, Arain, Cervantes, Oreskovic, Zuckerman,
125 2015; Beets, Bornstein, Dowda, & Pate, 2011; Davison et al., 2011; Ekelund, Brage, &
126 Wareham, 2004; Garriguet, Carson, Colley, Janssen, Timmons & Tremblay, 2016; Lampard,
127 Jurkowski, Davison, 2013; Mendoza, Zimmerman, & Christakis, 2007; Pate et al., 2015; Vale,
128 Santos, Soares-Miranda, Silva, & Mota, 2010; Vandewater, Bickham, & Lee, 2006).

129 Physical activity and sedentary behaviors are complex and influenced by interacting
130 multi-level factors that either facilitate or hinder PA and sedentary behaviors in young

131 children. The social-ecological model has been extensively used to help health professionals
132 and researchers identify and understand how the various multi-level influences interact to
133 form individual's PA **opportunities and** choices. Consistent evidence shows that addressing
134 multiple levels of the social-ecological model is associated with greater change in behaviors
135 including PA (Sallis, Cervero, Asher, Henderson, Kraft & Kerr, 2006). Moreover, the social-
136 ecological model recognizes **the context and interaction typical of preschool-aged children**
137 **and provide an important framework** for developing interventions that address the social and
138 physical environments and public policy for improving PA and reducing sedentary behaviors
139 of young children (Sallis, Cervero, Asher, Henderson, Kraft & Kerr, 2006).

140 Given the increasing prevalence of insufficient PA and increasing levels of sedentary
141 behaviors among young children, this review sought to elucidate early influences on PA and
142 sedentary behaviors of preschool-age children in high-income countries by using the social-
143 ecological model to identify and organize facilitators and/or barriers to PA and sedentary
144 behaviors of young children.

145 **METHODS**

146 We searched three electronic databases (PubMed, SPORTDiscus, and PsycINFO) to
147 identify studies published from January 2000 onward that addressed influences on PA and
148 sedentary behaviors among preschool-age children (age ≤ 5 years) in high-income countries.
149 We used the criteria from the Organization for Economic Cooperation and Development
150 (OECD) and The World Bank to define high-income countries, which defines high-income as
151 a country with a gross national income per capita above US\$ 12,475 in 2015, calculated using
152 the Atlas method (The World Bank, 2013). Additionally, we searched the references of
153 identified studies and relevant published reviews. We used the combination of the following

154 key search terms: *physical activity, inactivity, lifestyle, sedentary behavior, screen, preschool,*
155 *children, overweight, obese/obesity, prevention, and influences.* Studies that met the following
156 inclusion criteria were reviewed: (a) sample included preschoolers (age ≤ 5 years); (b)
157 measured PA and sedentary behaviors and/or factors associated with PA and sedentary
158 behaviors; (c) published in English; (d) used either quantitative or qualitative methods; (e)
159 **conducted in a high-income country.** Exclusion criteria included the following: (a) the study
160 sample was children with preexisting conditions (e.g., cardiac disease, hypertension, diabetes
161 mellitus, dyslipidemia, or mental illness) that could constrain PA; (b) review articles; and (c)
162 articles published only in languages other than English. One author read all the abstracts of
163 identified full-text papers meeting inclusion criteria. Identified papers were reviewed by two
164 of the authors, who used the social-ecological model as a framework to organize identified
165 factors influencing PA. Figure 1 shows the literature search strategy.

166 [Figure 1 here]

167 **Theoretical Model for Examining Influences on Preschool Children’s Physical Activity**

168
169 The social-ecological model provided the conceptual framework for this review
170 (McLeroy, Bibeau, Steckler, & Glanz, 1988) and was used to organize influences on
171 preschool-age children’s PA by level: (a) intrapersonal (**e.g., age, gender, material**
172 **circumstances, ethnicity, etc.**); (b) interpersonal (e.g., social support, networks, etc.); (c)
173 environmental (e.g., access and proximity to parks, etc.); (d) organizational (e.g., child care,
174 federally funded nutrition programs such as WIC, etc.); and (e) policy (e.g., state policies and
175 regulations related to nutrition and PA in child care settings, etc.). This framework posits that
176 factors at each level interact and influence health behaviors (McLeroy et al., 1998). Table 1
177 shows a breakdown of the studies reviewed according to the social-ecological model.

178

[Figure 2 and Table 1 here]

179 **RESULTS**

180 **Individual/Intrapersonal Influences**

181 **Socioeconomic and demographic factors.** Parents' socioeconomic status (SES) and
182 educational level are associated with preschool children's early PA behaviors (Dawson-Hahn,
183 Fesinmeyer, & Mendoza, 2015; Salmon, Owen, Crawford, Bauman, & Sallis, 2003). Jones,
184 Hendricks, and Draper (2014) found that among children aged 4–5 years old attending
185 preschools, parents with low-income reported that their children spent 93% of time indoors
186 compared to 79% reported by parents with mid/high-income. Parental socioeconomic status is
187 also associated with preschool children's early sedentary behaviors. Levin, Martin & Riner
188 (2004) found excessive TV viewing habits among a sample of low-income 4-year old children
189 enrolled in Head Start in South Carolina.

190 Differences in PA by sex exist (Vale et al., 2010; Van Cauwenberghe et al., 2012), and
191 across SES and demographic strata (Vale et al., 2010; Van Cauwenberghe, Jones, Hinkley,
192 Crawford, & Okely, 2012), with boys being more active than girls (Finn, Johannsen, &
193 Specker, 2002; Pate, Pfeiffer, Trost, Ziegler, & Dowda, 2004).

194 Children's PA and sedentary behaviors may also vary according to family structure.
195 Controlling for household SES and child age, Bagley, Salmon, and Crawford (2006)
196 determined that boys without siblings spent more time watching TV compared to those who
197 had siblings. They also found that girls from single-parent families spent significantly more
198 time watching TV compared to girls from two-parent families, and girls with siblings were
199 more physically active compared to those who were the only child.

200 **Ethnicity, acculturation, and place of birth.** Caucasian children have higher rates of
201 PA than African American or Hispanic children (Ariza, Chen, Binns, & Christoffel, 2004;
202 Fakhouri, Hughes, Brody, Kit, & Ogden, 2013; Kuepper-Nybelen et al., 2005; Kumanyika &
203 Grier, 2006).

204 A few studies have examined the associations between ethnicity and preschool
205 children's PA (Dawson-Hahn, Fesinmeyer & Mendoza, 2015; Fitzgibbon et al., 2011;
206 Fitzgibbon et al., 2006; O'Connor et al., 2014; Toselli, Zaccagni, Celenza, Albertini &
207 Gualdi-Russo, 2015). Dawson-Hahn, Fesinmeyer & Mendoza (2015) found that the majority
208 of Latino preschool-aged children attending Head Start programs exceeded US national and
209 international guidelines of physical activity duration. A study conducted in Italy among
210 parents of Italian and immigrant preschool-aged children found that physical activity was
211 significantly higher in Italians than in immigrants (Toselli et al., 2015).

212 Limited research exists examining the association between acculturation levels and
213 preschool-age children's PA, and the limited research shows mixed results (Soltero, Cerin,
214 Lee & O'Connor, 2016; O'Connor et al., 2014; Gallagher, 2010). A recent study with 240 of
215 preschool-aged children showed that Latino parents' acculturation levels moderated the
216 relationship between perceptions of disorder and crime, which in turn, influenced parenting
217 practices that discourage child physical activity due to safety concerns were associated with
218 increased perceptions of traffic hazards, physical and social disorder and perceived stranger
219 danger (Soltero, Cerin, Lee & O'Connor, 2016). Another study with also conducted with
220 Latino parents of preschool-aged children showed that cultural variables only had a weak
221 main effect on PA parenting practices, specifically on discouraging PA due to safety concerns
222 (O'Connor et al., 2014). A qualitative study with Mexican-American mothers showed that

223 mothers' acculturation levels influenced their views of the type of physical activity children
224 should engage in depended on the age and maturity of the child (Gallagher, 2010). Previous
225 research highlights a need for further study of associations between acculturation and
226 preschoolers' PA in Western cultures (O'Connor et al., 2014; Suen, Cerin, & Wua, 2015).

227 Cespedes, McDonald, Haines, Bottino, and Taveras (2013) examined obesity-related
228 behaviors in urban, low-income, and non-U.S.- and U.S.-born racial/ethnic minority
229 preschool-age children (34% Black, 52% Hispanic), and found that time spent in active play
230 was lower among children whose parents were born outside the U.S. than among those whose
231 parents were U.S.-born, after adjusting for parental education.

232 **Interpersonal Influences**

233 **Parental physical activity.** Parental PA and sedentary behaviors are important
234 determinants of their preschool-aged children's PA and sedentary behaviors. Ruiz, Gesell,
235 Buchowski, Lambert, and Barkin (2011) determined that Latino parents who are less
236 physically active and more sedentary had children who were more sedentary and less active
237 than Latino parents who were more physically active. A similar study found weak correlations
238 between mild and moderate parental PA and their 3- and 4-year-old children's PA (Taylor
239 et al., 2009). A study conducted in Canada by Carson, Stearns, and Janssen (2015) examined
240 the associations between parental PA and screen time behaviors and their young children's
241 behavior (61% aged 1–3, 35% aged 4–5) and found that parents in the lowest quartile
242 of PA were 2.77 times more likely to have a child in the lowest quartile of PA compared
243 with parents in the highest quartile. Relationships were stronger in two parent homes than in
244 single-parent homes. Furthermore, parents in the second, third, and fourth screen time
245 quartiles were significantly more likely to have a child in the highest quartile of screen time

246 compared with parents in quartile one. Similar, a recent study conducted in Australia found
247 that maternal self-reported co-participation in sedentary behavior and provision of child
248 opportunities for physical activity was associated with children's physical activity (Hnatiuk,
249 Ridgers, salmon & Hesketh, 2016).

250 **Parental beliefs and attitudes.** Parents' perceptions, beliefs, and attitudes toward PA
251 may influence their young children's PA behaviors (Dwyer, Higgs, Hardy, & Baur, 2008;
252 Hesketh et al., 2013; Hinkley, Salmon, Okely, Crawford, & Hesketh, 2011; Loprinzi & Trost,
253 2010; O'Connor, Chen, Baranowski, Thompson, & Baranowski, 2013; Zecevic, Tremblay,
254 Lovsin, & Michel, 2010). Parental belief that participating in PA is important is associated
255 with their children's participation in both organized and free-time PA (Sawyer et al., 2014).
256 Children whose parents hold positive attitudes toward PA (e.g., PA is important to overall
257 health) are more active compared to those whose parents do not hold these positive attitudes
258 (Sawyer et al., 2014). Similarly, Zecevic and colleagues (2010) determined that children of
259 parents who view PA as enjoyable engage in significantly more PA than children did whose
260 parents did not view PA positively.

261 **Parental concerns.** Parents of preschool-age children have reported that their
262 concerns about safety inhibit their children's PA (Dwyer et al., 2008; O'Connor, Chen, et al.,
263 2013; Soltero et al., 20016; Suen et al., 2015), including concerns related to neighborhood and
264 community safety (e.g., crime, traffic) (Dwyer et al., 2008; Soltero et al., 2016). In addition,
265 parental concerns about excess screen time are also associated with children's PA levels (He,
266 Irwin, Sangster Bouck, Tucker, & Pollett, 2005; De Decker et al., 2012).

267 **Parenting styles.** Parenting styles are psychological constructs that represent broad
268 and standard strategies that parents use in child rearing (Baumrind, 1971). Overall parenting

269 style encompasses broader patterns of how parents respond and demand to their children
270 (Baumrind, 1971). Four parenting styles have been defined: 1) authoritarian (demand
271 obedience); 2) authoritative (use reasoning); 3) permissive (acquiesce to child's demands) and
272 4) uninvolved. Only one study included in this review examined the association between
273 parenting styles and preschool-age children's PA levels, and no association was found
274 between parenting styles and time spent in active play nor did parenting styles moderate the
275 relationship between parental support and child active play (Schary, Cardinal, Loprinzi,
276 2012).

277 **Parenting Practices.** Parenting practices describe context-specific behaviors such as
278 what a parent does to facilitate physical activity. A growing, but still limited literature exists
279 on studies that have examined the influence of parenting practices on young children's PA
280 behaviors (Hesketh et al., 2013; O'Connor, Chen, et al., 2013; Oliver, Schofield, & Schluter,
281 2010; Zecevic, Tremblay, Lovsin and Michel, 2010; Dowda et al., 2011; Hesketh et al., 2014).
282 Results of these studies show that some parental practices, such as encouraging PA, setting
283 rules, providing transportation, **paying fees and tuition**, parental modeling of PA, and
284 engaging in PA activities with children are associated with PA behaviors of preschool-age
285 children (Hesketh et al., 2014; O'Connor, Chen, et al., 2013; Vanderwater et al., 2005; Veitch,
286 Hume, Salmon, Crawford, & Ball, 2013).

287 An important parenting practice is parental support for physical activity. Available
288 research shows that children are more likely to be active if parents are supportive of them
289 being physically active in a number ways, but especially by providing encouragement,
290 participating in PA together, taking children to places where they can be physically active,
291 and enrolling them in organized activities such as sports classes (Davison et al., 2011;

292 Zecevic, Tremblay, Lovsin and Michel, 2010). Nevertheless, only limited research has
293 examined the relationship between parental support of PA and preschool-age children's PA
294 levels (Dowda et al., 2011; Gubbles et al., 2011; Hinkley, Salmon, Okely, Crawford, 2013;
295 O'Connor et al., 2013; Scharby et al., 2012; Vanderwater et al., 2005; Zecevic et al., 2010).
296 O'Connor et al. (2013) examined parental encouragement and discouragement of PA among
297 Latino children aged 3-5 and determined that parental practices promoting PA, including
298 enrolling children in sports, participating in the child's activities, and modeling of PA, and
299 supporting children's PA behavior. Grigsby-Toussaint, Chi, and Fiese (2011) found that while
300 certain environmental factors, such as a built environment with high levels of green space are
301 important for PA, most increases in preschoolers' PA is due to parental support for PA. A
302 study with preschool-age boys determined that boys who received greater parental support for
303 PA were significantly more likely to engage in 1 hour or more of daily PA than those who did
304 not receive such support (Zecevic et al., 2010). A study by Østbye and colleagues (2013)
305 found that parental attitudes in support of PA were significantly associated with MVPA
306 among preschool-age children. Suen and colleagues (2015) examined parental practices
307 among Hong Kong preschoolers and determined that providing conditional, instrumental, and
308 motivational support to parents motivated them to encourage children to be physically active.
309 On the other hand, parental emphasis on academic achievement, lack of time and resources,
310 promotion of sedentary behaviors, and safety concerns discouraged PA (Suen et al., 2015).

311 Parental involvement is another influential parental practice as it relates to PA. A
312 study conducted in New Zealand by Oliver et al. (2010) found a positive association between
313 PA of parents and their preschool-aged children, which suggests the potential importance of
314 parental involvement in preschool-based PA intervention such as parents and children

315 participating together in activities sponsored by the intervention (e.g., Family Fun Nights) and
316 parents direct involvement in intervention activities beyond the intervention environment (e.g.
317 “try this at home”).

318 **Parental and/or family influences on television-viewing and other screen-viewing.**

319 Most studies focusing on parental and family influences on screen-viewing time of young
320 children have focused on TV (Bagley, Salmon & Crawford, 2006; Barr, Danziger, Hilliard,
321 Andolina, Ruskis, 2010; Certain & Kahn, 2002; Dalzell, Msall, High, 2000; Dawson-Hahn et
322 al., 2015; Downing, Hinkley, Hesketh, 2015; Dennison BA, Erb TA, Jenkins PL., 2002;
323 Jackson, Djafarian, Stewart, & Speakman, 2009; Thompson,
324 Polk, Cheah, Vandewater, Johnson, Chrismer, Tschann, 2015; Vandewater, Rideout, Wartella,
325 Huang, Lee, Shim, 2007).

326 A study by Jackson et al. (2009) of preschool-age children (2–6 years) found that
327 children who watched more TV were significantly less physically active than children who
328 watched less TV. Similarly, a recent cross-sectional study by Dawson-Hahn et al. (2015)
329 conducted among preschool-age children in found that watching TV was inversely associated
330 with PA. Parental attitudes, screen time, and having a television in the bedroom were positive
331 predictors of children’s excess screen time and inadequate PA.

332 Parental education and self-efficacy for PA were negative predictors of screen time
333 (Carson & Janssen, 2012). A study by Lampard et al (2013) found low-income preschool-age
334 children (2-5 years) were more likely to meet the American Academy of
335 Pediatrics screen time recommendation (no more than 1 hour per day) if their parent reported
336 high restriction of child screen time. Moreover, in multivariate analysis, less
337 parent screen time, fewer parent life pressures, and greater social support were associated with

338 parents' high restriction of screen time (Lampard et al., 2013). Downing, Hinkley and Hesketh
339 (2015) found that children whose parents limited television viewing spent significantly less
340 time in watching TV and in total screen time; however, overall sedentary behavior was
341 unaffected. Further studies need to be conducted on how family environment influences
342 sedentary behaviors among children.

343 Studies suggest that the TV viewing habits of parents and other family members in the
344 household (e.g., older siblings) likely contribute to the time preschoolers spend watching TV
345 (Downing, Hinkley, Hesketh, 2015; Djafarian, Stewart, & Speakman, 2009;
346 Thompson, Polk, Cheah, Vandewater, Johnson, Chrismer, Tschann, 2015). The more time that
347 parents spend watching TV the more time their preschool-aged children spend watching TV
348 (Vandewater, Rideout, Wartella, Huang, Lee, Shim, 2007). A similar pattern is found between
349 older siblings' time spent viewing TV and preschool children's time spent viewing
350 (Vandewater, Rideout, Wartella, Huang, Lee, Shim, 2007).

351 Current research also indicates that many preschoolers' daily screen time exceeds
352 recommendations. Vandewater, Shim, and Caplovitz (2004) determined that preschool
353 children watched more television than primary school children (2.19 hours/day vs. 1.91
354 hours/day), with 61.7% of children in the preschool group watching television for at least
355 2 hours per day.

356 Other screen-viewing behaviors common among young children include activities
357 such as DVDs/VHS, video games, computers, smartphones, etc (Bagley, Salmon & Crawford,
358 2006), but to date, these other screen-viewing behaviors have not been extensively examined
359 among preschool-age children.

360 **Siblings and peer influences.** Sibling and peer PA and sedentary behaviors appear to
361 be important influences on preschool-aged children’s PA and sedentary behaviors.
362 Preschoolers observe and imitate the behaviors of those who are similar to them (Ward et al.,
363 2017). Therefore, siblings and peers may be role models for preschoolers' physical activity. A
364 study conducted in child care centers in Canada found that peers influenced preschoolers’
365 physical activity over time (Ward et al., 2017).

366 **Environmental (Community/Neighborhood) Influences**

367 Promoting physical activities in neighborhood environments where children spend
368 significant amounts of time can contribute to increased PA levels (Goldfield et al., 2012).
369 Several studies show that being outdoors is the strongest correlate of PA among preschool
370 children and that activity levels correlate with the number of play spaces near their homes and
371 the amount of time spent in those spaces (Hart, Herriot, Bishop, & Truby, 2003; Lindsay,
372 Sussner, Greaney, & Peterson, 2009; McKenzie et al., 2008; Roemmich et al., 2006; Salmon
373 et al., 2013). Researching the relationships between built environments and PA is challenging,
374 and designing and implementing supportive environments, corrective programs and policies is
375 complex because environmental factors may vary across children of different demographics
376 (e.g., age, gender, race/ethnicity, SES) and cultural backgrounds (Vandewater et al., 2006).

377 **Weather and season.** Natural environments can present barriers to PA. McKee,
378 Murtagh, Boreham, Nevill, and Murphy (2012) examined the influence of season on
379 objectively assessed PA in preschool children in Minnesota and found that children take
380 approximately 2,000 (20%) fewer steps per day in winter than in Spring. A qualitative study
381 by Lindsay et al. (2009) determined that Latina mothers viewed weather as an important factor
382 influencing their preschool children’s PA habits in Massachusetts; during cold weather,

383 children spent more time indoors and engaged in less PA, whereas children spent more time
384 outdoors and in parks and recreational facilities during warmer weather. Similarly, additional
385 qualitative studies with parents of young children in Canada and Australia showed that colder
386 weather posed challenges for parents in keeping their preschoolers physically active (He et al.,
387 2005; Pearson, Salmon, Crawford, Campbell, & Timperio, 2011).

388 Weather is also an important influence on children's screen time. A qualitative study
389 conducted in six European countries by De Decker and colleagues (2012) found that weather
390 condition was one of the most important factors influencing children's screen time.
391 Researchers suggest that parents should be provided with guidance on alternatives for screen
392 activities and information on how to set rules for screen time to assist them in decreasing
393 their preschool children's screen time.

394 **Time outdoors.** The more time preschool children spend outdoors, the higher their PA
395 levels (Boldemann et al., 2006; Burdette, Whitaker, & Daniels, 2004; Hinkley, Crawford,
396 Salmon, Okely, & Hesketh, 2008). Several studies included in this review suggest that parents
397 and caregivers can and should encourage outdoor play (Anderson et al., 2008; Ergler, Kearns,
398 & Witten, 2013; Tandon, Saelens, Zhou, Kerr, & Christakis, 2013; Veitch, Salmon, & Ball,
399 2010). Questions of safety and accessibility, however, can make it more difficult for some
400 parents and children to spend time outdoors. Minority and low-income parents, for example,
401 are more likely to live in communities with fewer parks, sports facilities, bike paths, and other
402 places for children to be active and safe (Lindsay et al., 2009).

403 **Availability and access to PA programs, parks, and recreational facilities.**

404 Availability and access to PA programs, parks, and recreational facilities are important
405 influences on preschool children's PA. Results from studies reviewed suggest that to support

406 efforts that promote preschool-age children's PA activity through active play, age-appropriate,
407 outdoor play spaces with access to play equipment should be developed and maintained in
408 communities (Anderson et al., 2008; Burdette & Whitaker, 2005; Ergler, Kearns, & Witten,
409 2013; Tandon, Saelens, Zhou, Kerr, & Christakis, 2013; Veitch, Salmon, & Ball, 2010). The
410 concept of neighborhood greenness has also been correlated with preschool-age children and
411 PA. Grigsby-Toussaint et al. (2011) determined that families with preschool-age children who
412 have access to recreational facilities with higher levels of green space are more physically
413 active than are preschoolers without access.

414 **Neighborhood safety.** As discussed earlier, a number of studies have demonstrated
415 that perceived lack of neighborhood safety is a potential barrier to preschool children's PA
416 (Burdette & Whitaker, 2005; Goldfield et al., 2012; Lindsay et al., 2009; Salmon et al., 2003).
417 For example, a cross-sectional survey of 2,445 mothers of 2-3 year old children found that
418 perceived neighborhood safety was a barrier to PA (Burdette, Wadden, & Whitaker, 2006).
419 Likewise, a qualitative study conducted with Latina mothers in Massachusetts found that
420 mothers reported neighborhood safety as a barrier to PA engagement and to their preschool-
421 age children's active play (Lindsay et al., 2009). On the other hand, research indicates that
422 providing a safe play area and attendants to supervise children increases PA and decreases
423 screen time of young children attending schools near the play areas (Burdette et al., 2006).

424 **Organizational Influences**

425 **Early care and education (ECE) settings.** Currently, there are several types of early
426 care and education arrangements in the United States. Broadly, they can be broken down into
427 four types: **nurseries/preschools**, center-based child care/daycare, family child care homes,
428 and home-based care from nannies/babysitters. Center-based child care is generally provided

429 in a public building and children are usually grouped by age in classrooms with at least one
430 trained teacher. Most states regulate center-based childcare. Nursery schools and pre-schools
431 are educational establishments or learning spaces offering early childhood education to
432 children between the ages of 3 and 5 years. Like center-based child care, most states regulate
433 nurseries and pre-schools. Family child care is home-based, and providers care for children
434 other than their own in the providers' own home. Family child care may be licensed or
435 unlicensed, and despite the growing interest in this type of setting, further exploration of PA
436 and sedentary in family child care settings is needed because available studies show that PA
437 levels have been low among preschool-age children attending licensed family child care
438 homes (Lindsay et al., 2016; Vanderloo, Martynuik, & Tucker, 2015). Home-based care
439 provided by a nanny or a babysitter often involve child care hired on a scheduled full- or part-
440 time basis with care often provided in the child's home.

441 Results of studies included in this review suggest that day care centers and preschools
442 may be able to provide access to outdoor play spaces for young children (Burdette &
443 Whitaker, 2005; Grigsby-Toussaint et al., 2011), which could foster opportunities to be
444 physically active. A recent study conducted in the U.K. by Hesketh, Griffin & van Sluijs
445 (2015) found that preschool-aged children and particularly boys were less sedentary and more
446 active when in child care compared to at home. Despite opportunities for daycare/child care
447 centers and preschools to provide access to outdoor play spaces for young children some
448 studies have also documented children facing barriers to being physically active in these
449 settings. Using focus groups, Copeland, Kendeigh, Saelens, Kalkwarf, and Sherman (2012)
450 determined that preschool and daycare center teachers believe that PA is important for
451 developing children, but noted that children's inappropriate clothing (e.g., flip-flops or

452 sandals, dress/expensive clothes, no hat/gloves or coat during the winter) were barriers to
453 children's PA at the daycare centers (Copeland et al., 2012). Furthermore, results showed that
454 clothing choices were a source of conflict between parents and child-care providers (Copeland
455 et al., 2012). In addition, a recent cross-sectional study conducted in Australia by Hinkley,
456 Salmon, Crawford, Okely & Hesketh (2016) found that preschool-age children significantly less
457 active during the hours they spent in organized child care than outside care hours.

458 Quality of **early education and care** also appears to influence children's PA levels
459 when in these settings. Dowda, Pate, Trost, Almeida, and Sirard (2004) found that children
460 spent more time in sedentary activities at low-quality preschools than did children in high-
461 quality preschools.

462 Other important influences on levels of PA among preschool-age children in ECE
463 settings include adult support and availability of both outdoor space and play equipment.
464 Using direct observation at child care centers, Bower et al. (2008) determined that adult
465 support and availability of play equipment were associated with greater PA and lower
466 sedentary activity levels among preschoolers. Similarly, a study in the Netherlands focusing
467 on preschools and daycare centers found that locations with greater outdoor space and
468 equipment availability reported higher PA levels in the children (Gubbels, Van Kann, &
469 Jansen, 2012). Likewise a recent study by Schlechter, Rosenkranz, Fees & Dzewaltowski
470 (2017) found that providing more time outdoors and restructuring preschool activities from
471 whole group to small group could increase the amount of total physical activity that children
472 accumulate during preschool.

473 **Policy Influences**

474 **Need for PA policies in early care and education settings.** There has been a growing
475 interest in understanding how policies in ECE settings influence PA behavior and obesity

476 prevention efforts in young children in the U.S. (Larson, Ward, Neelon, & Story, 2011).
477 Some studies investigating the role of child care environments in influencing PA levels of
478 preschoolers noted these settings as important venues to promote and support PA among
479 young children, given families' increased reliance on these sites (Gubbels, Van Kann, &
480 Jansen, et al., 2012; Vanderloo et al., 2014). National trends in the U.S show that about 77%
481 of children age 3–5 years spend an average of 30 hours weekly in an ECE setting (Buscemi,
482 Kanwischer, Becker, Ward, & Fitzgibbon, 2015; McPherson & Homer, 2011). A study by
483 Duffey, Slining, and Benjamin Neelon (2014) investigating ECE policies found that no state
484 in the US had regulations for staff joining children in PA, taking away PA opportunities as a
485 punishment for poor classroom behavior (e.g., taking away recess), or providing
486 training/education on PA for childcare providers. Study results suggested that there is room
487 for improvement in childcare regulations related to PA for young children and that updated
488 regulations are needed (Duffey et al., 2014).

489 **National support for PA policies.** Some of the leading national health organizations
490 have come to consensus on strategies to support obesity prevention efforts through promotion
491 of PA, reduced screen time, and healthy eating in ECE settings (AAP, American Public Health
492 Association, [National Association of Pediatric Nurse Practitioners \(NAPNAP\)](#), and National
493 Resource Center for Health and Safety in Child Care and Early Education, 2012). With
494 support from the Health Resources and Services Administration, organizations such as the
495 American Academy of Pediatrics, [NAPNAP](#), Maternal Child Health Bureau, American Public
496 Health Association, and the National Resource Center for Health and Safety in Child Care and
497 Early Education have outlined national child care regulations that include PA guidelines for
498 children from birth to 6 years of age (AAP et al., 2012) attending ECE setting. These

499 guidelines recommend that preschool-age children engage in 90–120 minutes of age-
500 appropriate MVPA per 8-hour day in an ECE setting. If weather permits, preschool-age
501 children should be given two or three occasions of 60–90 minutes of outdoor play, and that
502 structured activities that promote bodily movement should be led by caregiver/teachers two or
503 more times per day (indoor or outdoor). Another recommendation is to have written policies
504 in ECE settings regarding children’s PA while at the ECE. The Institute of Medicine (IOM,
505 2011) has made policy and policy implementation recommendations for ECE settings, such as
506 the need to increase young children’s PA and reduce sedentary behavior, in order to guide
507 care providers and health professionals. The Society of Behavioral Medicine has
508 recommended that state and local policymakers use effective evidence-based models (e.g.,
509 Michelle Obama’s Let’s Move!, Childcare) to implement policies in ECE settings in order to
510 increase PA and reduce sedentary behavior (Buscemi et al., 2015). A recent study in the
511 United States suggested adoption of the Montessori school system as a strategy to promote PA
512 in preschools. Pate et al. (2015) determined that compared to students in traditional
513 preschools, children in Montessori preschools accumulated more light, MVPA, and total PA
514 after adjusting for BMI, sex, SES, and parental education (Pate et al., 2015). Different than
515 traditional school settings where children often sit at their desks for most of the time, children
516 in Montessori programs learn through action and self-discovery, choosing activities and
517 moving about freely during the course of the day (Pate et al., 2015).

518 In addition, it has been proposed that community policies concerning joint-use
519 agreements related to PA (e.g., school gyms), neighborhood design, and urban planning may
520 increase PA opportunities for preschool-age children (McPherson & Homer, 2011).

521 Research suggests that state regulations, outlined in the National Resource Center for
522 Health and Safety in Child Care and Early Education (2011), lack specificity about PA
523 frequency in ECE settings (Battista et al., 2014; Benjamin, Cradock, Walker, Slining, &
524 Gillman, 2008; Cradock, O'Donnell, Benjamin, Walker, & Slining, 2010). This is an area of
525 growing research interest, and additional information is critical to providing directions on
526 ways in which state regulations can help promote early physical activity among young
527 children.

528 **ECE policy change efforts.** ECE-level policies and efforts have been supportive
529 influences on preschool-age children's PA behaviors in the U.S. (Gubbels, Slining, et al.,
530 2012; Trost, Ward, & Senso, 2010), but states inconsistently regulate the implementation of
531 national policies on PA in ECE settings (Buscemi et al., 2015; Gubbels et al., 2012; Larson et
532 al., 2011; McPherson & Homer, 2011; Vanderloo et al., 2014), and many children do not meet
533 PA levels in childcare centers (Buscemi et al., 2015; Duffey et al., 2014; Vanderloo et al.,
534 2014). There have been attempts to develop and implement PA policies in ECEs. For
535 example, in 2007, New York City's ECE centers implemented new regulations put forth by
536 the New York City Department of Health and Mental Hygiene to reduce screen time, increase
537 PA, and provide healthier beverages (Nonas, Silver, Kettel Khan, & Leviton, 2004). However,
538 results of recent studies evaluating compliance with these recommendations found that
539 compliance was low for offered PA time (38.5%) and structured PA time (34.6%) (Lessard
540 et al., 2014; Nonas et al., 2004).

541 **Barriers to PA policy changes in ECE settings.** Some of the barriers to promoting
542 PA in ECE settings that influence adoption of PA regulations are inadequate outdoor/play
543 space, sedentary staff, staff not interested in PA and lack of appropriate equipment (Buscemi

544 et al., 2015; Nonas et al., 2004). Research gaps in preschool-age children's PA measurements
545 and PA program policy-assessment tools are areas to further explore in ECE sites (Kaciroti,
546 Staples-Watson, & Lumeng, 2012).

547 **DISCUSSION**

548 Using the social-ecological model as a framework, we appraised the current literature
549 to look at the context of and influences on PA for preschool-age children. Our review
550 provides significant evidence for several levels of action to improve the current state of PA for
551 this diverse population. Overall, the two important themes are the importance of increasing
552 access and achieving adult buy-in, including that of parents and child care providers. While
553 policies have focused primarily on the quantity of suggested PA, these twin themes, access to
554 and adult support of PA, could inform future policies and potentially have a positive impact
555 on this important issue.

556 Findings from studies included in this review elucidated multi-level factors at various
557 levels in the social-ecological model that influence preschool-age children's PA levels and
558 behaviors. At the individual level, factors associated with differing levels of PA include SES
559 and the child's sex (Annesi, Smith, & Tennant, 2013; Bagley et al., 2006; Cespedes et al.,
560 2013; Chuang, Sharman, Skala, & Evans, 2013; Kumanyika & Grier, 2006; Montgomery et
561 al., 2004; O'Connor et al., 2014; Pate et al., 2004; Salmon et al., 2003; Sisson et al., 2009;
562 Suen et al., 2015; Vale et al., 2010; Van Cauwenberghe et al., 2012). At the individual level,
563 some factors suggest that the lack of PA may have to do with lack of access or insufficient
564 family resources to make PA a priority (e.g., SES, family structure), while other factors
565 suggest parental values come into play (child's sex and cultural context).

566 At the interpersonal level, the presence of siblings, single- versus two-parent family
567 structure, and the parents' culture, parents' PA habits, attitudes toward PA, concerns about the
568 outside environment and their encouragement of PA—are important **factors influencing their**
569 **children's PA** (Carson et al., 2014; Davison & Birch, 2002; De Decker et al., 2012; Oliver et
570 al., 2010; Østbye et al., 2013; Pfeiffer, Dowda, McIver, & Pate, 2009; Rodriguez-Oliveros et
571 al., 2011). Family time spent watching TV is also an important interpersonal level factor
572 impacting the PA level of young children (Dawson-Hahn et al., 2015; Jackson et al., 2009). In
573 addition, studies suggest that parents can be important mediators of children's sedentary
574 behaviors and should promote other enjoyable alternatives to increase children's level of PA
575 (Salmon et al., 2005). At the interpersonal level, family factors that suggest a need for access
576 include concerns about neighborhood safety and traffic, whereas parents' PA habits and
577 perceptions suggest a need for intervention that address communication, education and
578 provision of guidance for parents regarding the importance of these factors in influencing their
579 children's PA.

580 At the environmental level, outdoor space—greenness of the environment, weather
581 and season, access to parks and PA programming, and safety and traffic—was the primary
582 factor (Boldemann et al., 2006; Burdette et al., 2004; Ergler et al., 2013; Grigsby-Toussaint et
583 al., 2011; McKee et al., 2012; McKenzie et al., 2008; Salmon et al., 2013; Tandon et al., 2013;
584 Veitch et al., 2010). At the environmental level, identified factors reflect a need for access to
585 safe and age-appropriate places for young children to be active (safety and traffic, weather and
586 seasons). The fact that “greenness” of the outdoor resources has an impact on PA suggests
587 that making the outdoor space desirable as well as useable is important. At the institutional

588 level, not all EEC facilities have equal resources, and, access is an issue (e.g., availability of
589 indoor and outdoor space for PA, access to age-appropriate equipment, etc.).

590 At the organizational level, in the context of the ECE settings, teacher/caregiver
591 support of PA, play equipment, and outdoor space were important factors associated with PA
592 of children while at the EEC setting (Bower et al., 2008; Copeland et al., 2012; Dowda et al.,
593 2004; Gubbels, Kremer, et al., 2012; Reilly, 2010).

594 At the policy level, there is much room for **improvement**. **While** the National
595 Association for Sport and Physical Education, AAP, and IOM have clear PA guidelines for
596 preschool-age children (AAP et al., 2012; IOM, 2011; NASPE, 2009), compliance is low
597 (Lessard et al., 2014; Nonas et al., 2004).

598 **CONCLUSIONS**

599 PA is a key component of overall physical, social, and mental health (AAP, 2006;
600 Goldfield et al., 2012; Hodges et al., 2013; O’Dwyer et al., 2012). Helping children set the
601 foundation for healthful PA habits early in life is a key component for achieving longer and
602 healthier lives for individuals and the general population. Promoting PA in early childhood
603 requires attention to the child and also caregivers—the home, early care and education (e.g.,
604 child care centers, family child care homes, etc.), and community settings in which care and
605 development of children take place. Despite an increasing number of studies focusing on the
606 PA of young children, much remains to be learned about the many factors—such as parenting
607 practices and styles, sociocultural factors, and environmental factors (including childcare
608 settings)—that influence the development and maintenance of PA and sedentary behaviors
609 among preschool-age children and, consequently, the development of obesity in childhood
610 (Lindsay, Sussner, Kim, & Gortmaker, 2006; Lindsay et al., 2009).

611 Since children's PA habits are initiated very early in life, early PA promotion
612 programs and intervention may not only have immediate health benefits, but may also help
613 reduce chronic disease risks when learned, healthful PA habits and preferences are carried into
614 adulthood (Davison, Edmunds, et al., 2011; Lindsay et al., 2006; Tremblay et al., 2011). This
615 literature provides a comprehensive synthesis of factors related to PA of young children at
616 multiple levels of influence that could be targeted in interventions.

617 **How Might this Information Affect Nursing Practice?**

630 This review provides a comprehensive synthesis of factors related to PA levels and
631 behaviors of young children that could assist pediatric nurses in their daily health promotion
632 and disease prevention efforts with families of young children and in the development of
633 childhood obesity prevention interventions.

634 With reducing and preventing childhood obesity remaining a public health priority,
635 nurses will continue to be engaged in childhood obesity prevention efforts through practice,
636 research, and education. PA assessment should be integrated as part of well-child visit
637 assessments in pediatric nursing primary care tasks of screening, communication and
638 anticipatory counseling. Integrating such assessment into electronic medical records would
639 help pediatric nurses routinely monitor children's PA behaviors as part of children's overall
640 health status assessment, creating opportunities to communicate to parents the importance of
641 early PA habits as part of their child's overall health. Given pediatric nurses' roles as primary
642 care providers and their frequent and continued contact with parents and children throughout
643 the early childhood years through well-child visits, immunization schedule, and minor acute
644 illnesses, they are well positioned to work with parents to promote and support the
645 development of healthful early physical activity behaviors of young children. **Pediatric nurses**

646 can play an important role in facilitating communication, education and provision of guidance
647 for parents regarding the importance of PA as well as factors influencing their children's PA
648 levels and behaviors. Nurses may be able to facilitate children's access to PA opportunities is
649 by providing parents with information and anticipatory guidance about various types of
650 physical activities which are developmentally appropriate for young children, as well as list of
651 local places free-of-cost or low-cost where young children and their families can be active.

652

653 REFERENCES

- 654 American Academy of Pediatrics (AAP). (2006). Active healthy living: Prevention of
655 childhood obesity through increased physical activity. *Pediatrics*, *117*, 1834-1842.
- 656 American Academy of Pediatrics (AAP), American Public Health Association, and National
657 Resource Center for Health and Safety in Child Care and Early Education. (2012).
658 *Preventing childhood obesity in early care and education programs: Selected*
659 *standards from caring for our children* (2nd ed.). Retrieved from
660 <http://nrckids.org/default>
661 [/assets/File/PreventingChildhoodObesity2nd.pdf](#)
- 662 Anand S, Krosnick JA. Demographic predictors of media use among infants, toddlers, and
663 preschoolers. *Am Behav Sci*. 2005;48(5):539–561. doi: 10.1177/0002764204271512.
- 664 Annesi, J. J., Smith A. E., & Tennant, G. A. (2013). Effects of the Start for Life treatment
665 on physical activity in primarily African American preschool children of ages 3-5
666 years. *Psychology, Health, and Medicine*, *18*(3), 300-309.
667 doi:10.1080/13548506.2012.712704

668 Ariza, A. J., Chen E. H., Binns H. J., Christoffel K. K. (2004) Risk factors for overweight in
669 five- to six-year-old Hispanic-American children: a pilot study. *J Urban*
670 *Health.*;81(1):150-61.

671 Asplund KM, Kair LR, Arain YH, Cervantes M, Oreskovic NM, Zuckerman KE. (2015).
672 Early Childhood Screen Time and Parental Attitudes
673 Toward Child Television Viewing in a Low-Income Latino Population Attending the
674 Special Supplemental Nutrition Program for Women, Infants, and Children.
675 *Child Obes.*,11(5):590-9. doi: 10.1089/chi.2015.0001. Epub 2015 Sep 21.

676 Bagley, S., Salmon, J., & Crawford, D. (2006). Family structure and children's television
677 viewing and physical activity. *Medicine & Science in Sports & Exercise*, 38(5), 910-
678 918.

679 Ball SC, Gillman MW, Mayhew M, Namenek Brouwer RJ, Benjamin Neelon SE. (2015)
680 Physical activity-related and weather-related practices of child care centers from
681 2 states. *J Phys Act Health.*, 12(2):238-44. doi: 10.1123/jpah.2013-0266. Epub 2014
682 Apr 17.

683 Barr R, Danziger C, Hilliard M, Andolina C, Ruskis J. (2010). Amount, content and context
684 of infant media exposure: a parental questionnaire and diary analysis. *Int J Early Years*
685 *Educ.* 18(2):107–122. doi: 10.1080/09669760.2010.494431.

686 Battista, R. A., Oakley, H., Weddell, M. S., Mudd, L. M., Greene, J. B., & West, S. T. (2014).
687 Improving the physical activity and nutrition environment through self-assessment
688 (NAP SACC) in rural area child care centers in North Carolina. *Preventive Medicine*,
689 67(Suppl 1), S10-16. doi:10.1016/j.ypped.2014.01.022.

690 Baumrind D. (1971). Current patterns of parental authority. *Developmental Psychology*
691 *Monograph*, Part 2, 4 (1): 1–103.

692 Beets, M. W., Bornstein, D., Dowda, M., & Pate, R. R. (2011). Compliance with national
693 guidelines for physical activity in U.S. preschoolers: Measurement and interpretation.
694 *Pediatrics*, 127(4), 658-664. doi:10.1542/peds.2010-2021

695 Bélanger M, Humbert L, Vatanparast H, Ward S, Muhajarine N, Chow AF, Engler-Stringer
696 R, Donovan D, Carrier N, Leis A. (2016). A multilevel intervention to
697 increase physical activity and improve healthy eating and physical literacy among
698 young children (ages 3-5) attending early childcare centres: the Healthy Start-Départ
699 Santé cluster randomised controlled trial study protocol. *BMC Public Health*.
700 12;16:313. doi: 10.1186/s12889-016-2973-5.

701 Benjamin, S. E., Cradock, A., Walker, E. M., Slining, M., & Gillman, M. W. (2008). Obesity
702 prevention in child care: A review of U.S. state regulations. *BMC Public Health*,
703 30(8), 188. doi:10.1186/1471-2458-8-188.

704 Bentley GF, Turner KM, Jago R. (2016) Mothers' views of their preschool child's screen-
705 viewing behaviour: a qualitative study. *BMC Public Health*, 16:718. doi:
706 10.1186/s12889-016-3440-z.

707 Boldemann, C., Blennow, M., Dal, H., Mårtensson, F., Raustorp, A., Yuen, K., & Wester, U.
708 (2006). Impact of preschool environment upon children's physical activity and sun
709 exposure. *Preventive Medicine*, 42, 301-308.

710 Bower, J. K., Hales, D. P., Tate, D. F., Rubin, D. A., Benjamin, S. E., & Ward, D. S. (2008).
711 The child care environment and children's physical activity. *American Journal of*
712 *Preventive Medicine*, 34(1), 23-29.

713 Burdette, H. L., Wadden, T. A., & Whitaker, R. C. (2006). Neighborhood safety, collective
714 efficacy, and obesity in women with young children. *Obesity, 14*(3), 518-525.

715 Burdette, H. L., & Whitaker, R. C. (2005). Resurrecting free play in young children: Looking
716 beyond fitness and fatness to attention, affiliation, and affect. *Archives of Pediatric
717 and Adolescent Medicine, 159*(1), 46-50.

718 Burdette, H. L., Whitaker, R. C., & Daniels, S. R. (2004). Parental report of outdoor playtime
719 as a measure of physical activity in preschool-aged children. *Archives of Pediatric and
720 Adolescent Medicine, 158*(4), 353-357.

721 Buscemi, J., Kanwisher, K., Becker, A. B., Ward, D. S., & Fitzgibbon, M. L. (2015). Society
722 of Behavioral Medicine position statement: Early care and education (ECE) policies
723 can impact obesity prevention among preschool-aged children. *Translational
724 Behavioral Medicine, 5*(1), 122-125.

725 Carlson SA, Fulton JE, Lee SM, Foley JT, Heitzler C, Huhman M. (2010). Influence of limit-
726 setting and participation in physical activity on youth screen time. *Pediatrics.*,
727 126(1):e89-96. doi: 10.1542/peds.2009-3374.
728

729 Carson, V., & Janssen, I. (2012). Associations between factors with the home setting and
730 screen time among children aged 0-5 years: A cross-sectional study. *BMC Public
731 Health, 12*, 539. doi:10.1186/1471-2458-12-539.

732 Carson, V., Stearns, J., & Janssen, I. (2015). The Relationship Between Parental Physical
733 Activity and Screen Time Behaviors and the Behaviors of their Young Children.
734 *Pediatr Exerc Sci., 27*(3):390-5. doi: 10.1123/pes.2014-0214.

735 Cerin E, Baranowski T, Barnett A, Butte N, Hughes S, Lee RE, Mendoza JA, Thompson D,
736 O'Connor TM. (2016). Places where prechoolers are (in) active: an observational

737 study on Latino preschoolers and their parents using objective measures. *Int J Behav*
738 *Nutr Phys Act.* 29;13:29. doi: 10.1186/s12966-016-0355-0.

739 Certain LK, Kahn RS. (2002). Prevalence, correlates, and trajectory of television viewing
740 among infants and toddlers. *Pediatrics.* 109(4):634–642. doi: 10.1542/peds.109.4.634

741 Cespedes, E. M., McDonald, J., Haines, J., Bottino, C. J., & Taveras, E. M. (2013). Obesity-
742 related behaviors of U.S. and non-U.S. born parents and children in low-income
743 households. *Journal of Developmental and Behavioral Pediatrics, 34*(8), 541-548.

744 Chen, T. A., O'Connor, T. M., Hughes, S. O., Frankel, L., Baranowski, J., Mendoza, J. A.,
745 Thompson, D., & Baranowski, T. (2013). TV parenting practices: Is the same scale
746 appropriate for parents of children of different ages? *International Journal for*
747 *Behavioral Nutrition and Physical Activity, 10*(41). doi:10.1186/1479-5868-10-41.

748 Cheng S, Maeda T, Yoichi S, Yamagata Z, Tomiwa K. (2010). Early television exposure and
749 children's behavioral and social outcomes at age 30 months. *J Epidemiol.* 20(Suppl
750 2):S482–S489.

751 Chuang R. J., Sharman S., Skala, K., & Evans, A. (2013). Ethnic differences in the home
752 environment and physical activity behaviors among low-income, minority
753 preschoolers in Texas. *American Journal of Health Promotion, 27*(4), 270-278.
754 doi:10.4278/ajhp
755 .110427-QUAN-171

756 Commonwealth of Australia, Department of Health and Ageing. (2010). *Move and play every*
757 *day: National physical activity recommendations for children 0 to 5 years.* Retrieved
758 from <http://www.health.gov.au/internet/main/publishing.nsf/content>

759 /9D831D9E6713F92ACA257BF0001F5218/\$File/0-5yrACTIVE_Brochure
760 _FA%20SCREEN.pdf

761 Copeland, K. A., Kendeigh, C. A., Saelens, B. E., Kalkwarf, H. J., & Sherman, S. N. (2012).
762 Physical activity in child-care centers: Do teachers hold the key to the playground?
763 *Health Education Research*, 27(1), 81-100.

764 Copeland KA, Sherman SN, Khoury JC, Foster KE, Saelens BE, Kalkwarf HJ. (2011). Wide
765 variability in physical activity environments and weather-related outdoor play policies
766 in childcare centers within a single county of Ohio. *Arch Pediatr Adolesc*
767 *Med.*,165(5):435-42. doi: 10.1001/archpediatrics.2010.267. Epub 2011 Jan 3.

768 Copeland KA, Khoury JC, Kalkwarf HJ. (2016). Child Care Center Characteristics Associated
769 With Preschoolers' Physical Activity. *Am J Prev Med.*, 50(4):470-9. doi:
770 10.1016/j.amepre.2015.08.028.

771 Cradock, A. L., O'Donnell, E. M., Benjamin, S. E., Walker, E., & Slining, M. (2010). A
772 review of state regulations to promote physical activity and safety on playgrounds
773 in child care centers and family child care homes. *Journal of Physical Activity and*
774 *Health*, 7(Suppl 1), S108-119.

775 Davison, K. K., & Birch, L. L. (2002). Obesogenic families: Parents' physical activity and
776 dietary intake patterns predict girls' risk of overweight. *International Journal of*
777 *Obesity*, 26, 1186-1193.

778 Davison, K. K., Edmunds, L. S., Wyker, B. A., Young, L. M., Sarfoh, V. S., & Sekhobo, J. P.
779 (2011). Feasibility of increasing childhood outdoor play and decreasing television
780 viewing through a family-based intervention in WIC, New York State, 2007-2008.
781 *Prevention of Chronic Disease*, 8, A54.

782 Dawson-Hahn, E. E., Fesinmeyer, M. D., & Mendoza, J. A. (2015). Correlates of physical
783 activity in Latino preschool children attending Head Start. *Pediatric Exercise Science*,
784 27(3), 372-379. doi:10.1123/pes.2014-0144.

785 Dalzell VP, Msall ME, High PC. (2000). Parental attitudes of television and videocassette
786 viewing of children aged birth to 36 months. *J Dev Behav Pediatr.* 21(5):390.

787 De Decker, E., De Craemer, M., De Bourdeaudhuij, I., Wijndaele, K., Duvinage, K.,
788 Koletzko, B., ... Cardon, G. (2012). Influencing factors of screen time in preschool
789 children: An exploration of parents' perceptions through focus groups in six European
790 countries. *Obesity Review*, (Suppl 1), 75-84. doi:10.1111/j.1467-789X.2011.00961.x.

791 Dennison BA, Erb TA, Jenkins PL. (2002). Television viewing and television in bedroom
792 associated with overweight risk among low-income preschool children.
793 *Pediatrics.* 109(6):1028-35.

794 Department of Health, Physical Activity, Health Improvement and Protection. (2011). *Start*
795 *active, stay active: A report on physical activity for health from the four home*
796 *countries' Chief Medical Officers.* London, England: Author.

797 Dowda, M., Pate, R., Trost, S., Almeida, M., & Sirard, J. (2004). Influences of preschool
798 policies and practices on children's physical activity. *Journal of Community Health*,
799 29, 183-196.

800 Dowda, M., Pfeiffer K. A., Brown W. H., Mitchell J. A., Byun W., Pate R. R. (2011)
801 Parental and environmental correlates of physical activity of children attending
802 preschool. *Arch Pediatr Adolesc Med.* 165(10):939-44. doi:
803 10.1001/archpediatrics.2011.84. Epub 2011 Jun 6.

804 Downing KL, Hinkley T, Hesketh KD. (2015). Associations of Parental Rules and
805 Socioeconomic Position With Preschool Children's Sedentary
806 Behaviour and Screen Time. *J Phys Act Health.*, 12(4):515-21. doi:
807 10.1123/jpah.2013-0427.

808 Duffey, K. J., Slining, M. M., & Benjamin Neelon, S. E. (2014). States lack physical
809 activity policies in child care that are consistent with national recommendations.
810 *Childhood Obesity*, 10(6), 491-500. doi:10.1089/chi.2014.0096

811 Dwyer, G. M., Higgs, J., Hardy, L. L., & Baur, L. A. (2008). What do parents and preschool
812 staff tell us about young children's physical activity: A qualitative study. *International*
813 *Journal of Behavioral Nutrition and Physical Activity*, 5, 66. doi:10.1186/1479-5868-
814 5-66

815 Dwyer, J., Needham, L., Simpson, J. R., & Heeney, E. S. (2008). Parents report intrapersonal,
816 interpersonal, and environmental barriers to supporting healthy eating and physical
817 activity among their preschoolers. *Applied Physiology, Nutrition, and Metabolism*,
818 33(2), 338-46. doi:10.1139/H07-195.

819 Edwards MJ, Jago R, Sebire SJ, Kesten JM, Pool L, Thompson JL. The influence of friends
820 and siblings on the physical activity and screen viewing behaviours of children aged 5-
821 6 years: a qualitative analysis of parent interviews. *BMJ Open.*, 5(5):e006593. doi:
822 10.1136/bmjopen-2014-006593.

823 Ekelund, U., Brage, S., & Wareham, N. J. (2004). Physical activity in young children. *Lancet*,
824 363(9415), 1163-1164.

825 Ergler, C. R., Kearns, R. A., & Witten, K. (2013). Seasonal and locational variations in
826 children's play: Implications for wellbeing. *Social Science Medicine*, 91, 178-185.

827 Fakhouri, T. H., Hughes, J. P., Brody, D. J., Kit, B. K., & Ogden, C. L. (2013). Physical
828 activity and screen-time viewing among elementary school-aged children in the United
829 States from 2009 to 2010. *JAMA Pediatrics*, *167*(3), 223-229.

830 Finch M, Wolfenden L, Falkiner M, Edenden D, Pond N, Hardy LL, Milat AJ, Wiggers J.
831 (2012). Impact of a population based intervention to increase the adoption of
832 multiple physical activity practices in centre based childcare services: a quasi
833 experimental, effectiveness study. *Int J Behav Nutr Phys Act.* *29*;9:101. doi:
834 10.1186/1479-5868-9-101.

835 Finn, K., Johannsen, N., & Specker, B. (2002). Factors associated with physical activity in
836 preschool children. *Journal of Pediatrics*, *140*(1), 81-85.

837 Fitzgibbon, M.L., Stolley, M.R., Schiffer, L.A., Braunschweig, C.L., Gomez, S.L., Van Horn
838 L., Dyer, A.R. (2011). Hip-Hop to Health Jr. Obesity Prevention Effectiveness Trial:
839 post-intervention results. *Obesity (Silver Spring)*.;19(5):994-1003. doi:
840 10.1038/oby.2010.314.

841 Fitzgibbon, M.L., Stolley, M.R., Schiffer, L.A., Van Horn L., Christoffel K.K., Dyer, A.R.
842 (2006). Hip-Hop to Health Jr. for Latino Preschool Children. *Obesity* *14* (9): 1617-
843 1625.

844 Gallagher MR. (2010). Maternal perspectives on lifestyle habits that put children of Mexican
845 descent at risk for obesity. *J Spec Pediatr Nurs.*;15(1):16-25. doi: 10.1111/j.1744-
846 6155.2009.00213.x.

847 Garriguet D, Carson V, Colley RC, Janssen I, Timmons BW, Tremblay MS. (2016). Physical
848 activity and sedentary behavior of Canadian children aged 3 to 5. *Health Rep.* *21*;
849 27(9):14-23.

850 Goldfield, G. S., Harvey, A., Grattan, K., & Adamo, K. B. (2012). Physical activity promotion
851 in the preschool years: A critical period to intervene. *International Journal of*
852 *Environmental Research and Public Health*, 9(4), 1326-1342.

853 Grigsby-Toussaint, D. S., Chi, S. H., & Fiese, B. H. (2011). Where they live, how they play:
854 Neighborhood greenness and outdoor physical activity among preschoolers.
855 *International Journal of Health Geography*, 10, 66.

856 Gubbels, J. S., Kremer, S. P., Van Kann, D. H., Stafleu, A., Candel, M. J., Dagnelie, P. C., ...
857 de Vries, N. K. (2012). Interaction between physical environment, social environment,
858 and child characteristics in determining physical activity at child care. *Health*
859 *Psychology*, 30(1), 84-90.

860 Gubbels, J. S., Van Kann, D. H., & Jansen, M. W. (2012). Play equipment, physical
861 activity opportunities, and children's activity levels at child care. *Journal of*
862 *Environmental and Public Health*, 2012. doi:10.1155/2012/326520

863 Gubbels J. S., Kremers S. P. J., Stafleu A., de Vries SI, Goldbohm RA, Dagnelie PC, de Vries
864 NK, van Buuren S, Thijs C. (2011). Association between parenting practices and
865 children's dietary intake, activity behavior and development of body mass index: the
866 KOALA birth cohort study. *International Journal of Behavioral Nutrition and*
867 *Physical Activity*. 8, article 18 doi: 10.1186/1479-5868-8-18.

868 Hnatiuk JA, Ridgers ND, Salmon J, Hesketh KD. (2016). Maternal correlates of young
869 children's physical activity across periods of the day. *J Sci Med Sport*. 14. pii: S1440-
870 2440(16)30119-0. doi: 10.1016/j.jsams.2016.06.014.

871 Hart, K. H., Herriot, A., Bishop, J. A., & Truby, H. (2003). Promoting healthy diet and
872 exercise patterns amongst primary school children: A qualitative investigation of
873 parental perspectives. *Journal of Human Nutrition and Dietetics*, *16*, 89-96.

874 He, M., Irwin, J. D., Sangster Bouck, L. M., Tucker, P., & Pollett, G. L. (2005). Screen-
875 viewing behaviors among preschoolers: Parents' perceptions. *American Journal of*
876 *Preventive Medicine*, *29*, 120-125. doi:10.1016/j.amprepe.2005.04.004

877 Hesketh KR, Griffin SJ, van Sluijs EM. (2015). UK Preschool-aged children's physical
878 activity levels in childcare and at home: a cross-sectional exploration. *Int J Behav Nutr*
879 *Phys Act.*, *26*, 12:123. doi: 10.1186/s12966-015-0286-1.

880 Hesketh, K. R., Goodfellow, L., Ekelund, U., McMinn, A. M., Godfrey, K. M., Inskip, H. M.,
881 ... van Sluijs, E. M. F. (2014). Activity levels in mothers and their preschool children.
882 *Pediatrics*, *133*(4), 973-980.

883 Hesketh, K. R., McMinn, A. M., Griffin, S. J., Harvey, N. C., Godfrey, K. M., Inskip, H. M.,
884 ... van Sluijs, E. M. (2013). Maternal awareness of young children's physical activity:
885 Levels and cross-sectional correlates of overestimation. *BMC Public Health*, *13*, 924.

886 Hinkley, T., Crawford, D., Salmon, J., Okely, A. D., & Hesketh, K. (2008). Preschool
887 children and physical activity: A review of correlates. *American Journal of Preventive*
888 *Medicine*, *34*(5), 435-441.

889 Hinkley, T., Salmon, J., Okely, A. D., Crawford, D., & Hesketh, K. (2011). Influences on
890 preschool children's physical activity: Exploration through focus groups. *Family and*
891 *Community Health*, *34*(1), 39-50.

892 Hinkley T., Salmon J., Okely A. D., Crawford D. (2013). The correlates of preschoolers'
893 compliance with screen recommendations exist across multiple domains. *Preventive*
894 *Medicine*. 57(3):212–219. doi: 10.1016/j.ypmed.2013.05.020.

895 Hinkley T, Salmon J, Crawford D, Okely AD, Hesketh KD. (2016). Preschool and childcare
896 center characteristics associated with children's physical activity during care hours: an
897 observational study. *Int J Behav Nutr Phys Act*. 11;13(1):117.

898 Hodges, E. A., Smith, C., Tidwell, S., & Berry, D. (2013). Promoting physical activity in
899 preschoolers to prevent obesity: A review of the literature. *Journal of Pediatric*
900 *Nursing*, 28(1), 3-19.

901 Horodyski MA, Stommel M, Brophy-Herb HE, Weatherspoon L. (2010). Mealtime
902 television viewing and dietary quality in low-income African American and Caucasian
903 mother-toddler dyads. *Matern Child Health J*. 14(4):548–556. doi: 10.1007/s10995-
904 009-0501-2.

905 Institute of Medicine (IOM). (2011). *Early childhood obesity prevention policies*.
906 Washington, DC: The National Academies Press.

907 Jackson, D. M., Djafarian, K., Stewart, J., & Speakman, J. R. (2009). Increased television
908 viewing is associated with elevated body fatness but not with lower total energy
909 expenditure in children. *American Journal of Clinical Nutrition*, 89(4), 1031-1036.
910 doi:10.3945/ajcn.2008.26746

911 Jones, S., Hendricks, S., & Draper, C. E. (2014). Assessment of physical activity and
912 sedentary behavior at preschools in Cape Town, South Africa. *Childhood Obesity*,
913 10(6), 501-510. doi:10.1089/chi.2014.0097

914 Kaciroti, N., Staples-Watson, J., & Lumeng, J. C. (2012). Enhancing self-regulation as a
915 strategy for obesity prevention in Head Start preschoolers: The growing healthy study.
916 *BMC Public Health*, 30(12), 1040.

917 Kelly, L. A., Reilly, J. J., Fisher, A., Montgomery, C., Williamson, A., McColl, J. H., ...
918 Grant, S. (2006). Effect of socioeconomic status on objectively measured physical
919 activity. *Archives of Disease in Childhood*, 91(1), 35-38.
920 doi:10.1136/adc.2005.080275.

921 Kourlaba G, Kondaki K, Liarigkovinos T, Manios Y. (2009). Factors associated with
922 television viewing time in toddlers and preschoolers in Greece: the GENESIS study. *J*
923 *Public Health (Oxf)* 31(2):222–230. doi: 10.1093/pubmed/udp011.

924 Kuepper-Nybelen, J., Lamerz, A., Bruning, N., Hebebrand, J., Herpertz-Dahlmann, B., &
925 Brenner, H. (2005). Major differences in prevalence of overweight according to
926 nationality in preschool children living in Germany: Determinants and public health
927 implications. *Archives of Disease in Childhood*, 90, 359-363.

928 Kumanyika, S., & Grier, S. (2006). Targeting interventions for ethnic minority and low-
929 income populations. *Future Child*, 16(1), 187-207.

930 Lampard AM, Jurkowski JM, Davison KK. (2013). The family context of low-income parents
931 who restrict child screen time. *Child Obes.*, 9(5):386-92. doi: 10.1089/chi.2013.0043.

932 Lapierre MA, Piotrowski JT, Linebarger DL. (2012). Background television in the homes of
933 US children. *Pediatrics*. 2012;130(5):839–846. doi: 10.1542/peds.2011-2581.

934 LaRowe TL, Tomayko EJ, Meinen AM, Hoiting J, Saxler C, Cullen B; Wisconsin Early
935 Childhood Obesity Prevention Initiative (WECOPI). (2016) Active Early: one-
936 year policy intervention to increase physical activity among early care and education

937 programs in Wisconsin. *BMC Public Health.*, 16:607. doi: 10.1186/s12889-016-3198-
938 3.

939 Larson, N., Ward, D. S., Neelon, S. B., & Story, M. (2011). What role can child-care settings
940 play in obesity prevention? A review of the evidence and call for research efforts.
941 *Journal of the American Dietetic Association*, 111(9), 1343-1362.

942 Lessard, L., Lesesne, C., Kakietek, J., Breck, A., Jernigan, J., Dunn, L., ... Khan, L. K.
943 (2014). Measurement of compliance with New York City's regulations on beverages,
944 physical activity, and screen time in early child care centers. *Prevention of Chronic
945 Disease*, 16(11), E183. doi:10.5888/pcd11.130433.

946 Levin S, Martin MW, Riner WF. (2004). TV viewing habits and body mass index among
947 South Carolina Head Start children. *Ethn Dis.* ;14(3):336-9.

948 Lindsay AC, Greaney ML, Wallington SF, Sands FD, Wright JA, Salkeld J. (2017).
949 Latino parents' perceptions of the eating and physical activity experiences of their pre-
950 school children at home and at family child-care homes: a qualitative study. *Public
951 Health Nutr.*;20(2):346-356. doi: 10.1017/S136898001600207X.

952 Lindsay, A. C., Sussner, K. M., Greaney, M. L., & Peterson, K. E. (2009). Influence of social
953 context on eating, physical activity, and sedentary behaviors of Latina mothers and
954 their preschool-age children. *Health Education and Behavior*, 36, 81-96.

955 Lindsay, A. C., Sussner, K. M., Kim, J., & Gortmaker, S. L. (2006). The role of parents in
956 preventing childhood obesity. *Future Child*, 16(1), 169-186.

957 Linebarger DL, Walker D. (2005) Infants' and toddlers' television viewing and language
958 outcomes. *Am Behav Sci.* 48(5):624-645. doi: 10.1177/0002764204271505.

959 Loprinzi, P. D., & Trost, S. G. (2010). Parental influences on physical activity behavior in
960 preschool children. *Preventive Medicine*, 50(3), 129-133.

961 Lumeng JC, Rahnama S, Appugliese D, Kaciroti N, Bradley RH. (2006). Television exposure
962 and overweight risk in preschoolers. *Arch Pediatr Adolesc Med.* 160(4):417–422. doi:
963 10.1001/archpedi.160.4.417.

964 Manios Y, Kondaki K, Kourlaba G, Grammatikaki E, Birbilis M, Ioannou E. (2009).
965 Television viewing and food habits in toddlers and preschoolers in Greece: the
966 GENESIS study. *Eur J Pediatr.* 168(7):801–808. doi: 10.1007/s00431-008-0838-3.

967 Masur EF, Flynn V. (2008). Infant and mother-infant play and the presence of the television. *J*
968 *Appl Dev Psychol.* 29(1):76–83. doi: 10.1016/j.appdev.2007.10.001.

969 McLeroy, K. R., Bibeau, D., Steckler, A., & Glanz, K. (1988). An ecological perspective on
970 health promotion programs. *Health Education Quarterly*, 15(4), 351-377.

971 McKee, D. P., Murtagh, E. M., Boreham, C. A., Nevill, A. M., & Murphy, M. H. (2012).
972 Seasonal and annual variation in young children’s physical activity. *Medicine &*
973 *Science in Sports & Exercise*, 44(7), 1318-1324.
974 doi:10.1249/MSS.0b013e3182464db5

975 McKenzie, T. L., Baquero, B., Crespo, N. C., Arredondo, E. M., Campbell, N. R., & Elder,
976 J. P. (2008). Environmental correlates of physical activity in Mexican American
977 children at home. *Journal of Physical Activity and Health*, 5(4), 579-591.

978 McLeroy, K. R., Bibeau, D., Steckler, A., & Glanz, K. (1998). An ecological perspective on
979 health promotion programs. *Health Education Quarterly*, 15(4), 351-377.

980 McLearn KT, Minkovitz CS, Strobino DM, Marks E, Hou W. (2006). The timing of maternal
981 depressive symptoms and mothers’ parenting practices with young children:

982 implications for pediatric practice. *Pediatrics*. 118(1):e174–e182. doi:
983 10.1542/peds.2005-1551.

984 McPherson, M. E., & Homer, C. J. (2011). Policies to support obesity prevention for children:
985 A focus on of early childhood policies. *Pediatric Clinics of North America*, 58(6),
986 1521-1541.

987 Mendelsohn AL, Dreyer BP, Brockmeyer CA, Berkule-Silberman SB, Huberman HS,
988 Tomopoulos S. (2011). Randomized controlled trial of primary care pediatric
989 parenting programs: effect on reduced media exposure on infants, mediated through
990 enhanced parent–child interaction. *Arch Pediatr Adolesc Med*. 165(1):42–48. doi:
991 10.1001/archpediatrics.2010.266.

992 Mendoza, J. A., Zimmerman, F. J., & Christakis, D. A. (2007). Television viewing, computer
993 use, obesity, and adiposity in US preschool children. *International Journal of*
994 *Behavioral Nutrition and Physical Activity*, 4, 44.

995 Montgomery, C., Reilly, J., Jackson, D., Kelly, L. A., Slater, C., Paton, J. Y., & Grant, S.
996 (2004). Relation between physical activity and energy expenditure in a representative
997 sample of young children. *American Journal of Clinical Nutrition*, 80(3), 591-596.

998 National Association for Sport and Physical Education (NASPE). (2009). *Active start: A*
999 *statement of physical activity guidelines for children from birth to age 5* (2nd ed.).
1000 Sewickley, PA: American Alliance for Health, Physical Education, Recreation, and
1001 Dance.

1002 National Resource Center for Health and Safety in Child Care and Early Education (NRC),
1003 University of Colorado Denver. (2011). *Achieving a state of healthy weight: A*

1004 *national assessment of obesity prevention terminology in Child Care Regulations*
1005 2010. Aurora, CO: Author.

1006 Nonas, C., Silver, L. D., Kettel Khan, L., & Leviton, L. (2004). Rationale for New York
1007 City's regulations on nutrition, physical activity, and screen time in early child care
1008 centers. *Prevention of Chronic Disease, 11*, E182. doi:10.5888/pcd11.130435

1009 O'Connor, T. M., Cerin, E., Hughes, S. O., Robles, J., Thompson, D., Baranowski, T., ...
1010 Shewchuk, R. M. (2013). What Hispanic parents do to encourage and discourage 3-4
1011 year old children to be active: A qualitative study using nominal group technique.
1012 *International Journal of Behavioral Nutrition and Physical Activity, 10*, 93.
1013 doi:10.1186/1479-5868-10-93

1014 O'Connor, T. M., Cerin, E., Lee, R. E., Parker, N., Chen, T. A., Hughes, S. O., ...
1015 Baranowski, T. (2014). Environmental and cultural correlates of physical activity
1016 parenting practices among Latino parents with preschool-aged children: Niños activos.
1017 *BMC Public Health, 14*, 707. doi:10.1186/1471-2458-14-707

1018 O'Connor, T. M., Chen, T. A., Baranowski, J., Thompson, D., & Baranowski, T. (2013).
1019 Physical activity and screen-media-related parenting practices have different
1020 associations with children's objectively measured physical activity. *Childhood Obesity,*
1021 9(5), 446-453.

1022 O'Dwyer, M. V., Fairclough, S. J., Knowles, Z., & Stratton, G. (2012). Effect of a family
1023 focused active play intervention on sedentary time and physical activity in preschool
1024 children. *International Journal of Behavioral Nutrition and Physical Activity, 9*(1), 1-
1025 13.

- 1026 Okely, A. D., Salmon, J., Trost, S. G., & Hinkley, T. (2008). *Discussion paper for the*
1027 *development of physical activity recommendations for children under five years.*
1028 Canberra, Australia: Commonwealth Department of Health and Ageing.
- 1029 Oliver, M., Schofield, G. M., & Schluter, P. J. (2010). Parent influences on preschoolers'
1030 objectively assessed physical activity. *Journal of Science and Medicine in Sport,*
1031 *13(4), 403-409.*
- 1032 Østbye, T., Malhotra, R., Stroe, M., Lovelady, C., Brouwer, R., Zucker, N., Fuemmeler, B.
1033 (2013). The effect of the home environment on physical activity and dietary intake
1034 in preschool children. *International Journal of Obesity (London), 37(10), 1314-21.*
1035 doi:10.1038/ijo.2013.76
- 1036 Pate, R. R., O'Neill, J. R., Brown, W. H., Pfeiffer, K. A., Dowda, M., & Addy, C. L. (2015).
1037 Prevalence of Compliance with a New Physical Activity Guideline for Preschool-Age
1038 Children. *Childhood Obesity, 11(4), 415-420.* doi:10.1089/chi.2014.0143
- 1039 Pate, R. R., Pfeiffer, K. A., Trost, S. G., Ziegler, P., & Dowda, M. (2004). Physical activity
1040 among children attending preschools. *Pediatrics, 114, 1258-1263.*
- 1041 Pearson, N., Salmon, J., Crawford, D., Campbell, K., & Timperio, A. (2011). Are parental
1042 concerns for child TV viewing associated with child TV viewing and the home
1043 sedentary environment? *International Journal of Behavioral Nutrition and Physical*
1044 *Activity, 8, 102.* doi:10.1186/1479-5868-8-102
- 1045 Pfeiffer, K. A., Dowda, M., McIver, K. L., & Pate, R. R. (2009). Factors related to objectively
1046 measured physical activity in preschool children. *Pediatric Exercise Science, 21(2),*
1047 *196-208.*

- 1048 Pratt, M., Epping, J. N., & Dietz, W. H. (2009). Putting physical activity into public health: A
1049 historical perspective from the CDC. *Preventive Medicine, 49*(4), 301-302.
- 1050 Prochaska, J.O.; DiClemente, C.C. The transtheoretical approach. In: Norcross, J.C.; Goldfried,
1051 M.R. (eds.) *Handbook of psychotherapy integration*. 2nd ed. New York: Oxford
1052 University Press; 2005. p.147–171.
- 1053 Rodríguez-Oliveros, G., Haines, J., Ortega-Altamirano, D., Power, E., Taveras, E. M.,
1054 González-Unzaga, M. A., & Reyes-Morales, H. (2011). Obesity determinants in
1055 Mexican preschool children: Parental perceptions and practices related to feeding and
1056 physical activity. *Archives of Medical Research, 42*(6), 532-539.
- 1057 Roemmich, J. N., Epstein, L. H., Raja, S., Yin, L., Robinson, J., & Winiewicz, D. (2006).
1058 Association of access to parks and recreational facilities with the physical activity of
1059 young children. *Preventive Medicine, 43*(6), 437-441.
- 1060 Rosenstock, I. (1974). *Historical Origins of the Health Belief Model*. *Health Education*
1061 *Monographs*. Vol. 2 No. 4.
- 1062 Ruiz, R., Gesell, S. B., Buchowski, M. S., Lambert, W., & Barkin, S. L. (2011). The
1063 relationship between Hispanic parents and their preschool-aged children’s physical
1064 activity. *Pediatrics, 127*(5), 888-895.
- 1065 Sallis JF, Cervero RB, Ascher W, Henderson KA, Kraft MK, Kerr J. (2006).
1066 An ecological approach to creating active living communities. *Annu Rev Public*
1067 *Health., 27*:297-322.
- 1068 Salmon, J., Ball, K., Crawford, D., Booth, M., Telford, A., Hume, C., ... Worsley, A. (2005).
1069 Reducing sedentary behaviour and increasing physical activity among 10-year-old

1070 children: Overview and process evaluation of the “switch-play” intervention. *Health*
1071 *Promotion International*, 20(1), 7-17.

1072 Salmon, J., Owen, N., Crawford, D., Bauman, A., & Sallis, J. F. (2003). Physical activity and
1073 sedentary behavior: A population-based study of barriers, enjoyment, and preference.
1074 *Health Psychology*, 22(2), 178-188.

1075 Salmon, J., Veitch, J., Abbott, G., ChinAPaw, M., Brug, J. J., teVelde, S. J., ... Ball, K.
1076 (2013). Are associations between the perceived home and neighbourhood environment
1077 and children’s physical activity and sedentary behaviour moderated by urban/rural
1078 location? *Health Place*, 24, 44-53. doi:10.1016/j.healthplace.2013.07.010

1079 Sawyer, A., Smith, L., Schrepft, S., van Jaarsveld, C. H., Wardle, J., & Fisher, A. (2014).
1080 Primary caregiver knowledge of paediatric physical activity recommendations in the
1081 United Kingdom and its association with caregiver behaviour: An observational study.
1082 *BMC Public Health*, 14, 795. doi:10.1186/1471-2458-14-795.

1083 Schary D. P., Cardinal B. J., Loprinzi P. D. (2012). Parental support exceeds parenting style
1084 for promoting active play in preschool children. *Early Child Development and*
1085 *Care*. 182(8):1057–1069. doi: 10.1080/03004430.2012.685622.

1086 Schlechter CR, Rosenkranz RR, Fees BS, Dzewaltowski DA. Preschool Daily Patterns
1087 of Physical Activity Driven by Location and Social Context. *J Sch Health*. 2017
1088 Mar;87(3):194-199. doi: 10.1111/josh.12486.

1089 Sigmundová D, Sigmund E, Badura P, Vokáčová J, Trhlíková L, Bucksch J. (2016).
1090 Weekday-weekend patterns of physical activity and screen time in parents and their
1091 preschoolers. *BMC Public Health*, 16:898. doi: 10.1186/s12889-016-3586-8.

1092 Sisson, S. B., Church, T. S., Martin, C. K., Tudor-Locke, C., Smith, S. R., Bouchard, C., ...
1093 Katzmarzyk, P. T. (2009). Profiles of sedentary behavior in children and adolescents:
1094 The U.S. National Health and Nutrition Examination Survey, 2001-2006. *International*
1095 *Journal of Pediatric Obesity*, 4(4), 353-359.

1096 Schmidt ME, Rich M, Rifas-Shiman SL, Oken E, Taveras EM. (2009). Television viewing in
1097 infancy and child cognition at 3 years of age in a US cohort. *Pediatrics*. 123(3):e370–
1098 e375. doi: 10.1542/peds.2008-3221.

1099 Smith BJ, Grunseit A, Hardy LL, King L, Wolfenden L, Milat A. (2010). Parental influences
1100 on child physical activity and screen viewing time: a population based study. *BMC*
1101 *Public Health*,10:593. doi: 10.1186/1471-2458-10-593.

1102 Stephens RL, Xu Y, Lesesne CA, Dunn L, Kakietek J, Jernigan J, Khan LK. (2014).
1103 Relationship between child care centers' compliance with physical activity regulations
1104 and children's physical activity, New York City, 2010. *Prev Chronic Dis.*, 11:E179.
1105 doi: 10.5888/pcd11.130432.

1106 Soltero EG, Cerin E, Lee RE, O'Connor TM. (2016). Associations Between Objective and
1107 Self-Report Measures of Traffic and Crime Safety in Latino Parents of Preschool
1108 Children. *J Immigr Minor Health*. 1-12. doi:10.1007/s10903-016-0498-8. Sep 28.
1109 [Epub ahead of print]

1110 Tandon PS, Walters KM, Igoe BM, Payne EC, Johnson DB. (2016). Physical
1111 Activity Practices, Policies and Environments in Washington State Child
1112 Care Settings: Results of a Statewide Survey. *Matern Child Health J.*, 2016 Jul 22.
1113 [Epub ahead of print]

1114 Tandon, P. S., Saelens, B. E., Zhou, C., Kerr, J., & Christakis, D. A. (2013). Indoor versus
1115 outdoor time in preschoolers at child care. *American Journal of Preventive Medicine*,
1116 44(1), 85-88.

1117 Taylor, R. W., Murdoch, L., Carter, P., Gerrard, D. F., Williams, S. M., & Taylor, B. J.
1118 (2009). Longitudinal study of physical activity and inactivity in preschoolers: The
1119 FLAME study. *Medicine & Science in Sports & Exercise*, 41(1), 96-102.

1120 The World Bank. Data High income: OECD. 2013.

1121 Thompson AL, Adair LS, Bentley ME. (2013). Maternal characteristics and perception of
1122 temperament associated with infant TV exposure. *Pediatrics*. 131(2):e390–e397. doi:
1123 10.1542/peds.2012-1224.

1124 Thompson DA, Christakis DA. (2007). The association of maternal mental distress with
1125 television viewing in children under 3 years old. *Ambulatory Pediatr*. 7(1):32–37. doi:
1126 10.1016/j.ambp.2006.09.007.

1127 Thompson D.A., Polk S, Cheah C.S., Vandewater E.A., Johnson S.L., Chrismer M.C.,
1128 Tschann J.M. (2015). Maternal Beliefs and Parenting Practices Regarding
1129 Their Preschool Child's Television Viewing: An Exploration in a Sample of Low-
1130 Income Mexican-Origin Mothers. *Clin Pediatr (Phila)*. 54(9):862-70. doi:
1131 10.1177/0009922815574074. Epub 2015 Feb 26.

1132 Thompson DA, Christakis DA. (2005). The association between television viewing and
1133 irregular sleep schedules among children less than 3 years of
1134 age. *Pediatrics*. 116(4):851–856. doi: 10.1542/peds.2004-2788.

1135 Thompson DA, Sibinga EM, Jennings JM, Bair-Merritt MH, Christakis DA. (2010).
1136 Television viewing by young Hispanic children: evidence of heterogeneity. *Arch*
1137 *Pediatr Adolesc Med.* 164(2):174–179. doi: 10.1001/archpediatrics.2009.257.

1138 Tomopoulos S, Dreyer BP, Valdez P. et al. (2007). Media content and externalizing behaviors
1139 in Latino toddlers. *Ambulatory Pediatr.* 7(3):232–238. doi:
1140 10.1016/j.ambp.2007.02.004.

1141 Tonge KL, Jones RA, Hagenbuchner M, Nguyen TV, Okely AD. (2017). Educator
1142 engagement and interaction and children's physical activity in early childhood
1143 education and care settings: an observational study protocol. *BMJ Open.*
1144 7;7(2):e014423. doi: 10.1136/bmjopen-2016-014423.

1145 Toselli S, Zaccagni L, Celenza F, Albertini A, Gualdi-Russo E. (2015). Risk factors of
1146 overweight and obesity among preschool children with different ethnic background.
1147 *Endocrine.* 2015 Aug;49(3):717-25. doi: 10.1007/s12020-014-0479-4.

1148 Tremblay, M. S., Leblanc, A. G., Carson, V., Choquette, L., Connor Gorber, S., Dillman, C.,
1149 ... Timmons, B. W. (2012). Canadian physical activity guidelines for the early years
1150 (aged 0 to 4 years). *Applied Physiology, Nutrition, and Metabolism*, 37(2), 370-391.

1151 Tremblay, M. S., LeBlanc, A. G., Kho, M. E., Saunders, T. J., Larouche, R., Colley, R. C., ...
1152 Gorber, S. C. (2011). Systematic review of sedentary behaviour and health indicators
1153 in school-aged children and youth. *International Journal of Behavioral Nutrition and*
1154 *Physical Activity*, 8, 98. doi:10.1186/1479-5868-8-98

1155 Trost, S. G., Ward, D. S., & Senso, M. (2010). Effects of child care policy and environment
1156 on physical activity. *Medicine & Science in Sports & Exercise*, 42(2), 520-525.
1157 doi:10.1249/MSS.0b013e3181cea3ef .

1158 Tucker P, Vanderloo LM, Burke SM, Irwin JD, Johnson AM. (2015). Prevalence and
1159 influences of preschoolers' sedentary behaviors in early learning centers: a cross-
1160 sectional study. *BMC Pediatr.*, 15:128. doi: 10.1186/s12887-015-0441-5.

1161 Vale, S., Santos, R., Soares-Miranda, L., Silva, P., & Mota, J. (2010). Compliance with
1162 physical activity guidelines in preschool children. *Journal of Sports Science*,
1163 28(6), 603-608.

1164 Van Cauwenberghe, E., Jones, R. A., Hinkley, T., Crawford, D., & Okely, A. D. (2012).
1165 Patterns of physical activity and sedentary behavior in preschool children.
1166 *International Journal of Behavioral Nutrition and Physical Activity*, 27(9), 138.

1167 Vanderloo LM, Tucker P. (2015). Weekly trends in preschoolers' physical activity and
1168 sedentary time in childcare. *Int J Environ Res Public Health*,12(3):2454-64. doi:
1169 10.3390/ijerph120302454.

1170 Vanderloo LM, Tucker P, Johnson AM, Burke SM, Irwin JD. (2015) Environmental
1171 Influences on Preschoolers' Physical Activity Levels in Various Early-Learning
1172 Facilities. *Res Q Exerc Sport.*, 86(4):360-70. doi: 10.1080/02701367.2015.1053105.
1173 Epub 2015 Aug 19.

1174 Vanderloo, L. M., Martynuik, O. J. M., & Tucker, P. (2015). Physical and sedentary activity
1175 levels among preschoolers in home-based child care: A systematic review. *Journal of*
1176 *Physical Activity and Health*, 12(6), 879-889. doi:10.1123/jpah.2013-0483

1177 Vanderloo, L. M., Tucker, P., Johnson, A. A., Van Zandvoort, M. M., Burke, S. M., & Irwin,
1178 J. D. (2014). The influence of center-based child care on preschoolers' physical
1179 activity levels: A cross-sectional study. *International Journal of Environmental*
1180 *Research and Public Health*, 11(2), 1794-1802.

- 1181 Vandewater, E. A., Bickham, D. S., & Lee, J. H. (2006). Time well spent? Relating television
1182 use to children's free-time activities. *Pediatrics*, *117*(2), 181-191.
- 1183 Vandewater, E. A., Shim, M. S., & Caplovitz, A. G. (2004). Linking obesity and activity level
1184 with children's television and video game use. *Journal of Adolescents*, *27*(1), 71-85.
- 1185 Vandewater E. A., Park S.-E., Huang X., Wartella E. A. (2005). "No—you can't watch that":
1186 parental rules and young children's media use. *American Behavioral*
1187 *Scientist*. *48*(5):608–623. doi: 10.1177/0002764204271497.
- 1188 Vandewater EA, Rideout VJ, Wartella EA, Huang X, Lee JH, Shim MS. (2007). Digital
1189 childhood: electronic media and technology use among infants, toddlers, and
1190 preschoolers. *Pediatrics*. *119*(5):e1006–e1015. doi: 10.1542/peds.2006-1804.
- 1191 Vandewater EA, Bickham DS, Lee JH, Cummings HM, Wartella EA, Rideout VJ. (2005).
1192 When the television is always on - Heavy television exposure and young children's
1193 development. *Am Behav Sci*. *48*(5):562–577. doi: 10.1177/0002764204271496.
- 1194 Veitch, J., Hume, C., Salmon, J., Crawford, D., & Ball, K. (2013). What helps children be
1195 more active and less sedentary? Perceptions of mothers living in disadvantaged
1196 neighbourhoods. *Child Care Health Development*, *39*(1), 94-102.
- 1197 Veitch, J., Salmon, J., & Ball, K. (2010). Individual, social and physical environmental
1198 correlates of children active free-play: A cross-sectional study. *International Journal*
1199 *of Behavioral Nutrition and Physical Activity*, *2*, 7-11.
- 1200 Ward S, Bélanger M, Donovan D, Boudreau J, Vatanparast H, Muhajarine N, Leis
1201 A, Humbert ML, Carrier N. (2017). "Monkey see, monkey do": Peers' behaviors
1202 predict preschoolers' physical activity and dietary intake in childcare centers. *Prev*
1203 *Med*. *10*;97:33-39. doi: 10.1016/j.ypmed.2017.01.001.

1204 Zecevic, C. A., Tremblay, L., Lovsin, T., & Michel, L. (2010). Parental influence on young
1205 children's physical activity. *International Journal of Pediatrics*, 2010, 468-526.
1206 doi:10.1155/2010/468526.

1207 Zimmerman FJ, Christakis DA. (2005). Children's television viewing and cognitive
1208 outcomes: a longitudinal analysis of national data. *Arch Pediatr Adolesc*
1209 *Med.* 159(7):619–625. doi: 10.1001/archpedi.159.7.619.

1210 Zimmerman FJ, Christakis DA, Meltzoff AN. (2007). Television and DVD/video viewing in
1211 children younger than 2 years. *Arch Pediatr Adolesc Med.* 161(5):473–479. doi:
1212 10.1001/archpedi.161.5.473.

1213

1214

1215

1216

1217

1218