

O. I. Kalchenko, S. O. Cherenok, A. V. Solovyov*, S. Yu. Suikov, V. I. Kalchenko

Institute of Organic Chemistry, National Academy of Sciences of Ukraine

Murmanska str., 5, Kyiv-94, 02660, Ukraine. E-mail: ioch@ioch.kiev.ua

* Department of Chemical and Biomolecular Engineering, University of California, Berkeley, USA, CA 94720-1460

The study of calixarenes complexation with phenols by RP HPLC

The Host-Guest complexation of octakis(diphenoxyphosphoryloxy)tetramethylcalix[4]resorcinarene, 5,17-bis-(N-tolyliminomethyl)-25,27-dipropoxycalix[4]arene and 5,11,17,23-tetrakis(diisopropoxyphosphonyl)-25,26,27,28-tetrapropoxycalix[4]arene with a series of 11 phenols (phenol, p-fluorophenol, p-chlorophenol, p-bromophenol, pyrogallol, p-cresol, p-aminophenol, p-nitrophenol, salicylic aldehyde, guaiacol and veratrole) has been studied by the high-performance liquid chromatography (RP HPLC) method. Chromatographic characteristics and $\log P$ of industrial phenols have been determined. Using the relationship of the phenol retention factor k' vs the calixarene concentration in the mobile phase the stability constants of the supramolecular complexes K_A (29-331 M^{-1}) have been determined. The stability constants of the calixarene complexes show that the Host-Guest interaction strongly depends on the nature of the substituents in the Host and Guest molecules. Calixresorcinarene functionalized by diphenoxyphosphoryl groups and calixarene containing tolyliminomethyl groups formed more stable complexes with some phenols compared to calixarene functionalized by diisopropoxyphosphonyl groups. In accordance with the molecular modeling data the complexation does not change the C_{2v} flattened-cone conformation of the calixarene skeleton. The Host-Guest complexes are stabilized by the intermolecular hydrogen bonds of phenolic OH groups with oxygen atoms of $P=O$ groups at the upper rim, and OH groups at the lower rim of the macrocycle. Hydrophobic interactions also participate in the complexation.

Key words: Calixarenes; phenols; inclusion complexes; binding constants; molecular modeling

O. I. Кальченко, С. О. Черенок, А. В. Соловйов, С. Ю. Суйков, В. І. Кальченко Дослідження комплексоутворення каліксаренів з фенолами методом ОФ ВЕРХ

Комплексоутворення типу гість-господар октакіс-(дифеноксифосфорилокси)-тетраметилкалікс[4]резорцинарену, 5,17-біс-(N-толілімінометил)-25,27-дипропоксикалікс[4]арену та 5,11,17,23-тетракіс(діізопропоксифосфоніл)-25,26,27,28-тетрапропоксикалікс[4]арену з низкою 11 фенолів (фенол, p-фторофенол, p-хлорофенол, p-бромофенол, пірогалол, p-кресол, p-амінофенол, p-нітрофенол, саліциловий альдегід, гваякол, вератрол) було досліджено методом високоефективної рідинної хроматографії (ОФ ВЕРХ). Визначені хроматографічні характеристики фенолів та розраховано значення їх ліпофільності $\log P$. З урахуванням залежності коефіцієнтів утримання фенолів k' від концентрації каліксаренів у рухомій фазі визначені константи стійкості їх супрамолекулярних комплексів K_A , які знаходяться в межах 29-331 M^{-1} . Значення констант стійкості комплексів вказують на те, що взаємодія типу гість-господар значною мірою зумовлена природою замісників у молекулах гостя та господаря. Каліксрезорцинарен, функціоналізований дифеноксифосфорилокси-групами, і каліксарен, що містив толіліміно-групи, утворювали стійкіші комплекси з деякими фенолами порівняно з каліксареном, що містив діізопропоксифосфонільні групи. За даними молекулярного моделювання комплексоутворення не змінює конформацію каліксаренового кістяка C_{2v} сплющений конус. Комплекси типу гість-господар стабілізуються міжмолекулярними водневими зв'язками фенольних ОН-груп з атомами кисню $P=O$ груп верхнього вінця і ОН-груп нижнього вінця макроциклу. Гідрофобні взаємодії також відіграють роль у комплексоутворенні.

Ключові слова: каліксарени; феноли; комплекси включення; константи зв'язування; молекулярне моделювання

O. I. Кальченко, С. О. Черенок, А. В. Соловьев, С. Ю. Суйков, В. И. Кальченко Исследование комплексообразования каликсаренов с фенолами методом ОФ ВЭЖХ

Комплексообразование типа гость-хозяин октакис-(дифеноксифосфорилокси)-тетраметилкаликс[4]резорцинарена, 5,17-бис-(N-толилиминуметил)-25,27-дипропоксикаликс[4]арена и 5,11,17,23-тетракис диизопропоксифосфонил)-25,26,27,28-тетрапропоксикаликс[4]арена с серией 11 фенолов (фенол, p-фторфенол, p-хлорфенол, p-бромфенол, пирагаллол, p-кресол, p-аминофенол, p-нитрофенол, салициловый альдегид, гваякол, вератрол) было исследовано методом высокоэффективной жидкостной хроматографии (ОФ ВЭЖХ). Определены хроматографические характеристики фенолов и рассчитаны значения их липофильности $\log P$. С использованием зависимости коэффициентов удерживания фенолов k' от концентрации каликсаренов в подвижной фазе определены константы устойчивости их супрамолекулярных комплексов K_A , которые находятся в пределах 29-331 M^{-1} . Значения констант устойчивости показали, что взаимодействие типа гость-хозяин в значительной мере зависит от природы заместителей в молекулах Гостя и Хозяина. Каликсрезорцинарен, функционализированный дифеноксифосфорилокси-группами, и каликсарен, содержащий толилиминуметильные группы, давали более устойчивые комплексы с некоторыми фенолами, чем каликсарен с диизопропоксифосфонильными группами. Согласно данным молекулярного моделирования комплексообразование не приводит к изменению конформации каликсаренового скелета уплощенный конус C_{2v} . Комплексы типа гость-хозяин стабилизируются межмолекулярными водородными связями фенольных ОН-групп с атомами кислорода $P=O$ групп верхнего обода макроцикла и ОН-групп его нижнего обода. Гидрофобные взаимодействия также играют роль в комплексообразовании.

Ключевые слова: каликсарены; фенолы; комплексы включения; константы связывания; молекулярное моделирование

Calixarenes [1], bowl-shaped macrocyclic compounds are versatile platforms in the design of molecular receptors capable to recognize, bind and separate different organic and inorganic molecules similar in size and properties [2]. Calixarenes are able to recognize amino acids [3, 4, 5], dipeptides [6], nucleotides, nucleosides [7] and environmentally hazardous molecules [8, 9, 10]. Apparently, the outstanding receptor properties of functionalized calixarenes toward different (bio)molecules make them highly promising Hosts for sensor technologies [11, 12, 13] or for bio-medical research [14, 15, 16].

The class of phenols is important as a component of many useful industrial materials, pharmaceuticals, herbicides, disinfectants, etc. Phenols are essential raw material and additives for laboratory processes, chemical industry, chemical engineering processes, wood and plastics processing [17].

Phenolic compounds are commonly synthesized industrially; however, they are also produced by plants and microorganisms [18]. Organisms that synthesize phenolic compounds do so in response to ecological pressures, such as pathogen and insect attack, UV radiation and wounding [19]. Phenols are found in the natural world, especially in the plant kingdom, such as fruits, vegetables, herbs, spices and unfiltered olive oil [20].

In animals and humans, after ingestion, natural phenols become part of the xenobiotic metabolism. These activated metabolites are conjugated with charged species, such as glutathione, sulfate, glycine or glucuronic acid. As they are present in food consumed in human diets and in plants used in traditional medicine of several cultures, their role in human health and disease is a subject of research. Some natural phenols can be used as biopesticides [21, 22].

Phenols (aspirin and paracetamol, first of all) are widely used as inflammatory drugs [23]. Crofelemer is applied for the treatment of diarrhea. Phenolic compound Combretastatin is active as an anticancer drug [24].

In this paper the results of studying the Host-Guest complexation of octakis(diphenoxyphosphoryloxy)-tetramethylcalix[4]resorcinarene (**CA-1**), 5,17-bis-(N-tolyliminomethyl)-25,27-dipropoxycalix[4]arene (**CA-2**) and 5,11,17,23-tetrakis(diisopropoxyphosphonyl)-25,26,27,28-tetrapropoxycalix[4]arene (**CA-3**) with a series of phenols (phenol **1**, *p*-fluorophenol **2**, *p*-chlorophenol **3**, *p*-bromophenol **4**, pyrogallol **5**, *p*-cresol **6**, *p*-aminophenol **7**, *p*-nitrophenol **8**, salicylic aldehyde **9**, guaiacol **10**, veratrole **11**) (Fig. 1) in the acetonitrile-water solution by the RP HPLC methods are presented. The stability constants of the complexes were determined. The mechanism of the Host-Guest complexation was studied by the molecular modeling method. The information on the supramolecular Host-Guest interaction of calixarenes with phenol derivatives can be useful for design of chemosensors and drug carriers.

Experimental Part

CA-1, CA-2, CA-3 were synthesized by the methods [25, 26, 27], respectively. Acetonitrile was obtained from Acros Organics (Thermo Fisher Scientific, New Jersey – USA), and phenols were purchased from Sigma-Aldrich (St. Louis, MO, USA).

RP HPLC analysis

RP HPLC analysis was performed using the liquid chromatographic system (Hitachi, Ltd., Tokyo, Japan). The column (250 × 4.6 mm i.d.) was packed with LiChrosorb RP-18 (Merck, Darmstadt, Germany). The UV detector was operated at 254 nm. The flow rate was 0.8 ml/min. Experiments were performed with isocratic conditions. All chromatograms were obtained at 26 °C. The mixture of acetonitrile–water (80 : 20, v/v) was used as a mobile phase. The mobile phases contained calixarene additives in the concentrations of 0.05–0.6 mM. The samples of analytes for injections were dissolved in the same mixture (*C* = 0.01 mM) of acetonitrile–water (80 : 20, v/v). The volume of the samples injected was 20 μL. For equilibration of the chromatographic column the mobile phase with the calixarene additive was fluxed for 3 h before analysis. Under these conditions the column was saturated with calixarene.

The stability constants K_A of the calixarene complexes with phenols were determined by the RP HPLC method described in [28]. The method is based on changing the phenols retention factor k' induced by the complexation with calixarenes in the mobile phase.

Determination of the $\log P$ values of phenols

The values of $\log P$ of phenols **1-11** (Tab. 1) were calculated from equation 1:

$$\log P = 2.11 \cdot (\log k'), \quad (1)$$


where the coefficient 2.11 is the ratio of the experimental value of $\log P$ of phenol 1.46 [29] to $\log k'$ of the phenol value of 0.69 determined by the RP HPLC method in this work. The reason of the experimental determination of lipophilicity was a mismatch of the literature data of the $\log P$ values of the phenols studied. For example, $\log P$ of *p*-nitrophenol **9** is 1.96 according to [30] and 1.68 according to [31].

Molecular modeling

Molecular modeling of calixarenes and their Host-Guest complexes were carried out using the HyperChem 8.0 program in the force field (PM3) [32]. The structures were calculated by the semi-empirical method. The RMS (the standard deviation of the word root mean square) gradient was equal to 0.01 kcal/Å mol.

Results and Discussion

Calixarenes were registered on the chromatograms by sharp peaks with retention factors 8.65 (**CA-1**), 0.89 (**CA-2**) and 2.38 (**CA-3**). As shown, addition of

Fig. 1. Calix[4]arenes **CA-1**, **CA-2** and **CA-3** (Hosts) and phenols **1-11** (Guests)**Table 1**

The values of $\log P$ of phenols **1-11**, the stability constants K_A (M^{-1}) (RSD = 9-15 %) and free Gibbs energies ΔG (kJ/mol) of their Host-Guest complexes with **CA-1**, **CA-2**, **CA-3**

No.	Guest	CA-1		CA-2		CA-3		log P
		K_A	ΔG	K_A	ΔG	K_A	ΔG	
1	Phenol	123	-11.9	211	-13.24	33	-8.65	1.46
2	<i>p</i> -Fluorophenol	102	-11.44	141	-12.24	140	-12.22	1.27
3	<i>p</i> -Chlorophenol	92	-11.18	240	-13.56	36	-8.86	1.53
4	<i>p</i> -Bromophenol	119	-11.82	235	-13.50	29	-8.33	1.49
5	Pyrogallol	116	-11.76	160	-12.55	nd	nd	1.40
6	<i>p</i> -Cresol	94	-11.24	216	-13.29	nd	nd	1.49
7	<i>p</i> -Aminophenol	306	-14.16	230	-13.45	331	-14.30	1.47
8	<i>p</i> -Nitrophenol	166	-12.64	220	-13.34	34	-8.72	1.47
9	Salicylic aldehyde	40	-9.12	141	-12.24	94	-11.24	1.04
10	Guaiacol	209	-13.21	324	-14.3	130	-12.04	2.22
11	Veratrole	80	-10.84	246	-13.62	263	-13.78	1.53




Fig. 2. Plots of $1/k'$ for phenols vs the **CA-1** concentration in the mobile phase ($r = 0.89-0.98$)

calixarene to the mobile phase decreases the retention factors k' of phenols **1-11** due to formation of the Host-Guest inclusion complexes. The linear character of plots of $1/k'$ vs the calixarene concentration (Fig. 2-4) indicates 1 : 1 stoichiometry of the Host-Guest supramolecular complexes formed. It allows using equation (2) correctly for calculation of the stability constants K_A :

$$1/k' = 1/k_0' + K_A \times [CA]/k_0' \quad (2)$$

where k_0' i k' – are capacity factors of the Guest molecule determined in the absence and the presence of the calixarene Host in the mobile phase.

The stability constants K_A of the calixarene complexes are within 29-331 M^{-1} (Tab. 1). This is close to the stability constant of the β -cyclodextrin – pyrogallol complex determined by the UV-visible ($K_A = 78 M^{-1}$) and fluorescence ($K_A = 227 M^{-1}$) spectroscopy in the water solution [33]. Free Gibbs energies DG of the cali-




Fig. 3. Plots of $1/k'$ for phenols vs the **CA-2** concentration in the mobile phase ($r = 0.97-0.99$)




Fig. 4. Plots of $1/k'$ for phenols vs the **CA-3** concentration in the mobile phase ($r = 0.93-0.96$)

xarene complexes calculated by equation (3) are from -8.33 to -14.30 kJ/mol (Tab. 1).

$$DG = -RT \cdot \ln K_A \quad (3)$$

To clarify the nature of the Host-Guest interaction the molecular modeling of the calixarene Hosts **CA-1**, **CA-2**, **CA-3** and their complexes with *p*-aminophenol and phenol was carried out (Fig. 5).

In accordance with the molecular modeling data the calixarene macrocyclic skeleton of **CA-1**, **CA-2** and **CA-3** and their Host-Guest complexes exist in the *flattened-cone* conformation with C_{2v} symmetry. Two aromatic rings in the distal position are “coplanar” oriented, but two other ones are “perpendicular” oriented to the main plane of the macrocycle. The dihedral angles characterized inclination of benzene rings A, B, C, D relatively to the main macrocycle plane formed by CH_2 links are presented in Tab. 2. As shown from Tab. 2, the complexation with phenols does not practically change dihedral angles in the calixarene macrocyclic skeleton.

In all complexes formation of the intermolecular hydrogen bonds between OH groups of phenols and oxygen atoms of calixarenes are observed ($H \cdots O$ distances are within 2.91-3.19 Å). In complex (a) OH and NH_2 groups of *p*-aminophenol form the hydrogen bonds with oxygen atoms of distal P = O groups of **CA-1**. In complex (b) phenol forms the hydrogen bond with the oxygen atom of the OH group at the lower rim of calixarene **CA-2**. In complex (c) phenol forms the hydrogen bond with the P = O group at the upper rim of the macrocycle **CA-3**.

Short contacts between calixarenes and phenol molecules (Fig. 5) indicate that the Host-Guest complexes can be additionally stabilized by various non-valence Van der Waals, π - π , and hydrophobic interactions. Fig. 6 presents the linear dependences of the




Fig. 5. Energy minimized structures of the Host-Guest complexes: **CA-1** with *p*-aminophenol (a), **CA-2** and **CA-3** with phenol (b, c). Phenyl substituents at the upper rim of **CA-1** are omitted for clarity. Hydrogen bonds are shown as dotted lines




Fig. 6. Plots $\log K_A$ of **CA-1** ($r = 0.60$), **CA-2** ($r = 0.92$) and **CA-3** ($r = 0.96$) of the complexes on $\log P$ of phenols 1-8

stability constants K_A of **CA-1**, **CA-2** and **CA-3** from $\log P$ of the phenol derivatives confirming the role of the hydrophobic interaction in the process of the Host-Guest complexation.

Conclusions


Calix[4]arenes form stable supramolecular Host-Guest complexes with phenols in acetonitrile-water solutions. The stability constants of the complexes determined by the RP HPLC method show that the complex stability is strongly dependent on the nature of the substituents in the Host and Guest molecules. In accordance with the molecular modeling the complexation does not change the C_{2v} flattened-cone conformation of the calixarene skeleton. The Host-Guest

References

- Gutsche, C. D. Calixarenes revisited. Monographs in supramolecular chemistry / C. D. Gutsche; ed. J. F. Stoddart. – Cambridge, 1998. doi: 10.1039/9781847550293.
- Calixarenes and beyond / eds. P. Neri, J. L. Sessler, M.-X. Wang. – Springer, 2016. doi: 10.1007/978-3-319-31867-7.
- Complexation of calix[4]arenehydroxymethyl-phosphonic acids with amino acids. Binding constants determination of the complexes by HPLC method / O. Kalchenko, S. Cherenok, O. Yushchenko, V. Kalchenko // J. Incl. Phenom. – 2012. – Vol. 76, Issue 1-2. – P. 29-36. doi: 10.1007/s10847-012-0169-x.
- Kalchenko, O. I. A comparative study of the determination of the stability constants of inclusion complexes of *p*-sulphonatocalix[4]arene with aminoacids by RP HPLC and ^1H NMR / O. I. Kalchenko, F. Perret, A. W. Coleman // J. Chem. Soc., Perkin Trans. 2. – 2001. – Issue 3. – P. 258-263. doi: 10.1039/b005497f.
- Recognition of amino acids by functionalized calixarenes / L. Muthiac, J. H. Lee, J. S. Kim, J. Vicens // Chem. Soc. Rev. – 2011. – Vol. 40, Issue 5. – P. 2777. doi: 10.1039/c0cs00005a.
- Binding of dipeptides and tripeptides containing lysine or arginine by *p*-sulphonatocalixarenes in water: NMR and microcalorimetric studies / N. Douteau-Guevel, F. Perret, A. W. Coleman et al. // J. Chem. Soc. Perkin Trans. 2. – 2002. – Issue 3. – P. 524-532. doi: 10.1039/b109553f.
- Sansone, F. Calixarenes in bioorganic and biomimetic chemistry / F. Sansone, M. Segura, R. Ungaro // Calixarenes 2001. – Springer, 2001. – P. 496-512. doi: 10.1007/0-306-47522-7_27.
- Complexation of calix[4]arenephosphonous acids with 2,4-dichlorophenoxyacetic acid and atrazine in water / O. I. Kalchenko, A. V. Solovyov, S. A. Cherenok et al. // J. Incl. Phenom. – 2003. – Vol. 46, Issue 1. – P. 19-25.

Table 2

The inclination (dihedral angles) of benzene rings A, B, C, D in calixarenes **CA-1**, **CA-2**, **CA-3** and their complexes with *p*-aminophenol and phenol



Calixarenes or their Host-Guest complexes	Dihedral angles, (°)			
	A	B	C	D
CA-1	149	103	148	107
CA-1 – <i>p</i> -aminophenol	148	115	147	104
CA-2	148	91	148	91
CA-2 – phenol	147	101	146	101
CA-3	140	88	128	94
CA-3 – phenol	137	99	130	102

complexes are stabilized by the intermolecular hydrogen bonds of phenolic OH groups with oxygen atoms of P = O groups at the upper rim of the macrocycle and OH groups at the lower rim of the macrocycle. Hydrophobic interactions also participate in the Host-Guest complexation processes.

Conflicts of Interest: authors have no conflict of interest to declare.

9. Kalchenko, V. Calixarene receptors of environmentally hazardous and bio-relevant molecules and ions / V. Kalchenko // *Pure Appl. Chem.* – 2008. – Vol. 80, Issue 7. – P. 1449–1458. doi: 10.1351/pac200880071449.
10. Complexation and extraction of non-ferrous metals by calix[n]arene phosphine oxides. / G. A. Kostin, T. V. Us, T. M. Korda et al. // *J. Incl. Phenom.* – 2010. – Vol. 68, Issue 1–2. – P. 131–137. doi: 10.1007/s10847-010-9755-y.
11. Diamond, D. Peer reviewed: calixarenes: designer ligands for chemical sensors / D. Diamond, K. Nolan // *Anal. Chem.* – 2001. – Vol. 73, Issue 1. – P. 22A–29A. doi: 10.1021/ac012376g.
12. Calix[n]arenes as Protein Sensors / A. W. Coleman, F. Perret, M. Moussa et al. // *Top. Curr. Chem.* – 2007. – Vol. 277. – P. 31–88.
13. Zadnarm, R. Nanomolar protein sensing with embedded receptor molecules / R. Zadnarm, T. Schrader // *J. Amer. Chem. Soc.* – 2005. – Vol. 127, Issue 3. – P. 904–915. doi: 10.1021/ja045785d.
14. Perret, F. Biochemistry of the para-sulfonato-calix[n]arenes / F. Perret, A. N. Lazar, A. W. Coleman // *Chem. Commun.* – 2006. – Issue 23. – P. 2425. doi: 10.1039/b600720c.
15. Rodik, R. V. Calixarenes in bio-medical researches / R. V. Rodik, V. I. Boyko, V. I. Kalchenko // *Curr. Med. Chem.* – 2009. – Vol. 16, Issue 13. – P. 1630–1655. doi: 10.2174/092986709788186219.
16. Da Silva, E. Biopharmaceutical applications of calixarenes / E. Da Silva, A. N. Lazar, A. W. Coleman // *J. Drug Deliv. Sci. and Technol.* – 2004. – Vol. 14, Issue 1. – P. 3–20. doi: 10.1016/s1773-2247(04)50001-1.
17. Weber, M. Phenol. Ullmann's encyclopedia of industrial chemistry / M. Weber, M. Kleine-Boymann. – Wiley-VCH, 2004. doi: 10.1002/14356007.a19_299.pub2.
18. Hättenschwiler, S. The role of polyphenols in terrestrial ecosystem nutrient cycling / S. Hättenschwiler, P. M. Vitousek // *Trends in Ecology & Evolution.* – 2000. – Vol. 15, Issue 6. – P. 238–243. doi: 10.1016/s0169-5347(00)01861-9.
19. Klepacka, J. Phenolic compounds as cultivar- and variety-distinguishing factors in some plant products / J. Klepacka, E. Gujska, J. Michalak // *Plant Foods for Human Nutrition.* – 2011. – Vol. 66, Issue 1. – P. 64–69. doi: 10.1007/s11130-010-0205-1.
20. Gutfinger, T. Polyphenols in olive oils / T. Gutfinger // *J. of the American Oil Chemists Society.* – 1981. – Vol. 58, Issue 11. – P. 966–968. doi: 10.1007/bf02659771.
21. Mishra, B. B. Natural products: an evolving role in future drug discovery / B. B. Mishra, V. K. Tiwari // *Eur. J. Med. Chem.* – 2011. – Vol. 46, Issue 10. – P. 4769–4780. doi: 10.1016/j.ejmech.2011.07.057.
22. Wildman, R. E. C. Handbook of nutraceuticals and functional foods / R. E. C. Wildman. – 2nd ed. – CRC Press, 2006. doi: 10.1201/9781420006186.
23. Frank, E. Paracetamol: a curriculum resource / E. Frank. – Cambridge, 2002. – P. 764.
24. Lead identification of conformationally restricted β -lactam type combretastatin analogues: Synthesis, antiproliferative activity and tubulin targeting effects / M. Carr, L. M. Greene, A. J. S. Knox et al. // *Eur. J. Med. Chem.* – 2010. – Vol. 45, Issue 12. – P. 5752–5766. doi: 10.1016/j.ejmech.2010.09.033.
25. Study of the complexation of octakis(diethoxyphosphoryloxy)-tetramethylcalix[4]resorcinarene with benzene derivatives by RP HPLC method / O. I. Kalchenko, A. V. Solovyov, J. Lipkowski, V. I. Kalchenko // *J. Incl. Phenom.* – 1999. – Vol. 34, Issue 3. – P. 259–266.
26. Upper rim alpha-hydroxy of alpha-aminophosphonic acid derivatives of calix[4]arenes / L. N. Markovsky, V. I. Kalchenko, A. V. Solovyov et al. // *Anales de Quimica.* – 1998. – Vol. 94, Issue 3. – P. 164–170.
27. Synthesis and complexation of amphiphilic calix[4]arene phosphonates with organic molecules in solutions and Langmuir-Blodgett films / A. V. Solovyov, S. O. Cherenok, O. I. Kalchenko et al. // *J. Mol. Liq.* – 2011. – Vol. 159, Issue 2. – P. 117–123. doi: 10.1016/j.molliq.2010.12.007.
28. Host-Guest interactions of calix[4]resorcinarenes with benzene derivatives in conditions of reversed-phase high-performance liquid chromatography / J. Lipkowski, O. I. Kalchenko, J. Slowikowska et al. // *J. Phys. Org. Chem.* – 1998. – Vol. 11, Issue 6. – P. 426–437. doi: 10.1002/(sici)1099-1395(199806)11:6<426::aid-poc963>3.3.co;2-i.
29. Hansch, C. Exploring QSAR: hydrophobic, electronic, and steric constants (ACS Professional Reference Book) / C. Hansch, A. Leo, D. H. Hoekman. – American Chemical Society, 1995. – 348 p.
30. Garst, J. Accurate, wide-range, automated, high-performance liquid chromatographic method for the estimation of octanol/water partition coefficients I: Effect of chromatographic conditions and procedure variables on accuracy and reproducibility of the method / J. Garst, E. Wilson // *J. Pharm. Sci.* – 1984. – Vol. 73, Issue 11. – P. 1616–1623. doi: 10.1002/jps.2600731133.
31. Unger, S. H. Simple procedure for determining octanol- aqueous partition coefficients, distribution and ionization coefficients with reverse phase high performance liquid chromatography / S. H. Unger, J. R. Cook, J. S. Hollenberg // *J. Pharm. Sci.* – 1978. – Vol. 67, Issue 10. – P. 1364–1367. doi: 10.1002/jps.2600671008.
32. HyperChem Community. – Available at : <http://www.hyper.com/Download/AllDownloads/tabid/470/Default.aspx>.
33. Inclusion complex of 1,2,3-trihydroxybenzene with α - and β -cyclodextrin / T. Stalin et al. // *Indian Journal of Chemistry.* – 2006. – Vol. 45, Issue 5. – P. 1113–1120.

References

1. Gutsche, C. D. (1998). *Calixarenes revisited. Monographs in supramolecular chemistry.* J. F. Stoddart (Ed.). Cambridge. doi: 10.1039/9781847550293.
2. Neri, P., Sessler, J. L., Wang, M.-X. (Eds.). (2016). *Calixarenes and Beyond.* Springer. doi: 10.1007/978-3-319-31867-7.
3. Kalchenko, O., Cherenok, S., Yushchenko, O., Kalchenko, V. (2012). Complexation of calix[4]arenehydroxymethylphosphonic acids with amino acids. Binding constants determination of the complexes by HPLC method. *Journal of Inclusion Phenomena and Macrocyclic Chemistry*, 76 (1–2), 29–36. doi: 10.1007/s10847-012-0169-x.
4. Kalchenko, O. I., Perret, F., Morel-Desrosiers, N., Coleman, A. W. (2001). A comparative study of the determination of the stability constants of inclusion complexes of p-sulfonatocalix[4]arene with amino acids by RP-HPLC and ¹H NMR. *Journal of the Chemical Society, Perkin Transactions 2*, 3, 258–263. doi: 10.1039/b005497f.
5. Mutihac, L., Lee, J. H., Kim, J. S., Vicens, J. (2011). Recognition of amino acids by functionalized calixarenes. *Chemical Society Reviews*, 40 (5), 2777. doi: 10.1039/c0cs00005a.
6. Douteau-Guével, N., Perret, F., Coleman, A. W., Morel, J.-P., Morel-Desrosiers, N. (2002). Binding of dipeptides and tripeptides containing lysine or arginine by p-sulfonatocalixarenes in water: NMR and microcalorimetric studies. Electronic supplementary information (ESI) available: chemical shifts experienced by different protons of KK, RR, KKK or RRR in the presence of increasing amounts of 14 or 16; heat effects observed upon titration of 14, 16 or 18 by KK, RR, KKK or RRR; COSY and ROESY 2D ¹H NMR spectra for 14 complexed with KK or RR. See <http://www.rsc.org/suppdata/p2/b1/b109553f/>. *Journal of the Chemical Society, Perkin Transactions 2*, 3, 524–532. doi: 10.1039/b109553f.
7. Sansone, F., Segura, M., Ungaro, R. (2001). *Calixarenes in Bioorganic and Biomimetic Chemistry.* Calixarenes 2001. Springer, 496–512. doi: 10.1007/0-306-47522-7_27.
8. Kalchenko, O. I., Solovyov, A. V., Cherenok, S. A., Starodub, N. F., Kalchenko, V. I. (2003). Complexation of calix[4]arenephosphonous acids with 2,4-dichlorophenoxyacetic acid and atrazine in water. *Journal of Inclusion Phenomena and Macrocyclic Chemistry*, 46 (1), 19–25.
9. Kalchenko, V. I. (2008). Calixarene receptors of environmentally hazardous and biorelevant molecules and ions. *Pure and Applied Chemistry*, 80 (7). doi: 10.1351/pac200880071449.
10. Kostin, G. A., US, T. V., Korda, T. M., Torgov, V. G., Kuratieva, N. V., Miroshnichenko, S. I., Kalchenko, V. I. (2010). Complexation and extraction of non-ferrous metals by calix[n]arene phosphine oxides. *Journal of Inclusion Phenomena and Macrocyclic Chemistry*, 68 (1–2), 131–137. doi: 10.1007/s10847-010-9755-y.

11. Diamond, D., Nolan, K. (2001). Peer Reviewed: Calixarenes: Designer Ligands for Chemical Sensors. *Analytical Chemistry*, 73 (1), 22A–29A. doi: 10.1021/ac012376g.
12. Coleman, A. W., Perret, F., Moussa, M., Dupin, M., Guo, Y., Perron, H. (2007). Calix[n]arenes as Protein Sensors. *Topics in Current Chemistry*, 277, 31–88.
13. Zadnard, R., Schrader, T. (2005). Nanomolar Protein Sensing with Embedded Receptor Molecules. *Journal of the American Chemical Society*, 127 (3), 904–915. doi: 10.1021/ja045785d.
14. Perret, F., Lazar, A. N., Coleman, A. W. (2006). Biochemistry of the para-sulfonato-calix[n]arenes. *Chemical Communications*, 23, 2425. doi: 10.1039/b600720c.
15. Rodik, R., Boyko, V., Kalchenko, V. (2009). Calixarenes in Bio-Medical Researches. *Current Medicinal Chemistry*, 16 (13), 1630–1655. doi: 10.2174/092986709788186219.
16. Da Silva, E., Lazar, A. N., Coleman, A. W. (2004). Biopharmaceutical applications of calixarenes. *Journal of Drug Delivery Science and Technology*, 14 (1), 3–20. doi: 10.1016/s1773-2247(04)50001-1.
17. Weber, M., Weber, M., Kleine-Boymann, M. (2004). *Phenol. Ullmann's Encyclopedia of Industrial Chemistry*. Wiley-VCH. doi: 10.1002/14356007.a19_299.pub2.
18. Hättenschwiler, S., Vitousek, P. M. (2000). The role of polyphenols in terrestrial ecosystem nutrient cycling. *Trends in Ecology & Evolution*, 15 (6), 238–243. doi: 10.1016/s0169-5347(00)01861-9.
19. Klepacka, J., Gujska, E., Michalak, J. (2011). Phenolic Compounds as Cultivar- and Variety-distinguishing Factors in Some Plant Products. *Plant Foods for Human Nutrition*, 66 (1), 64–69. doi: 10.1007/s11130-010-0205-1.
20. Gutfinger, T. (1981). Polyphenols in olive oils. *Journal of the American Oil Chemists Society*, 58 (11), 966–968. doi: 10.1007/bf02659771.
21. Mishra, B. B., Tiwari, V. K. (2011). Natural products: An evolving role in future drug discovery. *European Journal of Medicinal Chemistry*, 46 (10), 4769–4807. doi: 10.1016/j.ejmech.2011.07.057.
22. Wildman, R. (Ed.). (2006). *Handbook of Nutraceuticals and Functional Foods. Modern Nutrition*. (2nd ed.). CRC Press. doi: 10.1201/9781420006186.
23. Frank, E. (2002). *Paracetamol: a curriculum resource*. Cambridge, 764.
24. Carr, M., Greene, L. M., Knox, A. J. S., Lloyd, D. G., Zisterer, D. M., Meegan, M. J. (2010). Lead identification of conformationally restricted β -lactam type combretastatin analogues: Synthesis, antiproliferative activity and tubulin targeting effects. *European Journal of Medicinal Chemistry*, 45 (12), 5752–5766. doi: 10.1016/j.ejmech.2010.09.033.
25. Kalchenko, O. I., Solovyov, A. V., Lipkowski, J., Kalchenko, V. I. (1999). Study of the complexation of octakis(diethoxyphosphoryloxy)-tetramethyl-calix[4]resorcinarene with benzene derivatives by RP HPLC method. *Journal of Inclusion Phenomena and Macrocyclic Chemistry*, 34 (3), 259–266.
26. Markovskiy, L. N., Kalchenko, V. I., Solovyov, A. V., Finocchiaro, P., Failla, S., Atamas, L. I., Consiglio, G., Tsybmal, I. F. (1998). Upper rim alpha-hydroxy of alpha-aminophosphonic acid derivatives of calix[4]arenes. *Anales de Quimica*, 94 (3), 164–170.
27. Solovyov, A. V., Cherenok, S. O., Kalchenko, O. I., Atamas, L. I., Kazantseva, Z. I., Koshets, I. A., Kalchenko, V. I. (2011). Synthesis and complexation of amphiphilic calix[4]arene phosphonates with organic molecules in solutions and Langmuir-Blodgett films. *Journal of Molecular Liquids*, 159 (2), 117–123. doi: 10.1016/j.molliq.2010.12.007.
28. Lipkowski, J., Kalchenko, O. I., Slowikowska, J., Kalchenko, V. I., Lukin, O. V., Markovskiy, L. N., Nowakowski, R. (1998). *Journal of Physical Organic Chemistry*, 11 (6), 426–435. doi: 10.1002/(sici)1099-1395(199806)11:6<426::aid-poc963>3.3.co;2-i.
29. Hansch, C., Leo, A., Hoekman, D. H. (1995). *Exploring QSAR: hydrophobic, electronic, and steric constants (ACS Professional Reference Book)*. American Chemical Society, 348.
30. Garst, J. E., Wilson, W. C. (1984). Accurate, Wide-Range, Automated, High-Performance Liquid Chromatographic Method for the Estimation of Octanol/Water Partition Coefficients I: Effect of Chromatographic Conditions and Procedure Variables on Accuracy and Reproducibility of the Method. *Journal of Pharmaceutical Sciences*, 73 (11), 1616–1623. doi: 10.1002/jps.2600731133.
31. Unger, S. H., Cook, J. R., Hollenberg, J. S. (1978). Simple Procedure for Determining Octanol-Aqueous Partition, Distribution, and Ionization Coefficients by Reversed-Phase High-Pressure Liquid Chromatography. *Journal of Pharmaceutical Sciences*, 67 (10), 1364–1367. doi: 10.1002/jps.2600671008.
32. HyperChem Community. Available at: <http://www.hyper.com/Download/AllDownloads/tabid/470/Default.aspx>.
33. Stalin, T., Vasantha Rani, P., Shanthi, B., Sekar, A., Rajendiran, N. (2006). Inclusion complex of 1,2,3-trihydroxybenzene with α - and β -cyclodextrin. *Indian Journal of Chemistry*, 45 (5), 1113–1120.

Надійшла до редакції 10.01.2017 р.

Acknowledgement

This work is supported by the State Fund for Fundamental Research of Ukraine.