

O. M. Agafonov¹, S. I. Okovytyy², M. Ye. Blazheyevskiy¹¹ National University of Pharmacy, Ukraine53, Pushkinska str., Kharkiv, 61002. E-mail: chuh2018@ukr.net² Oles Honchar Dnipro National University, Ukraine

The comparative quantum chemical study of the epoxidation reaction mechanism of eugenol and *isoeugenol* with peracetic and perbenzoic acids

Aim. To study the kinetics of the epoxidation reaction for eugenol and *isoeugenol* with perbenzoic acid, carry out the comparative quantum chemical study of the epoxidation reaction mechanism of eugenol and *isoeugenol* isomers (2-*cis* and 2-*trans*) with peracetic and perbenzoic acids.

Results and discussion. The kinetics of the epoxidation reaction of isomeric terpenoids eugenol and *isoeugenol* with perbenzoic acid in the medium of methylene chloride medium at 293 K was studied using the method of iodometric titration. It was shown that the rate constant of the epoxidation reaction for eugenol was in 5.5 times higher than for *isoeugenol*. According to the results of quantum chemical calculations using the UBH&HLYP/6-31G (d) approximation, the structures of transition states of eugenol and *isoeugenol* formed during the epoxidation reactions studied were proposed, and the activation energies for the corresponding reactions were calculated. Based on the results of the studies conducted it was found that the ratio of the activation energies during the interaction of eugenol and *isoeugenol* with peracetic and perbenzoic acids showed the higher reactivity of *isoeugenol*.

Experimental part. To study the kinetics of the epoxidation reaction the method of iodometric titration was used. The method of the functional density (software Gaussian 09, approximation UBH&HLYP/6-31G (d)) was applied for calculation.

Conclusions. The results of the quantum chemical study of the epoxidation reaction mechanism of eugenol and *isoeugenol* are consistent with the kinetic data experimentally obtained; it confirms the correctness of using the UBH&HLYP/6-31G (d) approximation for studying the features of epoxidation of isomeric terpenoids with organic peracids.

Key words: terpenoids; quantum chemical research; reaction mechanism; reaction kinetics; epoxidation; peracetic acid; perbenzoic acid

О. М. Агафонов¹, С. І. Оковитий², М. Є. Блажеєвський¹¹ Національний фармацевтичний університет, Україна² Дніпровський національний університет імені Олеся Гончара, Україна

Порівняльне квантово-хімічне дослідження механізму реакції епоксидування евгенолу та ізоєвгенолу надацетатною та надбензойною кислотами

Мета. Дослідити кінетику реакції епоксидування евгенолу та ізоєвгенолу надбензойною кислотою, провести порівняльне квантово-хімічне дослідження механізму реакції епоксидування евгенолу та двох ізомерів ізоєвгенолу (2-*цис* та 2-*транс*) надацетатною та надбензойною кислотами.

Результати та обговорення. Методом йодометричного титрування вивчено кінетику реакції епоксидування ізомерних терпеноїдів евгенолу та ізоєвгенолу надбензойною кислотою в середовищі метиленхлориду при 293 К і показано, що константа швидкості реакції епоксидування евгенолу в 5,5 разів вище, ніж для ізоєвгенолу. За результатами квантово-хімічних розрахунків з використанням наближення UBH&HLYP/6-31G (d) запропоновано структури перехідних станів евгенолу та ізоєвгенолу, що утворюються в ході вивчених реакцій епоксидування, і розраховано значення енергій активації для відповідних реакцій. Виходячи з результатів проведених досліджень встановлено, що співвідношення енергій активації при взаємодії евгенолу та ізоєвгенолу з надацетатною і надбензойною кислотами свідчить про більш високу реакційну здатність ізоєвгенолу.

Експериментальна частина. Для вивчення кінетики реакції епоксидування використовували метод йодометричного титрування. Для розрахунків використовували метод функціоналу густини (наближення UBH&HLYP/6-31G (d)) програми Gaussian 09.

Висновки. Результати квантово-хімічного дослідження механізму реакції епоксидування евгенолу та ізоєвгенолу узгоджуються з експериментально отриманими кінетичними даними, що підтверджує коректність використання наближення UBH&HLYP/6-31G (d) для вивчення особливостей епоксидування ізомерних терпеноїдів за допомогою органічних надкислот.

Ключові слова: терпеноїди; квантово-хімічне дослідження; механізм реакції; кінетика реакції; епоксидування; надацетатна кислота; надбензойна кислота

А. М. Агафонов¹, С. И. Оковитый², Н. Е. Блажеевский¹¹ Национальный фармацевтический университет, Украина² Днепропетровский национальный университет имени Олеся Гончара, Украина

Сравнительное исследование механизма реакции эпексидирования эвгенола и изоэвгенола перуксусной и пербензойной кислотами

Цель. Изучить кинетику реакции эпексидирования эвгенола и изоэвгенола пербензойной кислотой, провести сравнительное квантово-химическое исследование механизма реакции эпексидирования эвгенола и двух изомеров изоэвгенола (2-*цис* и 2-*транс*) перуксусной и пербензойной кислотами.

Результаты и обсуждение. Методом йодометрического титрования изучена кинетика реакции эпексидирования изомерных терпеноидов эвгенола и *изоэвгенола* пербензойной кислотой в среде хлористого метилена при 298 К и показано, что константа скорости реакции эпексидирования эвгенола в 5,5 раз выше, чем *изоэвгенола*. По результатам квантово-химических расчетов с использованием приближения UBH&HLYP/6-31G (d) предложены структуры переходных состояний эвгенола и *изоэвгенола*, образующиеся в ходе изученных реакций эпексидирования, и рассчитаны значения энергий активации для соответствующих реакций. Исходя из результатов проведенных исследований установлено, что соотношение энергий активации при взаимодействии эвгенола и *изоэвгенола* с перуксусной и пербензойной кислотами свидетельствует о более высокой реакционной способности *изоэвгенола*.

Экспериментальная часть. Для изучения кинетики реакции эпексидирования использовали метод йодометрического титрования. Для расчета использовали метод функционала плотности (приближение UBH&HLYP/6-31G (d)) программы Gaussian 09.

Выводы. Результаты квантово-химического исследования механизма реакции эпексидирования эвгенола и *изоэвгенола* согласуются с экспериментально полученными кинетическими данными, что подтверждает корректность использования приближения UBH&HLYP/6-31G (d) для изучения особенностей эпексидирования изомерных терпеноидов с помощью органических перкислот.

Ключевые слова: терпеноиды; квантово-химическое исследование; механизм реакции; кинетика реакции; эпексидирование; перуксусная кислота; пербензойная кислота

The study of transformation reaction mechanisms is increasingly important to both among theoretic and experimental chemists. By the means of quantum chemical calculations one may predict the nature, character of interaction, energy and geometric parameters of a transition state and end products of a reaction [1–3]. The specified density functional theory is used to study the electronic and geometric structure and relative stability of different compounds [1–5]. The B3LYP functional is the most extensively used among all hybrid density functional methods; it allows describing correctly biradical structures and is quite economic in terms of cost of computer time, which allows its use in the study of sufficiently complex organic compounds and reactions [4–5].

Eugenol (**1**) is a natural compound obtained from various plants, particularly from black cloves. Although its consumption in a small amount provides antibacterial and antifungal benefits, exceeding the limit may cause various harmful effects [6–11]. Eugenol-rich cloves have been used for food preservation and in various medical areas, owing to its antibacterial, anti-septic and anti-inflammatory properties [6–14]. Eugenol sources, such as clove oil, cinnamon, basil and nutmeg oil, are some of the basic ingredients in mouth washes, soaps, tooth pastes, perfumes and in various veterinary medications [15]. Moreover, eugenol antibacterial and antifungal activities are very helpful in treating many digestive problems, such as diarrhea, gas bloating, ulcers and Candida [6–11]. Eugenol-rich black paper and clove can increase the release of gastric acid, which helps in digesting food and kills intestinal bacteria in the stomach. Clove tea can be used to dilate blood vessels, increase the circulation and the body temperature. As an antioxidant, eugenol can prevent or decrease harmful oxidations in the body [16–17].

Although eugenol has been nominated to be safe to consume in suggested limits by Food and Drug Administration, exceeding the limits may cause serious

and fatal results, such as arrhythmia, kidney damage, digestive problems, increase in heart rate and blood pressure, dizziness, vomiting and liver failure [18]. Eugenol is quickly absorbed and digested in the liver, and 95 % of the dose is excreted within one day. However, it is reported that ingesting clove oil as small as 8 mL can cause fatal sickness by damaging the liver and nervous system [19]. The study of the eugenol epoxidation reaction mechanism of various isomers with organic peracids is of interest to scientists [20].


We earlier studied the epoxidation reaction mechanism of eugenol and *isoeugenol* with peracetic acid [21].

The aim of this article was to study the kinetics of the epoxidation reaction for eugenol (**1**) and isomers of *isoeugenol* – *cis*-2 (**2**) and *trans*-2 (**3**) – with perbenzoic acid and carry out the comparative quantum chemical study of the epoxidation reaction mechanisms of eugenol and *isoeugenol* with peracetic and perbenzoic acids (PBA) (Fig. 1).

The kinetic study of epoxidation reactions of both terpenoids with perbenzoic acid (PBA) was carried out in methylene chloride as a solvent in the second-order conditions under the temperature of 293 K. The reactions were studied by quantification of the unreacted PBA as a function of time using the method of iodometric titration. Fig. 2 shows the kinetic curves of the epoxidation reaction of eugenol and *isoeugenol* with perbenzoic acid.

By the results of titration the curves of dependence of $\ln[c/[b-(a-c)]]$ on time were plotted (a – is the initial molar concentration of PBA, mole/L; b – is the molar concentration of terpenes, mole/L; c – is the current molar concentration of PBA, mole/L).

The slope values for such curves were used for calculations of the second order rate constants k , L/(mole·min), for epoxidation reactions of eugenol and *isoeugenol* with perbenzoic acid in methylene chloride. The values of the rate constants were 0.44 L/(mole·min) for *isoeugenol* and 0.016 L/(mole·min)

Fig. 1. The structure of terpenoids – eugenol (1) and *isoeugenol* isomers (2–3)**Table 1**

The values of activation parameters of epoxidation of eugenol (4) and *isoeugenol* (5, 6) epoxidation with peracetic acid designed to approximate UBH&HLYP/6-31G (d)

Transition state	ΔH_{act} kJ/mole	ΔG_{act} kJ/mole
4	122.39	137.37
5	101.43	117.18
6	101.29	112.40

for eugenol, i. e. the rate constant of epoxidation of *isoeugenol* was in 5.5 times more than for its isomer (eugenol).

The method of the functional density (approximation UBH&HLYP/6-31G (d)) [22] was used for the quantum chemical calculation of activation parameters of the epoxidation reaction (Tab. 1–2) and modeling the transition states of molecules (Fig. 3–4).

As seen from Tab. 1, the values of activation barriers for epoxidation of *cis*- and *trans*-forms of *isoeugenol* were approximately 20 kJ/mole lower than that for eugenol coinciding with the experimental data.

The transition state of the eugenol epoxidation is in line with the synchronous process, in which the C–O-moeity of epoxide cycle has quite a close value. It is important to note that the structure has a closed electronic shell, i. e. in contrast to the reactions of epoxidation of other olefins the biradical transition state is not observed. According to the calculations, the activation parameters for epoxidation of *cis*- and *trans*-forms of *isoeugenol* have quite similar values, and it confirms the insignificant effect of stereoisomeric forms of molecules on their reactivity. The geometric parameters and spin density values on the corresponding atoms are also similar. As can be seen in Fig. 3, transition states **4** and **5** are close to planar character (see angles $C_{\alpha}C_{\beta}OH$), they are asymmetric

Table 2

The values of the activation parameters of epoxidation of eugenol (7) and *isoeugenol* (8, 9) with perbenzoic acid designed to approximate UBH&HLYP/6-31G (d)

Transition state	ΔH_{act} kJ/mol	ΔG_{act} kJ/mol
excluding the spin correction		
7	118.69	129.29
8	105.23	116.73
9	100.06	112.87
including the spin correction		
8	81.24	92.74
9	77.23	90.04




Fig. 2. Kinetic curves for the epoxidation reaction of eugenol and *isoeugenol* with perbenzoic acids ($c(\text{PBA}) = 0.1350$ mole/L; $c(\text{eugenol}) = 1.65\%$; $c(\text{isoeugenol}) = 1.65\%$; $T = 293$ K)




Fig. 3. The structure, geometric parameters (Å, deg.) and spin density values for some atoms of epoxidation transition states of eugenol (4) and isoeugenol (5, 6) with peracetic acid




Fig. 4. The structure, geometric parameters (Å, deg.) and spin density values for some atoms of epoxidation transition states of eugenol (7) and isoeugenol (8, 9) with perbenzoic acid

with a significant advantage in the bond formation of C_{α} -O above with C_{β} -O.

The analysis of the wave function transition states **5** and **6** show their biradicality. The highest spin density is localized on atoms C_2 and O_{27} . By coupling of the aromatic olefinic fragment there is a partial delocalization of the spin density at the *ortho*- and *para*-Carbon atoms.

The transition state **7** of eugenol epoxidation with perbenzoic acid is in line with the synchronous process, in which the C-O moiety of the epoxide cycle has quite similar value and a closed electronic shell. The analysis of the wave function transition states **8** and **9** shows their biradicality.

According to the calculation, the lowest activation barrier value is characterized by a reaction occurred through the transition state **9**. This trend persists even after consideration of the spin correction.

Experimental part

The following compounds were used in the work: eugenol (99 %, Sigma-Aldrich), *isoeugenol* (mixture of *cis*- and *trans*-isomers, 99 %, Sigma-Aldrich), benzoyl peroxide (≥ 98 %, Sigma-Aldrich), perbenzoic acid (98 % – determined by the iodometric titration), obtained by the method [23], acetic acid (at least 98 %), methylene chloride (p.a., LAB-SCAN).

To determine the *active oxygen content* for perbenzoic acid transfer 10.00 mL of its chloroform solution in a 100 mL Erlenmeyer flask, add 5 mL of 5 % KI solution and 5 mL of 30 % (w/w) acetic acid. Shake and allow to stand in a dark place for 30 minutes. Then titrate the liberated iodine with 0.1000 mole/L $Na_2S_2O_3$ solution using a 10 mL microburette (graduation is 0.02 mL). 1.00 mL of 0.1000 mole/L $Na_2S_2O_3$ solution corresponds to 0.0069 g of perbenzoic acid.

Diluted acetic acid: mix 31.3 wt. p. of acetic acid and 68.7 wt. p. of distilled water to obtain the acetic acid content of 29.5–30.5 % (w/w).

Potassium iodide solution, 5 %: dissolve 5.0 g of potassium iodide in 50 mL of freshly boiled and cooled water, and then dilute the solution obtained with

the same solvent to 100 mL. The solution should be colorless.

Sodium thiosulphate standard solution ($c(Na_2S_2O_3 \cdot 5H_2O) = 0.1000$ mole/L): prepare from 0.1 M Normadose® (standard titer) in freshly boiled and cooled distilled water with addition of 0.5 g of sodium carbonate in a 1000 mL measuring flask and dilute to the volume at 293 K.

Air thermostat TS-80m was used for maintaining the desired temperature of the reaction mixture.

The study of the epoxidation reaction kinetics

procedure: dissolve 0.8240 g of eugenol (or *isoeugenol*) in 49.2 mL of 0.1 mole/L solution of perbenzoic acid previously prepared in methylene chloride and mix to homogeneity. Over the determined period of time transfer 5.00 mL of the reaction mixture in a 150 mL Erlenmeyer flask using a pipette, add 2 mL of diluted acetic acid and 5 mL of 5 % potassium iodide solution while shaking. Titrate the liberated iodine with 0.1000 mole/L sodium thiosulphate solution using a microburette till the mixture becomes colorless. Repeat the titration without eugenol (control titration).

The method of the functional density (software Gaussian 09, approximation UBH&HLYP/6-31G(d)) was used for the quantum chemical calculation.

Conclusions

The study of epoxidation reaction mechanism of isomeric terpenoids eugenol and *isoeugenol* by density functional theory using the UBH&HLYP/6-31G (d) approach for the software Gaussian 09 has shown that the ratio of the activation energy for interaction of eugenol and *isoeugenol* with both acids (peracetic and perbenzoic) indicate higher reactivity of *isoeugenol*. These results are consistent with the kinetic data experimentally obtained; it confirms the correctness of the UBH&HLYP/6-31G (d) approach used to study the epoxidation characteristics of isomeric alkenes with peroxyacids.

Conflict of interests: authors have no conflict of interests to declare.

References

1. Koch, W. A chemist's guide to density functional theory / W. Koch, M. C. Holthausen. – 2nd ed. – Weinheim : Wiley-VCH Verlag GmbH, 2001. – 306 p. <https://doi.org/10.1002/3527600043>
2. Shi, H. Mechanism on epoxidation of alkenes by peracids: A protonation-promoted pathway and its quantum chemical elucidation / H. Shi, Z. Zhang, Y. Wang // J. Mol. Catal. A: Chem. – 2005. – Vol. 238, Issue 1–2. – P. 13–25. <https://doi.org/10.1016/j.molcata.2005.04.046>
3. A quantum-chemical study of the oxidation of ethylene by peroxyacetic acid derivatives / N. B. Filimonova, A. V. Vorob'ev, K. V. Bozhenko et al. // Russ. J. Phys. Chem. B. – 2010. – Vol. 4, Issue 2. – P. 235–241. <https://doi.org/10.1134/s1990793110020089>
4. Becke, A. D. Density-functional exchange-energy approximation with correct asymptotic behavior / A. D. Becke // Phys. Rev. A. – 1988. – Vol. 38, Issue 6. – P. 3098–3100. <https://doi.org/10.1103/physrev.38.3098>
5. Andersson, M. P. New scale factors for harmonic vibrational frequencies using the B3LYP density functional method with the triple- ζ basis set 6-311+G(d,p) / M. P. Andersson, P. Uvdal // J. Phys. Chem. A. – 2005. – Vol. 109, Issue 12. – P. 2937–2941. <https://doi.org/10.1021/jp045733a>
6. Kumara, A. Eugenol and its derivatives as antimicrobial agents / A. Kumara, D. Sahoob // J. Antimicrob. Photon. – 2013. – Vol. 128. – P. 133–140.
7. Antibacterial activity of terpenes and terpenoids present in essential oils / A. C. Guimarães, L. M. Meireles, M. F. Lemos et al. // Molecules. – 2019. – Vol. 24, Issue 13. – P. 2471. <https://doi.org/10.3390/molecules24132471>
8. Eugenol and derivatives activity against Mycobacterium tuberculosis, nontuberculous mycobacteria and other bacteria / A. L. De Almeida, K. R. Caleffi-Ferracioli, R. B. de L. Scodro et al. // Future Microbiol. – 2019. – Vol. 14. – P. 331–344. <https://doi.org/10.2217/fmb-2018-0333>

9. Hu, Q. Progress on the antimicrobial activity research of clove oil and eugenol in the food antisepsis field / Q. Hu, M. Zhou, S. Wei // J. Food Sci. – 2018. – Vol. 83, Issue 6. – P. 1476–1483. <https://doi.org/10.1111/1750-3841.14180>
10. Antimicrobial activity of eugenol and essential oils containing eugenol : A mechanistic viewpoint / A. Marchese, R. Barbieri, E. Coppo et al. // Crit. Rev. Microbiol. – 2017. – Vol. 43, Issue 6. – P. 668–689. <https://doi.org/10.1080/1040841x.2017.1295225>
11. In silico and in vitro investigation of the antifungal activity of isoeugenol against *Penicillium citrinum* / S. B. Ferreira, T. B. Dantas, D. de Figueiredo Silva et al. // Curr. Top. Med. Chem. – 2018. – Vol. 18, Issue 25. – P. 2186–2196. <https://doi.org/10.2174/1568026619666181130141818>
12. Anti-oxidant and anti-inflammatory effects of cinnamaldehyde and eugenol on mononuclear cells of rheumatoid arthritis patients / S. Mateen, M. T. Rehman, S. Shahzad et al. // Eur. J. Pharmacol. – 2019. – Vol. 852. – P. 14–24. <https://doi.org/10.1016/j.ejphar.2019.02.031>
13. Synthesis of eugenol derivatives and its anti-inflammatory activity against skin inflammation / A. K. Maurya, K. Agarwal, A. C. Gupta et al. // Nat. Prod. Res. – 2018. – P. 1–10. <https://doi.org/10.1080/14786419.2018.1528585>
14. An overview on the anti-inflammatory potential and antioxidant profile of eugenol / J. N. Barboza, C. da Silva Maia Bezerra Filho, R. O. Silva et al. // Oxid. Med. Cell. Longev. – 2018. – Vol. 2018. – P. 1–9. <https://doi.org/10.1155/2018/3957262>
15. Tsuchiya, H. Anesthetic agents of plant origin: a review of phytochemicals with anesthetic activity / H. Tsuchiya // Molecules. – 2017. – Vol. 22, Issue 8. – P. 1369. <https://doi.org/10.3390/molecules22081369>
16. Phytochemical analysis, cytotoxic and antioxidant activities of *Myrtus communis* essential oil from Morocco / Y. Harassi, M. Tilaoui, A. Idir, J. Frédéric, S. Baudino, S. Ajouai, H. A. Mouse, A. Ziad // J. Complement. Integr. Med. – 2019. <https://doi.org/10.1515/jcim-2018-0100>
17. Anti-metastatic and anti-proliferative activity of eugenol against triple negative and HER2 positive breast cancer cells / M. L. Abdullah, M. M. Hafez, A. Al-Hoshani, O. Al-Shabanah // BMC Complement Altern. Med. – 2018. – Vol. 18, Issue 1. – P. 321. <https://doi.org/10.1186/s12906-018-2392-5>
18. Federation of American Societies for Experimental Biology. (1973). Evaluation of health aspects of oil of cloves as a food ingredient. Life Sciences Research Office. Report no SCOGS-19; U.S. Food and Drug Administration. [Електронний ресурс]. – Режим доступу : http://www.faseb.org/Portals/2/PDFs/LSRO_Legacy_Reports/1973_SCOGS-19%20Oil%20of%20Cloves.pdf
19. Kegley, S. Marin Municipal Water District. Herbicide Risk Assessment. Chapter 6. Clove Oil (Eugenol) / S. Kegley, E. Conlisk, M. Moses. – Berkeley: Pesticide Research Institute, 2010. – 25 p. [Електронний ресурс]. – Режим доступу : <https://www.marinwater.org/DocumentCenter/View/259/Herbicide-Risk-Assessment-Chapter-1-Summary-January-10-2010>
20. Elgendy, E. M. Oxidation reactions of some natural volatile aromatic compounds : anethole and eugenol / E. M. Elgendy, S. A. Khayyat // Russ. J. Organ. Chem. – 2008. – Vol. 44, Issue 6. – P. 823–829. <https://doi.org/10.1134/s1070428008060079>
21. Naboka, V. M. Quantum-chemical investigation of the mechanism of eugenol and isoeugenol epoxidation by peracetic acid / V. M. Naboka, O. M. Agafonov // Topical issues of new drugs development: Abstracts of International Scientific and Practical Conference of Young Scientists and Students (April 23, 2015, Kharkiv). – Kharkiv : NUPh, 2015. – P. 90–91.
22. Gaussian 09, Revision A.02 / M. J. Frisch, G. W. Trucks, H. B. Schlegel et al. – Gaussian, Inc., Wallingford CT, 2009.
23. Губен-Вейль. Методы органической химии: в 2 т. / Губен-Вейль. – М. : Химия, 1967. – Т. 2. – 1032 с.

References

1. Koch, W., Holthausen, M. C. (2001). *A chemist's guide to density functional theory*, 2nd ed. Weinheim: Wiley-VCH Verlag GmbH, 306. <https://doi.org/10.1002/3527600043>
2. Shi, H., Zhang, Z., Wang, Y. (2005). Mechanism on epoxidation of alkenes by peracids: A protonation-promoted pathway and its quantum chemical elucidation. *Journal of Molecular Catalysis A: Chemical*, 238(1-2), 13–25. <https://doi.org/10.1016/j.molcata.2005.04.046>
3. Filimonova, N. B., Vorob'ev, A. V., Bozhenko, K. V., Moiseeva, N. I., Dolin, S. P., Gekhman, A. E., Moiseev, I. I. (2010). A quantum-chemical study of the oxidation of ethylene by peroxyacetic acid derivatives. *Russian Journal of Physical Chemistry B*, 4(2), 235–241. <https://doi.org/10.1134/s1990793110020089>
4. Becke, A. D. (1988). Density-functional exchange-energy approximation with correct asymptotic behavior. *Physical Review A*, 38(6), 3098–3100. <https://doi.org/10.1103/physreva.38.3098>
5. Andersson, M. P., Uvdal, P. (2005). New Scale Factors for Harmonic Vibrational Frequencies Using the B3LYP Density Functional Method with the Triple- ζ Basis Set 6-311+G(d,p). *The Journal of Physical Chemistry A*, 109(12), 2937–2941. <https://doi.org/10.1021/jp045733a>
6. Kumara, A., Sahoob, D. (2013). Eugenol and its derivatives as Antimicrobial Agents. *Journal of Antimicrobials Photon*, 128, 133–140.
7. Guimarães, A. C., Meireles, L. M., Lemos, M. F., Guimarães, M. C. C., Endringer, D. C., Fronza, M., Scherer, R. (2019). Antibacterial Activity of Terpenes and Terpenoids Present in Essential Oils. *Molecules*, 24(13), 2471. <https://doi.org/10.3390/molecules24132471>
8. De Almeida, A. L., Caleffi-Ferracioli, K. R., de L. Scodro, R. B., Baldin, V. P., Montaholi, D. C., Spricigo, L. F., Nakamura-Vasconcelos, S. S., Hegeto, L. A., Sampiron, E. G., Costacurta, G. F., Dos, S., Yamazaki, D. A., Gauze, F. G., Siqueira, V. L., Cardoso, R. F. (2019). Eugenol and derivatives activity against *Mycobacterium tuberculosis*, nontuberculous mycobacteria and other bacteria. *Future Microbiology*, 14, 331–344. <https://doi.org/10.2217/fmb-2018-0333>
9. Hu, Q., Zhou, M., Wei, S. (2018). Progress on the Antimicrobial Activity Research of Clove Oil and Eugenol in the Food Antisepsis Field. *Journal of Food Science*, 83(6), 1476–1483. <https://doi.org/10.1111/1750-3841.14180>
10. Marchese, A., Barbieri, R., Coppo, E., Orhan, I. E., Daglia, M., Nabavi, S. F., Izadi, M., Abdollahi, M., Nabavi, S. M., Ajami, M. (2017). Antimicrobial activity of eugenol and essential oils containing eugenol: A mechanistic viewpoint. *Critical Reviews in Microbiology*, 43(6), 668–689. <https://doi.org/10.1080/1040841x.2017.1295225>
11. Ferreira, S. B., Dantas, T. B., de Figueiredo Silva, D., Ferreira, P. B., de Melo, T. R., de Oliveira Lima E. (2018). In Silico and In Vitro Investigation of the Antifungal Activity of Isoeugenol against *Penicillium citrinum*. *Current Topics in Medicinal Chemistry*, 18(25), 2186–2196. <https://doi.org/10.2174/1568026619666181130141818>
12. Mateen, S., Rehman, M. T., Shahzad, S., Naeem, S. S., Faizy, A. F., Khan, A. Q., Khan, M. S., Husain, F. M., Moin, S. (2019). Anti-oxidant and anti-inflammatory effects of Cinnamaldehyde and Eugenol on mononuclear cells of rheumatoid arthritis patients. *European Journal of Pharmacology*, 852, 14–24. <https://doi.org/10.1016/j.ejphar.2019.02.031>
13. Maurya, A. K., Agarwal, K., Gupta, A. C., Saxena, A., Nooreen, Z., Tandon, S., Ahmad, A., Bawankule, D. U. (2018). Synthesis of eugenol derivatives and its anti-inflammatory activity against skin inflammation. *Natural Product Research*, 1–10. <https://doi.org/10.1080/14786419.2018.1528585>
14. Barboza, J. N., da Silva Maia Bezerra Filho, C., Silva, R. O., Medeiros, J. V. R., de Sousa, D. P. (2018). An Overview on the Anti-inflammatory Potential and Antioxidant Profile of Eugenol. *Oxidative Medicine and Cellular Longevity*, 2018, 1–9. <https://doi.org/10.1155/2018/3957262>
15. Tsuchiya, H. (2017). Anesthetic Agents of Plant Origin: A Review of Phytochemicals with Anesthetic Activity. *Molecules*, 22(8), 1369. <https://doi.org/10.3390/molecules22081369>

16. Harassi, Y., Tilaoui, M., Idir, A., Frédéric, J., Baudino, S., Ajouai, S., Mouse, H. A., Ziad, A. (2019). Phytochemical analysis, cytotoxic and antioxidant activities of *Myrtus communis* essential oil from Morocco. *Journal of Complementary and Integrative Medicine*. <https://doi.org/10.1515/jcim-2018-0100>
17. Abdullah, M. L., Hafez, M. M., Al-Hoshani, A., Al-Shabanah, O. (2018). Anti-metastatic and anti-proliferative activity of eugenol against triple negative and HER2 positive breast cancer cells. *BMC Complementary and Alternative Medicine*, 18(1), 321. <https://doi.org/10.1186/s12906-018-2392-5>
18. Federation of American Societies for Experimental Biology. (1973). Evaluation of health aspects of oil of cloves as a food ingredient. Life Sciences Research Office. Report no SCOGS-19; U.S. Food and Drug Administration. Available at: http://www.faseb.org/Portals/2/PDFs/LSRO_Legacy_Reports/1973_SCOGS-19%20oil%20of%20Cloves.pdf
19. Kegley, S., Conlisk, E., Moses, M. (2010). *Marin Municipal Water District. Herbicide Risk Assessment. Chapter 6. Clove Oil (Eugenol)*. Berkeley: Pesticide Research Institute, 25. Available at: <https://www.marinwater.org/DocumentCenter/View/259/Herbicide-Risk-Assessment-Chapter-1-Summary-January-10-2010>
20. Elgendy, E. M., Khayyat, S. A. (2008). Oxidation reactions of some natural volatile aromatic compounds: anethole and eugenol. *Russian Journal of Organic Chemistry*, 44(6), 823–829. <https://doi.org/10.1134/s1070428008060079>
21. Naboka, V. M., Agafonov, O. M. (2015). Quantum-chemical investigation of the mechanism of eugenol and isoeugenol epoxidation by peracetic acid. *Topical issues of new drugs development*, Kharkiv, 2015, 90–91.
22. Frisch, M. J., Trucks, G. W., Schlegel, H. B., Scuseria, G. E., Robb, M. A., Chessexman, J. R. (2009). *Gaussian 09, Revision A.02*. Gaussian, Inc., Wallingford CT.
23. Guben-Vejl'. (1967). *Metody organicheskoy himii: Vol. 1-2*. Moscow: Himija, 2, 1032.

Надійшла до редакції 07.06.2019 р.