

In vivo administration of fucoidan from *Turbinaria decurrens* protects shrimps from white spot syndrome virus

*S. Meenakshi¹, R. Saravanan², T. Balasubramanian³, & A. Palavesam¹

¹Department of Animal Science, Manonmaniam Sundaranar University, Tirunelveli, India

²Sri K.G.S Arts College, Srivaikuntam, India

³Faculty of Marine Sciences, Annamalai University, India

[E-mail: smeenakshi85@gmail.com]

Received 06 March 2017; revised 02 June 2017

Fucoidan was extracted from the brown seaweed *Turbinaria decurrens* by hot water extraction and characterized with HPLC, FTIR, NMR and GPC to study the impact against the White Spot Syndrome Virus (WSSV), bacteria and fungi. Fucoidan was fed for shrimps along with the diet (before and after WSSV infection). In the end of the study, the survival percentage of shrimp was 51% (in the case of 5-8 g) and 97% (in the case of 12-15 g), respectively. Among the 10 bacterial pathogens and 7 fungal pathogens, fucoidan inhibited 2 bacterial pathogen such as *S. aureus* (0.26±0.02 mm) and *E. faecalis* (1.3±0.1 mm) and *A. Niger* (0.84 ± 0.03 mm) in fungal pathogens with clear zone of inhibition.

[Keywords: Fucoidan, Brown seaweeds, WSSV, Antimicrobial activity, Prawn feed]

Introduction

WSSV is a viral disease affecting most of the commercially culturing marine shrimp species in the world. WSSV infection is characterized by a rapid disease on set and high mortalities. The principal clinical sign of WSSV is the presence of white spot on the carapace and the body colour of diseased shrimps becomes pale or reddish in colour¹. Out breaks of WSSV causing serious mortality to cultured shrimp have been occurring in Asia since 1992. In order to prevent WSSV infection in cultured shrimp, the present study is planned to use the purified fucoidan from the brown seaweed *T. decurrens*.

Fucoidan represent 40% of the dry weight of isolated cell walls of brown seaweeds^{2,3}. It is composed of fucose, uronic acids, galactose, xylose and sulfated fucose^{4,5}. They are comprised of long chains of linked sugar molecules, along with sulfate groups which make them negatively charged. They composed primarily of highly branched (1→2) or (1→3) linked α -L-fucose-4-sulfate units⁶. Structures and chemical composition of polysaccharide vary among the species and sometimes among different parts of the plant^{6,7}. This structural complexity, chemical composition of fucoidan of algal origin may vary depending on the method of extraction⁸. Fucoidan interfere in several cellular processes such as migration, adhesion, proliferation and apoptosis^{2,9}.

Thus, each new fucoidan of algal origin has unique compound with unique structural features which may display varied bioactivity and would be a potentially a new drug.

Fucoidan from algae possess important pharmacological activities such as antioxidant¹⁰, anticoagulant and antithrombotic^{11,12}, antiproliferative, anti-inflammatory, antiviral¹³, antitumour and immunomodulatory¹⁴, hepatoprotective¹⁵, neuroprotective¹⁶. *Turbinaria* species have been reported to have wide range of pharmacological activities. Only limited number of studies has been done in *Turbinaria decurrens*. So, the present study was planned to isolate fucoidan from *T. decurrens* and to study against WSSV infection in *P. monodon*.

Materials and Methods

The brown seaweed *T. decurrens* belonging to the family phaeophyceae was collected in yervadi, India. Specimens were identified and authenticated by phycologist Dr.P.Anantharaman, Faculty of Marine Sciences, Annamalai University, Parangipettai, India. The collected *T. decurrens* was initially washed with seawater to remove the macroscopic epiphytes and other extraneous matter and then rinsed in distilled water. Specimen was shade dried and coarsely powdered. 100g of dried seaweed powder was depigmented with acetone followed by hot water

extraction at 90-95°C for 3-4 h. Brown colored syrup was then filtered through whatmann No.3 filter paper, concentrated to ¼ of the original volume, cooled and precipitate was centrifuged at 5000 rpm, dehydrated with diethyl ether and further purified in HPLC (data not shown) to get a sulphated polysaccharide fucoidan¹⁷. FT-IR spectroscopy of sulphated polysaccharide revealed the presence of hydrogen bonded compound, possible acid or amide salt, ionized compound and aliphatic tertiary amine salt. 1H- NMR analysis shows the presences of signals corresponding to fucose. 3 OH groups appeared at 3.95 ppm, C=C-H (1H) (acetylenic group) signals at 3.45 ppm, C≡C-H (1H) (methylenic proton) signals at 2.35 ppm and 2.30 ppm C-H2 protons appeared at 1.34 ppm (2H). Presence of these signals clearly indicates that the isolated polysaccharide from *T. decurrens* is fucoidan. Biochemical estimation of the isolated fucoidan is carbohydrate 59.62%, sulfate 26.52%, and Uronic acid 6.3%. Molecular weight of the compound was determined in gel permeation chromatography as 2,34,000 Da¹⁶. All the chemicals were purchased from sigma chemical company (St. Louis, Mo) and Himedia (Mumbai, India).

Healthy shrimps weighing 5 – 8 and 12 – 15 g were used in this study. Both groups were performed separately in duplicate with 15 shrimps each treatment. Fucoidan was fed by mixing it with shrimp diet in 100, 200 and 400 mg/ kg (5-8 g) and 100 and 200 mg/ kg (12-15) of shrimp respectively. Fucoidan was fed for 4 days before WSSV infection. The shrimp were challenged by incubation in the virus solution for 2.5 h. This virus concentration caused the control shrimp to die in 3–5 days. The fucoidan was fed continuously for 15 days. The same diet without fucoidan was fed to the control groups, which were also challenged with WSSV. The shrimp were monitored for 10 days after infection and all mortality were recorded.

The WSSV virus solution was prepared by using gills, heart and lymphoid organ of WSSV-infected shrimp, crushed in K-199 (1% w/v M199, 1.88M NaCl, 0.06 M CaCl₂·2H₂O, 0.1M L-glutamine, 9.14 mM Herpes and 10% (v/v) salt mixture which consists of 0.05 M KCl, 0.12 M MgSO₄·7H₂O, 0.16M MgCl₂·6H₂O and 3.2 mM NaH₂PO₄·2H₂O, pH 7.3-7.6) at 1:2 (W/V). Suspension was precipitated by centrifugation at 3000×g for 10min; the supernatant fluid was then centrifuged at 8000×g, 4°C for 39 min, filtered through a 0.45µm membrane and stored in 15% glycerol at -70°C until used¹⁸.

Test group of the shrimp weight was 12 – 15 g fed with fucoidan 200 mg/ kg of body weight and WSSV infected (group 3). Phagocytic activity was compared with that of the normal diet control (group 1) and WSSV infected shrimp (group 2). Haemolymph (0.5 ml) was collected on day 7 of the culture and mixed with 0.5 ml of KC-199 (Itami *et al.*, 1994). Haemocytes were separated by centrifugation at 6500 rpm at 4 °C, then washed with KC-199. Haemocytes were then suspended ~10⁷ cells in 0.2ml and mixed with 0.2 ml of latex beads (~10⁸/ml, particle diameter 1.094µm) on a clean glass slide. Mixture was incubated in a moisture chamber at room temperature for 30 min. The cells were fixed by 2.5% glutaraldehyde for 5 min, after which the nonadherent cells were removed with 0.85% NaCl, then followed by air-drying, and staining with Wright stain. Numbers of ingested cells and ingesting cells were counted from 200 cells and calculated for percentage of phagocytosis, phagocytic index (PI) and average number of the beads ingested per cell (ABPC) as follows¹⁹

$$\left(\text{Percentage of phagocytosis} \right) = \left(\frac{\text{number of cells ingesting bead}}{\text{number of cell observed}} \right) \times 100$$

$$\left[\text{Phagocytic index (PI)} \right] = \left(\frac{\text{number of cells ingesting bead}}{\text{number of cell observed}} \right) \times \left(\frac{\text{number of bead ingested}}{\text{number of cells observed}} \right) \times 100$$

$$\left[\text{Average number of the bead ingested per cell (ABPC)} \right] = \left(\frac{\text{number of bead ingested}}{\text{number of cell ingesting bead}} \right)$$

The antimicrobial activity of the fucoidan from *T. decurrens* was assayed using the paper disc diffusion method. Human pathogens were obtained from the Rajamuthaihal medical college, Annamalai University. Bacterial pathogens namely *Escherichia coli*, *Aeromonas hydrophila*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Enterococcus faecalis*, *Enterobacter aerogenes*, *Micrococcus luteus*, *Salmonella typhi*, *Bacillus subtilis* and *B. megaterium* were used to study the antibacterial activity using muller hinton agar medium. Fungal pathogens namely

Aspergillus niger, *Aspergillus flavus*, *Penicillium* sp, *Trichoderma* sp, *Trichothecium* sp, *Hormodendrum* sp and *Rhizopus* sp were used to study the antifungal activity using potato dextrose agar medium. 1, 1.5 and 2 mg/ml of fucoidan from *T. decurrens* were dissolved in sterile distilled water. 50µl from each concentration was poured into sterile filter paper discs (5 mm diameter) and dried for 30 min, prior to the inoculation. The disc was placed onto the surface of bacterial and fungal pathogens swabbed petriplates and incubated for 24-72 hours. After incubation, the clearance zones around the discs were measured in millimeters and the assays were carried out in triplicate. Erythromycin was used as control for antibacterial and nystatin for antifungal activity.

Results

Polysaccharide was extracted using hot water and yield of crude polysaccharide was 10.43 g/kg. The crude polysaccharide was precipitated with calcium chloride to separate fucoidan from alginates. The precipitated polysaccharide was partially purified in DEAE cellulose52 using 2M sodium chloride and the yield was 6.46 g/kg. The active fraction was identified by carbohydrate estimation using phenol sulfuric acid method and it was used for the further study. Partially purified sample was further purified in HPLC using C18 column and the yield was 3.38 g/kg.

100, 200, 400 mg/kg of body weight of fucoidan was fed to 5-8 g and 12-15 g of shrimps before and after immersing in WSSV solution. Survival rate of 5-8 g of shrimp was 5.8%, 19.3% and 61% respectively. In case of 12-15 g of shrimp was 54% and 97% survived respectively. For shrimps, which did not get fucoidan, 50% were dead 3 – 5 days after WSSV infection and 100% after 15 days. Phagocytic activity (Table 1) for the fucoidan fed shrimps with WSSV infection was found to have a higher phagocytic index, a higher percentage of phagocytosis and a higher ABPC than both of the control groups.

Antimicrobial activity was determined in fucoidan from *T. decurrens* at 1, 1.5 and 2 mg/ml (Table 2). Antibacterial activity was observed at 2 mg/ml against

S. aureus and *E. faecalis* with a clear zone of 0.26±0.02 mm and 1.3±0.1 mm respectively. In antifungal activity clear zone of inhibition was observed only against *A. niger* at 2mg/ml with 0.84 ± 0.03 mm.

Discussion

Viral and bacterial infections together with poor water quality are the main reason for shrimp mortality. Reinforcing resistance against the invading pathogens of shrimp by improving shrimp immunity is an acceptable strategy for defending diseases. Unlike the mammals and fishes, crustaceans over lacks a specific immune system²⁰. Usage of natural immunostimulant as a better remedy than administrating vaccines and antibiotics is accepted for the control of pathogen in aquaculture.

Among the different extraction method for polysaccharide, hot water extraction contributed higher yield^{17,21}. The present study supports the above statement by providing a good yield (10.43 g/kg) of polysaccharide using hot water extraction. The crude polysaccharide was further purified in HPLC to isolate fucoidan and the yield was 3.38 g/kg. Due to the high antioxidant potential, fucoidan from *T. decurrens* showed promising activity against the liver damage¹⁵ and parkinsons disease¹⁶.

Fucoidan from *Cladosiphon okamuranus* was reported to control WSSV in *P. japonicus*²² and crude fucoidan of *Sargassum polycystum* was reported to

Table 2 — Antimicrobial activity of fucoidan from *T. decurrens*

S. No.	Pathogens	1 mg/ml	1.5 mg/ml	2 mg/ml
Bacterial pathogens				
1.	<i>Escherichia coli</i>	-	-	-
2.	<i>Aeromonas hydrophila</i>	-	-	-
3.	<i>Staphylococcus aureus</i>	-	-	+
4.	<i>Pseudomonas aeruginosa</i>	-	-	-
5.	<i>Enterococcus faecalis</i>	-	-	+
6.	<i>Enterobacter aerogenes</i>	-	-	-
7.	<i>Micrococcus luteus</i>	-	-	-
8.	<i>Salmonella typhi</i>	-	-	-
9.	<i>Bacillus subtilis</i>	-	-	-
10.	<i>Bacillus megaterium</i>	-	-	-
Fungal pathogens				
1.	<i>Aspergillus niger</i>	-	-	+
2.	<i>Aspergillus flavus</i>	-	-	-
3.	<i>Penicillium</i> sp	-	-	-
4.	<i>Trichoderma</i> sp	-	-	-
5.	<i>Trichothecium</i> sp	-	-	-
6.	<i>Hormodendrum</i> sp	-	-	-
7.	<i>Rhizopus</i> sp	-	-	-

Table 1 — Phagocytic activity of shrimp

Immunity indexes	Group1 (control shrimps)	Group 2 (WSSV infected)	Group 3 (fed with fucoidan and WSSV infected)
Phagocytic index (n=10)	1.2 ±0.4	0.41±0.31	3.41±1.23
%phagocytosis (n=10)	6.27±2.43	3.63±1.32	10.4±1.98
ABPC (n=10)	2.0±0.62	1.5±1.2	3.1±0.75

control WSSV in *P. monodon*¹⁸. Fucoïdan from *T. decurrens* fed to the WSSV infected shrimps reduced the mortality rate in both groups (5– 8 and 12 – 15 g) of the experiment. Mechanism of inhibition of WSSV is unclear. Fucoïdan from *C. okamuranus* was also reported to control WSSV in *P. japonicus*^{18,22} but the mechanism of the fucoïdan extracted from *C. okamuranus* were not disclosed. The 26.52% sulfate of the Fucoïdan from *T. decurrens* inhibited WSSV and the sulfate content effects on antiviral (WSSV) activity as previously proposed by Witvrouw & De Clercq²³ therefore, the lower dose may be manipulated by hypersulfation of the fucoïdan, or the fucoïdan from other sources with a higher content of sulfate are needed.

The cell extracts and active constituents of various algae have been shown to have antibacterial activity *in vitro* against Gram-positive and Gram-negative bacteria²⁴. Fucoïdan from *T. decurrens* showed antibacterial activity against *S. aureus* and *E. faecalis* and antifungal activity *A. niger* with minimum zone of inhibition. The polysaccharide extracted from algal species showed a high content of sulfate, but the sulfate content did not show a clear relation to the antimicrobial activity. In contrast, Berteau and Mulloy² reported that the antimicrobial activity of the polysaccharides was related to their chemical structure and ester sulfate groups. On the other hand, several species of brown seaweed that have a high sulfate content have been reported to have antimicrobial activities^{25,26}. Fucoïdan from macroalgae possess antibacterial activity^{18,27}. Crude fucoïdan from brown algae *Sargassum polycystum* at 12 mg/ml inhibited the growth of *V. harveyi*, *S. aureus* and *E. coli* which was the supportive evidence for the present study and the concentrations were higher than the present study.

Conclusion

In future Fucoïdan will be as one of the immunostimulant which increases phagocytic activity of the shrimps. From this study it was concluded that Fucoïdan from *T. decurrens* can be used as the immunostimulant and as in feed formulation to control WSSV in culture ponds.

Acknowledgement

The first author thanks The University Grant Commission for the financial support through Dr. D.S. Kothari Postdoctoral Fellowship during

manuscript production and the authorities of Annamalai University for providing the necessary facilities to carry out the work.

References

- 1 Takahashi, Y., Itami, T., Kondo, M., Maeda, M., Fujji, R., Tomonaga, S., Supamattaya, K. & Boonyaratpalin, S., Electron microscopic observation of bacilliform virus infection in kuruma shrimp (*Penaeus japonicus*). *Fish Pathol.* 19 (1994) 121-125.
- 2 Berteau, O. & Mulloy, B., Sulfated fucans, fresh perspectives: Structures, functions, and biological properties of sulfated fucans and an overview of enzymes active toward this class of polysaccharide. *Glycobiology*, 13(2003) 29-40.
- 3 Kloareg, B., Isolation and analysis of cell walls of the brown marine algae *Pelvetia canaliculata* and *Ascophyllum nodosum*. *Physiol. Veg.* 22 (1984) 47-56.
- 4 Percival, E., Richard, H. & Dowell, M. C., Chemistry and Enzymology of Marine Algal Polysaccharides. Academic Press, London, 219 (1967) 68 pp.
- 5 Leite, E L., Medeiros, M G L., Rocha, H A O., Farias, G G M., Silva, L F., Chavante, S F., Abreu, LRD., Dietrich, C P. & Nader, H B., Structure and pharmacological activities of sulphated xylofucoglucuronan from the alga *Spatoglossum schröderi*. *Plant Sci*, 132 (1998) 215-228.
- 6 Ermakova, S., Sokolova, R., Kim, S M., Um, B H., Isakov, V. & Zvyagintseva, T., Fucoïdins from brown seaweeds *Sargassum hornery*, *Eclonia cava*, *Costaria costata*: structural characteristics and anticancer activity. *Appl Biochem Biotechnol*, 164 (6) (2011) 841-850.
- 7 Chevolut, L., Foucault, A., Kervarec, N., Siquin, C., Fischer, A M. & Boisson-Vidal, C., Further data on the structure of brown seaweed fucans: relationships with anticoagulant activity. *Carbohydr Res*, 319 (1999): 154-165.
- 8 Grauffel, V., Kloareg, B., Mabeu, S., Durand, P. & Jozefonvicz, J., New natural polysaccharides with potent antithrombotic activity: Fucans from brown algae. *Biomaterials*, 10 (1989) 363-369.
- 9 Rocha, AB., Lopes, RM. & Schwartsmann, G., Natural products in anticancer therapy. *Curr Opin Pharmacol*, 1 (2001) 364-9.
- 10 Wang, J., Zhang, Q., Zhang, Z., Li, & Z., Antioxidant activity of sulphated polysaccharide fractions extracted from *Laminaria japonica*. *Int. J. Biol. Macromolec.* 42 (2008) 127-132.
- 11 Zoysa, M., Nikapitiya, C., Jeon, Y J., Jee Y & Lee, J., Anticoagulant activity of sulfated polysaccharide isolated from fermented brown seaweed *Sargassum fulvellum*. *J. Appl. Phycol.* 20 (2008) 67-74.
- 12 Rocha de Souza, M C., Marques, C T., Dore, C M G., Ferreira da Silva, F R., Rocha, H A O. & Leite, E L., Antioxidant activities of sulphated polysaccharides from brown and red seaweeds. *J. Appl. Phycol.* 19 (2007) 153-160.
- 13 Hayashi, K., Nakano, T., Hashimoto, M., Kanekiyo, K. & Hayashi, T., Defensive effects of a fucoïdan from brown alga *Undaria pinnatifida* against herpes simplex virus infection. *Int. Immunopharmacol.* 8 (2008) 109-116.
- 14 Kima, M H. & Joo, H G., Immunostimulatory effects of fucoïdan on bone marrow-derived dendritic cells. *Immunol. Lett*, 115 (2008) 138-143.

- 15 Meenakshi, S., Umayaparvathi, S., Saravanan, R., Manivasagam, T. & Balasubramanian T., Hepatoprotective effect of fucoidan isolated from the seaweed *Turbinaria decurrens* in ethanol intoxicated rats., *Int. J. Biol. Macromolec*, 67 (2014) 367–372.
- 16 Meenakshi, S., Umayaparvathi, S., Saravanan, R., Manivasagam, T. & Balasubramanian, T., Neuroprotective effect of fucoidan from *Turbinaria decurrens* in MPTP intoxicated Parkinsonic mice *Int. J. Biol. Macromolec*, 86 (2016) 425-433.
- 17 Ananthi, S., Ragavendran, H R B., Gopalan Sunil, A., Gayathri, V., Ramakrishnan, G. & Vasanthi, H R., *In vitro* antioxidant and *in vivo* anti-inflammatory potential of crude polysaccharide from *Turbinaria ornata* (Marine brown alga). *Food Chemi Toxicol*, 48 (2010) 187-192.
- 18 Chotigeat, W., Tongsupa, S., Supamataya, K. & Phongdara, A., Effect of Fucoidan on Disease Resistance of Black Tiger Shrimp. *Aquaculture*, 233 (2004) 23-30.
- 19 Itami, T., Takahashi, Y., Tsuchihira, E. & IgusaKondo, H M., Enhancement of disease resistance of kuruma prawn *Penaeus japonicus* and increase in phagocytic activity of prawn hemocytes after oral administration of h 1,3 – glucan (Schizophyllan). In: Chou, L.M., Munro, A.D., Lam, T.J., Chen, T.W., Cheong, L.K.K., Ding, J.K., Hooi, K.K., Khoo, H.W., Phang, V.P.E., Shim, K.F., Tan, C.H. (Eds.): *The 3rd Asian Fisheries Forum. Asian Fisheries Society* Manila, Philippines, (1994):375-378.
- 20 Raa J., The use of immune-stimulants in fish and shellfish feeds. In: *Avances en Nutricion Acuicola V. Memorias del V Simposium Internacional de Nutricion Acuicola*. Cruz-Suarez L. E., Ricque-Marie D., Tapia-Salazar M., Olvera-Novoa M. A., CiveraCerecedo R. (eds), 19-22 November, 2000, Merida, Yucatan Mexico, pp.47-56.
- 21 Rioux, L E., Turgeon, S L. & Beaulieu, M., Characterization of polysaccharides extracted from brown seaweeds, *Carbohydrate Polymers*, 69 (2007) 530-537.
- 22 Takahashi, Y., Uehara, K., Watanabe, R., Okumura, T., Yamashita, T., Omura, H., Yomo, T., Kanemitsu, A., Kawano, T., Narasaka, H., Suzuki, N. & Itami, T., Efficacy of oral administration shrimp in Japan. In: Flegel, T.W. (Ed.): *Advances in Shrimp Biotechnology: National Center for Genetic Engineering and Biotechnology*. Bangkok, pp. (1998) 171-173.
- 23 Witvrouw, M. & De Clercq, E., Sulfated polysaccharides extracted from sea algae as potential antiviral drugs. *Gen. Pharmacol*, 29 (1997) 497-511.
- 24 Ely, R., Supriya, T. & Naik, C G., Antimicrobial activity of marine organisms collected off the coast of South East India. *J. Experimental Biol. Ecol*, 309 (2004) 121-127.
- 25 Adhikari, U., Mateu, C G., Chattopadhyay, K., Pujol, C A., Damonte, E B. & Ray, B., Structure and antiviral activity of sulfated fucans from *Stoechospermum marginatum*. *Phytochemistry*, 67(2006): 2474-2482.
- 26 Asker, M M S., Mohamed, S F., Ali, F M. & El-Sayed, O H., Chemical structure and antiviral activity of water-soluble sulfated polysaccharides from *Sargassum latifolium*. *J. Appl. Sci. Res*, 3 (2007) 1178-1185.
- 27 Kantachumpoo, A. & Chirapart, A., Components and Antimicrobial Activity of Polysaccharides Extracted from Thai Brown Seaweeds. *Nat. Sci*, 44(2010) 220-233.