

Stock structure analysis of *Nemipterus bipunctatus* (Valenciennes, 1830) from three locations along the Indian coast

Vaisakh G.^{1*}, S.K. Chakraborty², A. K. Jaiswar², Sibina Mol S.³, Renjith R. K.⁴, & Sreekanth G. B.⁵

¹ICAR-Central Inland Fisheries Research Institute RRC, Vadodara, Gujarat, 390022, India

²ICAR-Central Institute of Fisheries Education, Mumbai, Maharashtra, 400 061, India

³ICAR-Central Inland Fisheries Research Institute RRC, Bengaluru, Karnataka, 390022, India

⁴ICAR-Central Institute of Fisheries Technology, Willington Island, Kochi, 682 029, India

⁵ICAR-Central Coastal Agricultural Research Institute, Old Goa, Goa, 403 402, India

*[E-mail: vaisakh.login@gmail.com]

Received 04 May 2018; revised 18 June 2018

Present study was done to identify the occurrence of various stocks of *Nemipterus bipunctatus* along the Indian coast, based on their body and skull shape morphometrics. Fish samples were collected from three locations along the Indian coast viz. Chennai along the East coast and Mumbai and Veraval on the West coast. Twenty truss distances from nine-point truss network of body and twenty-one truss distances from eleven-point truss network of the skull were measured from each fish sample. The canonical discriminant analysis showed that the truss distances belong to the anterior region and caudal peduncle of body and olfactory region of skull were significant in separating the fish stocks. The artificial neural network analysis revealed 91.4 % and 86.14 % well classification of the specimen, based on the truss distances of body and skull respectively. The results from the study indicated that there is a significant difference among the stocks of *N. bipunctatus*.

[Keywords: Body morphometry; Delagoa threadfin bream; *Nemipterus bipunctatus*; Skull morphometry; Stock structure; Truss network analysis]

Introduction

A fish stock can be defined as a subset of species which encompass an intraspecific group of randomly mating individuals with spatio-temporal integrity¹. As an intraspecific group of individuals, fisheries management measures without considering the stock structure of a fish species may consequently lead to its overexploitation². Hence, it is important to perceive more information on stock structure of a target fish species for the development of well outlined policies and sustainable management measures in order to optimize the yield of its multiple stocks which are differentially exploited³.

Stock identification studies based on meristics, morphometrics, otolith microchemistry, molecular genetics etc., were practiced in fishes⁴⁻⁷. Morphometric analysis offers more efficacious, facile and powerful techniques which essentially discriminates 'phenotypic stocks' that are groups of individuals with homogenous growth, mortality and reproductive rates⁸. A better description of shape is possible with the analysis of morphometric characters using truss network system⁹. Also truss network analysis helps to disclose more

complex aspects of shape differences in intraspecific population, in comparison to traditional morphometric methods¹⁰.

Threadfin breams under the family *Nemipteridae* constitute nearly 18 % of the total demersal finfish landings and 4.5 % of total marine landings in India¹¹. *N. bipunctatus* is one of the important threadfin bream species available in Indian waters¹¹. Considerable landing of this species was started reporting from some maritime states of India such as Maharashtra, Tamil Nadu and Gujarat in recent years¹¹. It is a benthic carnivorous species, found on sand or mud bottoms in depths between 18 to 100 m. Studies were conducted on the biology and fisheries of *N. bipunctatus* in India^{12,13}. But in spite of its commercial importance, extensive study has not been undertaken to identify stock structure of this species along the Indian coast. The present paper seeks to study the stock structure of *N. bipunctatus* from three sites along Indian coast based on truss network analysis so that the management measures in future can be designed based on assessments from independent fish stocks.

Materials and Methods

Sampling

Samples of *Nemipterus bipunctatus* along Indian coast were collected from three different locations two along the North west coast, Mumbai (Maharashtra State) and Veraval (Gujarat State), and Chennai (Tamil Nadu State) along the South east coast (Fig. 1). The sampling sites were selected as per the information available from the reports of Central Marine Fisheries Research Institute (CMFRI) and the number of sampling sites was limited within the scope of the study. A total of 303 fish Samples were collected randomly during October to December 2012. Fish samples were collected from the commercial landings after obtaining information about fishing area from the fishermen. The sampling period was designed based on the spawning season of the fish, so that the mixing of the putative stock is less during that period¹⁴. The peak spawning season of *N. bipunctatus* was reported from September to January along the Vizhinjam coast¹² and a prolonged spawning season from October to February off Tuticorin coast¹⁵.

Digitization of fish samples

The samples were first cleaned in running water, wiped with clean thin cotton towel and placed on a flat platform with laminated graph paper, keeping left




Fig. 1 — Location selected for sampling of Delagoa thread fin bream, *Nemipterus bipunctatus*

side up. The graph paper with fixed horizontal and vertical grids was used in calibrating the coordinates of digital images. The fins were kept erect by pinning so as to make the origin and insertion points clear, which is essential while selecting homologous landmarks around the outline of fish form. A block of expanded polystyrene (2 cm thickness) was placed beneath the graph sheet to facilitate pinning. Digital images of each fish sample were captured using a Sony cyber shot DSC-S300 digital camera (Sony, Japan).

Collection and digitization of fish skull

The head of fish samples was lopped after the digitization of fish samples and kept in 95 % alcohol for hardening. It was then boiled just enough to loosen the muscles and washed in a mild jet of water to remove the loosened muscles and skin. The skull was carefully removed and air dried without exposing it to direct heat or sun light. Air dried skulls were transferred to 1 % KOH in distilled water and kept for 24 h. It was then washed thoroughly again in mild pressure of water and air dried properly before staining. A stock dye of Alizarin Red was prepared for staining¹⁶. Dried skulls were transferred to 1 % KOH and Alizarin Red was added drop wise (3-4 drop). The stained skulls were kept for three days for drying.

Out of 303 fish samples, 231 intact skulls could be obtained, which were used for further analysis. For digitization the stained skull was placed on a flat platform with laminated graph paper in such a way as to obtain the dorsal view of the skull. A light source was provided below and above the laminated plane to improve the clarity of the image. Digital images of each skull were captured using a Sony cyber shot DSC-S300 digital camera (Sony, Japan).

Measurement of truss distances of body and skull

The geometric morphometric analysis called truss network analysis based on a series of truss network measurements that form a regular pattern of connected quadrilaterals, was selected for both body and skull of fish. These measurements were calculated on the basis of morphologically significant anatomical locations on fish body or skull called as 'landmarks'. The landmarks were digitized on the image using the software "tpsDig2 V2.1"¹⁷ and the data were encrypted into the tps files in the form of X-Y coordinates. A total of twenty truss distances were measured along the entire body surface on left side of the fish based on nine landmarks (Fig. 2). For

the digitized skull image another truss network was formed by colligating 11 landmarks, forming 21 truss distances (Fig. 3). Paleontological Statistics (PAST) software¹⁸ was used to extract the truss distances between pre-determined anatomical landmarks on each fish and skull specimen.

Analysis of truss morphometric data

The truss distances were first tested for outliers, based on Cook's distance estimates using PROC ROBUSTREG procedure of SAS Version 9.3¹⁹ to




Fig. 2 — Truss network series of body with interconnected landmarks




Fig. 3 — Truss network series of skull with interconnected landmarks

remove potential outliers from further analysis. A total of 23 fish samples were excluded from further analysis as they may have led to biased inferences. There were significant correlations between the standard length of fish and truss distances on the body and the standard skull length and truss distances of the skull. To overcome the length dependency, the transformation of absolute measurements into size independent shape variables was carried out by a modified formula originally given by Ihssen *et al.*¹ and Hurlburt and Clay²⁰:

$$D_{\text{trans}} = D \times \left(\frac{SL_{\text{mean}}}{SL} \right)^b$$

where,

D_{trans} : transformed truss distance

D : original truss distance

SL : standard length of fish*

SL_{mean} : mean standard length

b : Within-group slope of the regression of $\log D$ on $\log SL$

* Standard skull length is used for transformation of skull truss measurements

The effectiveness of transformation of truss distances of body and skull was ascertained by estimating the correlation coefficients of transformed truss measurements with the standard length of fish and standard skull length data, respectively. The transformed truss distances from body and skull were subjected to canonical discriminant function analysis using the CANDISC procedure of SAS¹⁹ to extract linear combinations of the canonical variables to obtain the maximum differences between various groups. Cumulative variance explained by the canonical axes and biologically meaningful groupings of the traits loaded on each canonical axis were taken into consideration for grouping the samples.

The truss distances loaded on the first two canonical axes were selected based on the Hatcher's scratching procedure²¹. The Artificial Neural Network (ANN) analysis was employed using NEURAL MODEL option in JMP 8¹⁹ to find out the accuracy of classifications of observations among different locations and coasts on the basis of the first two canonical axes scores. ANN is a nonlinear mathematical structure capable of representing complex nonlinear process that relates the inputs to the outputs of a system²². The analysis was separately carried out for transformed body truss distances and transformed skull truss distances. The structure of a neural net

consists of connected units referred to as “nodes” or “neurons”. Each neuron performs a portion of the computations inside the net. A neuron takes some numbers as inputs, performs a relatively simple computation on these inputs, and returns as an output. Output value of a neuron is passed on as one of the inputs for another neuron, except for neurons that generate the final output values of the entire system. In the present study, we used a single layered feed forward network (SLFN) of 3 hidden nodes with the following specifications (Fig. 4).

- 1) Input layer: The first two canonical axes scores (Can1 and Can 2)
- 2) Hidden layer: 3 hidden nodes; H1, H2 and H3
- 3) Output: Region (Location/coast)
- 4) Over all penalty =0.01
- 5) Number of tours = 8
- 6) Iterations = 500
- 7) Training data set = 65 % of the observations
- 8) Cross validation = Random Hold back

Results

The correlation coefficients between size adjusted variables and standard length reduced considerably, indicating that the data transformation was effective in removing the effect of size. In the present study, the mean standard length of all the three populations was significantly different from each other (Table 1). The canonical discriminant analysis on truss distances of body and skull revealed that there is a significant variation between stocks from all the different sampling locations and the variables with high loadings on the first two canonical axes were found useful in distinguishing these samples. The ANN analysis based on first two canonical scores of the body truss measurements differentiated the




Fig. 4 — Single layered feed forward artificial neural network with 3 hidden nodes (H1, H2 and H3).

individuals to their respective populations with an accuracy of 91.4 % and the accuracy of classification based on skull truss distances was 86.14 %.

Differentiation of the stocks based on truss network analysis of body

Eigen value for first canonical axis was 2.22 and that for the second canonical axis was 0.9238 and they explained 70 and 22 percent of the between group variations, respectively. The pooled within group correlation of canonical axes revealed that the truss distances 1-2, 2-3, 2-6, 2-7, 2-8, 3-4 and 3-5 contributed to the first canonical axis (Can 1) and truss distances 3-7, 3-8, 3-9, 4-5, 4-6, 5-6 and 6-7 contributed to the second canonical axis. The bivariate scatter plot of canonical axes (Can1 and Can2) figured out a better discrimination of the *N. bipunctatus* stocks from three locations with limited mixing among each other (Fig. 5a).

Differentiation of the stocks based on truss network analysis of skull

Eigen value for first canonical axis was 0.9088 with a between group variability of 62 % and the Eigen value for the second canonical axis was 0.5505 with a between group variability of 35 %. In Canonical discriminant analysis for skull the truss distances 1-2, 1-11, 3-4, 3-9, 4-10 and 9-10, which are truss distances from the olfactory region and anterior portion of orbital region of the skull contributed to the first canonical axis (can 1). The truss distances contributed to second canonical axis (can 2) are; 3-11, 4-5, 4-8, 5-6, 5-7, 6-7, 6-8, 7-8 and 8-9 which are on orbital and otic region of the skull. The bivariate scatter plot of the canonical axes (Can1 and Can2) depicted a better discrimination between the fish stocks with limited overlapping (Fig. 5b).

Classification by ANN

The first two canonical axes scores were taken for classifying samples among different stocks. They clearly discriminated fishes to their respective locations, with an accuracy of 91.4 % (Table 1). Similarly, for the skull specimens, accuracy of classification is 86.14 % (Table 1).

Discussion

The canonical discriminant analysis of transformed truss distances of body and skull showed that there is a significant morphometric heterogeneity among *N. bipunctatus* stocks from the three selected locations.




Fig. 5a — Stock wise canonical discriminant plot between first and second canonical coefficient for the truss distances of fish body.




Fig. 5b — Stock wise canonical discriminant plot between first and second canonical coefficient for the truss distances of skull

The classification using ANN has also confirmed the above result. The truss distances are transformed into size independent shape variables during statistical analysis as the morphometric variations should be attributable only to the body shape differences, and not relative to the size of the fish²³. There are no significant differences between the male and female specimen of the species and that rejected the chance of occurrence of sexual dimorphism in the species. Therefore, all the specimen was clubbed together for the truss network analysis as the shape difference was not evident between male and female specimen.

Morphological difference between the three fish stocks may be attributed to the difference in their location of existence²⁴. Organisms of same species ensue morphological variations in wild while experiencing different ecological factors²⁵. This may be due to extended use of morphological structures in that particular condition²⁶ or an important adaptive strategy for populations experiencing inconsistent environments^{27,28}. Moreover, phenotypically plastic traits of marine organisms are potentially affected by environmental factors such as temperature and salinity, which preclude adaptation to a particular habitat²⁹. Difference in physical and ecological conditions are evident from Bay of Bengal and Arabian Sea situated along the East and West coast of India, respectively³⁰⁻³³. These divergent environmental conditions may cause morphological variations to the fish species living in that habitat.

Table1 — Cross-validation of individuals classified by neural network analysis. Percentage of fish from each location or coast (in rows) classified by ANN to their respective groups (in columns) based on body truss and skull truss distances, collected from the respective locations.

Location	n	SL*	Body truss distances			Total rate of classification (%)	Total rate of misclassification (%)
			Veraval	Mumbai	Chennai		
Veraval	100	15.45 ^a	90.5	3.2	6.3	91.4	8.6
Mumbai	103	14.41 ^b	7.7	91.2	1.1		
Chennai	100	13.59 ^c	6.4	1.1	92.5		
Location	n	SKL**	Skull truss distances			Total rate of classification (%)	Total rate of misclassification (%)
			Veraval	Mumbai	Chennai		
Veraval	71	3.23 ^a	85.3	9.8	4.9	86.1	13.9
Mumbai	75	2.96 ^b	12.6	81.3	6.1		
Chennai	85	2.72 ^c	2.7	5.3	92		

*SL - The mean standard length in cm, n - the total number of samples

**SKL - The mean skull standard length in cm

*means bearing the same superscript within a subgroup are not significantly different from each other (p ≤ 0.01).

Body Morphometry

The analysis of the shape morphometry of the body revealed that the majority of the variations were associated with the traits of head region and the posterior body portion including the caudal region. The morphometric variations in the anterior body portion and caudal region was reported in the stocks of *Megalaspis cordyla* populations in Arabian Sea and Bay of Bengal based on the truss network analysis⁵. The variations in the head region may be attributed to the change in feeding pattern of the fish³⁴. *N. bipunctatus* is a carnivore fish which mainly feeds on benthic crustaceans, finfishes, cephalopods and polychaete worms³⁵. The study on the feeding biology from west coast of India more clearly depict that this species is mainly a zoo-benthic feeder and occasionally column feeder³⁶. Spatial variation in the feeding pattern of a related species *N. japonicus* is reported along East and West coasts of India³⁴. Moreover, the variation of feeding pattern of fishes totally depends on the food availability and environmental conditions at a particular habitat³⁷.

Head related variations also suggest the impact of diverse ecological conditions to which the populations are exposed³⁸. Morphological variation on the caudal region may be attributed to the variation in turbulence of water or the fluctuation in the temperature and salinity of the water^{39,40}. The Bay of Bengal is more turbulent than Arabian Sea⁴¹ and other hydrological conditions such as water temperature, salinity, productivity, etc., also vary between the Arabian Sea and Bay of Bengal^{42,43}.

Skull Morphometry

The analysis of skull shape was exercised widely on many major vertebrate phyla for studying the spatio-temporal variations in the morphology⁴⁴⁻⁴⁶, but much work was not done on truss network analysis of fish skull. Also, it is clearly pointed out that the skull can be chosen for morphometric investigation because of its presumed ability to reflect any local adaptation in feeding biology with respect to different environment⁴⁷. Here the truss network analysis of skull clearly depicts a significant difference in the anterior and otic regions of skull. As *N. bipunctatus* is a carnivore's fish species, the different feeding strategies in changing ecological conditions may bring changes to skull structures. Various researches also suggest that predatory or carnivore's fishes may show some morphometric variation on the skull as an adaptive strategy for prey capture and drag reduction

in different habitats^{48,49}. In addition to feeding pattern of fishes, the influence of environmental parameters of the water body may also bring shape variations on the skull^{50,51}. The latitudinal and longitudinal variation of environmental parameters are evident from the literatures on Arabian Sea and Bay of Bengal as discussed early. A significant difference in the skull morphometrics of the different stocks of oceanodromous Baltic Sea herring from different spawning locations, across various temperature gradients were also reported⁴⁷. Moreover, the truss network analysis of the entire body of *N. bipunctatus* discloses a significant variation in head region among different stocks. The possibilities of variations in the morphometry of skull due to changes in head region were reported in aquatic animals⁵². The present study using truss network analysis revealed more complex variation in skull structure that can be an evidence for the occurrence of heterogeneous stocks.

Conclusion

Significant variation in the phenotype of *N. bipunctatus* based on the truss network analysis of whole body of fish and skull well elucidate the existence of three different stocks along the Indian coast. The occurrence of same species in multifarious environmental condition may leads to the variation in the phenotypes, which is argued to be the optimal strategy to suit the local environment. Moreover, truss network analysis of the skull is performed for the first time in fish. The result of the present work is useful in developing the preliminary guidelines for the assessment of exploited populations of *N. bipunctatus*. It is mattering much to acquire the necessary knowledge about the stock structure of a commercially exploited species for proper management before exploitation proceeds too far.

Acknowledgement

Authors like to thank the Director and faculty of ICAR-Central Institute of Fisheries Education (Indian Council of Agricultural Research), Mumbai, India, as the study was carried out with financial and technical support from the institute. We extend our thanks to the faculty of the Central Marine Fisheries Research Institute, India for their help and support throughout the sampling period.

References

- 1 Ihssen, P.E., Booke, H.E., Casselman, J.M., McGlade, J.M., Payne, N.R. and Utter, F.M., Stock identification: materials

- and methods. *Can. J. Fish. Aquat. Sci.*, 38 (12) (1981) 1838-1855.
- 2 Devaraj, M. and Vivekanandan, E., Marine capture fisheries of India: Challenges and opportunities. *Curr. Sci.*, 76(3) (1999) 314-332.
 - 3 Larkin, P.A., A perspective on population genetics and salmon management. *Can. J. Fish. Aquat. Sci.*, 38 (1981) 1469-1475.
 - 4 Turan, C., Stock identification of Mediterranean horse mackerel (*Trachurus mediterraneus*) using morphometric and meristic characters. *ICES J. Mar. Sci.*, 61(5) (2004) 774-781.
 - 5 Sajina, A.M., Chakraborty, S.K., Jaiswar, A.K., Pazhayamadam, D.G. and Sudeshan, D., Stock structure analysis of *Megalaspis cordyla* along Indian coast based on truss network analysis. *Fish. Res.*, 108 (2011) 100-105.
 - 6 Sen, S., Jahageerdar, S., Jaiswar, A.K., Chakraborty, S.K., Sajina, A.M. and Dash, G.R., Stock structure analysis of *Decapterus russelli* (Ruppell, 1830) from east and west coast of India using truss network analysis. *Fish. Res.*, 112 (1-2) (2011) 38-43.
 - 7 Canas, L., Stransky, C., Schlickeisen, J., Sampedro, M.P. and Farina, A.C., Use of the otolith shape analysis in stock identification of anglerfish (*Lophius piscatorius*) in the Northeast Atlantic. *ICES J. Mar. Sci.*, 69 (2012) 250-256.
 - 8 Booke, H.E., The conundrum of the stock concept- are nature and nurture definable in fishery science. *Can. J. Fish. Aquat. Sci.*, 38 (1981) 1479-1480.
 - 9 Dwivedi A.K. and Dubey V.K., Advancements in morphometric differentiation: a review on stock identification among fish populations. *Rev. Fish. Biol. Fisher.*, 23 (2013) 23-39.
 - 10 Strauss, R.E. and Bookstein, F.L., The truss: body form reconstructions in morphometrics. *Syst. Zool.*, 31 (1982) 113-135.
 - 11 CMFRI, *Annual Report 2012-13*. (Central Marine Fisheries Research Institute, Cochin) 2013, pp. 200.
 - 12 Mohan, M. and Velayudhan, A.K., Spawning biology of *N. delagoae* (Smith) at Vizhinjam. *Indian J. Fish.*, 28 (1 & 2) (1986) 26-34.
 - 13 Kizhakudan, S.J. and Rajapackiam, S., Length-weight relationship in six species of threadfin breams occurring in the trawl landings at Chennai. *J. Mar. Biol. Assoc. India*, 53 (2) (2011) 268 - 271.
 - 14 Cadrin, S.X., Advances in morphometric identification of fishery stocks. *Fish. Biol. Fisher.*, 10 (2000) 91-112.
 - 15 Hamsa, K.M.S.A., Kasim, M.H. and Arumugam, G., The fishery, biology and stock assessment of *Nemipterus delagoae* smith off Tuticorin, Gulf of Mannar. *Bulletin of Central Marine Fisheries Research Institute*, 47 (1994) 112-120.
 - 16 Hollister, G., Clearing and drying of fish for bone study. *Zoologica*, 12 (1934) 89-101.
 - 17 Rohlf, F.J., *tpsDig, Digitize Landmarks and Outlines, Version 2.05*. (Stony Brook, NY: Department of Ecology and Evolution, State University of New York) 2006.
 - 18 Hammer, O., Harper, D.A.T. and Ryan, P.D., PAST: Paleontological Statistics Software Package for Education and Data Analysis. *Palaentologia Electronica*, 4(1) (2001) 9.
 - 19 SAS Institute Inc., *What's New in SAS® 9.3.*, (Cary, NC: SAS Institute) 2012, pp. 272.
 - 20 Hurlburt, T. and Clay, D., Morphometric and meristic differences between shallow and deep-water population of white hake (*Urophycis tenuis*) in the southern gulf of St. Lawrence. *Can. J. Fish. Aquat. Sci.*, 55 (1998) 2274 - 2282.
 - 21 Hatcher, L., *A Step by Step Approach to Using SAS for Factor Analysis and Structural Equational Modeling*, (SAS Institute Inc., Cary, NC) 2003, pp. 57-125.
 - 22 Pulido-Calvo, I. and Portela, M.M., Application of neural approaches to one-step daily flow forecasting in Portuguese watersheds. *J. Hydrol.*, 332 (2007) 1-15.
 - 23 Turan, C., A note on the examination of morphometric differentiation among fish populations: The Truss System. *Turk. J. Zool.*, 23 (1999) 259-264.
 - 24 Barlow, G.W., Causes and significance of morphological variations in fishes. *Syst. Biol.*, 10(3) (1961) 105-117.
 - 25 Sreekanth, G.B., Chakraborty, S.K., Jaiswar, A.K., Renjith, R.K., Ratheesh Kumar, Sandeep, K.P., Vaisakh, G., Sunil, S.A., Manju Lekshmi, N., and Deepak, G.P., Can the *Nemipterus japonicus* stocks along Indian coast be differentiated using morphometric analysis? *Indian J. Mar. Sci.*, 44 (4) (2015).
 - 26 Ehlinger, T.J., Habitat choice and phenotype-limited feeding efficiency and habitat choice in bluegill: Individual differences and trophic polymorphism. *Ecology*, 71 (1990) 886-896.
 - 27 Stearns, S.C., A natural experiment in life-history evolution: field data on the introduction of mosquitofish (*Gambusia affinis*) to Hawaii. *Evolution*, 37(3) (1983) 601-617.
 - 28 Wimberger, H.P., Plasticity in the skull morphology of the Neotropical cichlids, *Geophagus brasiliensis* and *G. steindachneri*. *Evolution*, 45(7) (1991) 1545-1563.
 - 29 Guinand, B., Lemaire, C. and Bonhomme, F., How to detect polymorphisms undergoing selection in marine fishes? A review of methods and case studies, including flatfishes. *J. Sea Res.*, 51 (2004) 167- 182.
 - 30 Jayaraman, R. and Gogate, S.S., *Salinity and temperature variations in the surface waters of the Arabian Sea off the Bombay and Saurashtra Coasts*. (Proceedings of the Indian Academy of Sciences - Section B, 45(4)) 1957, pp. 151-164.
 - 31 Balaramamurty, C., *Studies in Physical oceanography of the Western Bay of Bengal*. Ph.D. Thesis, Andhra University, Waltair, 1958.
 - 32 Kumar, S.P., Narvekar, J., Nuncio, M., Kumar, A., Ramaiah, N., Sardesai, S., Gauns, M., Fernandes, V. and Paul, J., Is the biological productivity in the Bay of Bengal light limited? *Curr. Sci.*, 98 (10) (2010) 1331-1339.
 - 33 Pakajakshan, T. and Gosh, A.K., Surface Layer Temperature Inversion in the Arabian Sea during Winter. *J. Oceanogr.*, 48(3) (1992) 293-304.
 - 34 Sreekanth G.B., Deepak G Pazhayamadom, Sushanta Kumar Chakraborty, Ashok Kumar Jaiswar, Renjith, R.K., Ratheesh Kumar R., Anuraj, A., Site specific differences in food and feeding biology of *Nemipterus japonicus* (Bloch, 1791) along Indian coast. *Indian J. Fish.*, 59 (3) (2012) 25-31.
 - 35 Froese, R. and Pauly, D., Editors. FishBase. World Wide Web electronic publication. www.fishbase.org, version (02/2018).
 - 36 Muthiah, C. and Pillai, S.K., A new distributional record of *N. delagoae* (Smith) from Bombay waters with notes on its biology. *J. Mar. Biol. Assoc. India*, 21 (1-2) (1979) 174- 178.

- 37 Singh, C.P., Ram, R.N. and Singh, R.N., Food and feeding pattern of *Channa punctatus* in two different habitats at Tarai region of Uttarakhand. *J. Environ. Biol.*, 34 (2012) 789-792.
- 38 Palma, J. and Andrade, J.P., Morphological study of *Diplodus sargus*, *Diplodus puntazzo*, and *Lithognathus mormyrus* (Sparidae) in the Eastern Atlantic and Mediterranean Sea. *Fish. Res.*, 57 (2002) 1-8.
- 39 Imre, I., McLaughlin, R.L. and Noakes, D.L.G., Phenotypic plasticity in brook charr: changes in caudal fin induced by water flow. *J. Fish Biol.*, 61 (2002) 1171-1181.
- 40 Omoniye, T. and Agbon, A.O., Morphometric Variations in *Sarotherodon melanotheron* (Pisces: Cichlidae) from Brackish and Fresh Water Habitats in South-western Nigeria. *West African J. Appl. Ecol.*, 12 (2007) 89-95.
- 41 Chamarthi, S., Ram, P.S. and Josyula, L., Effect of river discharge on Bay of Bengal circulation. *Mar. Geodesy*, 31 (3) (2008) 160-168.
- 42 Gomes, H.R., Goes, J.I. and Saino, T., Influence of physical processes and freshwater discharge on the seasonality of phytoplankton regime in the Bay of Bengal; *Cont. Shelf. Res.*, 20 (2000) 313-330.
- 43 Madhupratap, M., Gauns, M., Ramaiah, N., Prasanna Kumar, S., Muraleedharan, P.M., de Souza, S.N., Sardesai, S. and Usha, M., Biogeochemistry of the Bay of Bengal: Physical, chemical and primary productivity characteristics of the central and western Bay of Bengal during summer monsoon 2001. *Deep Sea Res.*, 50(2) (2003) 881-896.
- 44 Wehausen, J.D. and Ramey, R.R., Cranial morphometric and evolutionary relationships in the northern range of *Ovis Canadensis*. *J. Mammal.*, 81(1) (2000) 145-161.
- 45 Gomercic, T., Farkas, V., Gomeric, M.D., Huber, D., and Gomercic, H., *Cranial morphometry of adult Mediterranean monk seal (Monachusmonachus) from the Adriatic Sea*. (Proceedings of the international scientific meeting of Anatomy and Physiology., June 12-13-2009, Zagreb, Croatia) 2009, pp 120-128.
- 46 Wiig, O., Jorgensen, M.P.H., Laidre, K.L., Garde E. and Reeves, R.R., Geographic variation in cranial morphology of narwhals (*Monodon monoceros*) from Greenland and the eastern Canadian Arctic. *Polar Biol.*, 35(1) (2011) 63-71.
- 47 Jorgensen, H., Cino, P., Micheal, M.H., Daniel, E.R. and Volker, L., Genetic and environmental correlation of morphological variation in marine fish: The case of Baltic Sea Herring (*Clupea Harengus*). *Can. J. Fish. Aquat. Sci.*, 65 (3) (2008) 389-400.
- 48 De Schepper, N., De Kegel, B. and Adriaens, D., Morphological specialization in Heterocongrinae (Anguilliformes: Congridae) related to burrowing and feeding. *J. Morphol.*, 268 (2007) 343-356.
- 49 Eagderia, S. and Adriaens, D., Head morphology of the duckbill eel, *Hoplunnis punctata* (Regan, 1915; Nettastomatidae: Anguilliformes) in relation to jaw elongation. *Zoology*, 113 (2010) 148-157.
- 50 AL-Harbi, A. H., Skeletal Deformities in Cultured Common Carp, *Cyprinus carpio* L. *Asian Fish. Sci.*, 14 (2001) 247-254.
- 51 Wintzer, A.P. and Motta, P.J., Diet-induced phenotypic plasticity in the skull morphology of hatchery-reared Florida largemouth bass, *Micropterus salmoides floridanus*. *Ecol. Freshw. Fish*, 14 (2005) 311-318.
- 52 Jeferson, T.A., Preliminary analysis of geographic variations in cranial morphometrics of the finless Propoise (*Neophocaena phocaenoides*). *Raffles bull. Zool.*, 10 (2002) 3-14.