

NOTES

Variations in palpal ornamentation of
Anopheles fluviatilis species T and U
(Diptera: Culicidae) and their
taxonomic consequence

Gunjan Sharma, Manila Lather & Om P Singh*

National Institute of Malaria Research, Sector 8, Dwarka,
New Delhi-110 077, India

Received 14 September 2018; revised 16 April 2019

Anopheles fluviatilis sensu lato James is a highly efficient malaria vector in Indian subcontinent and Iran which is comprised of at least four sibling species provisionally designated as species S, T, U and V. An important morphological characteristic for differentiation of this species complex from other closely related anopheline species complex, the Minimus Complex, is the ratio of length of subapical pale band to dark band intervening apical and subapical pale bands on the maxillary palps of female mosquito. Here, we report variation in the subapical pale band in *An. fluviatilis*, especially in species U, to the extent that palps of some specimens resemble members of Minimus Complex, inferring that palpal ornamentation may not be reliable characteristics for identification of *An. fluviatilis*. Taxonomic consequence of such variation is discussed.

Keywords: *Anopheles minimus*, Fluviatilis Complex, hypomelanism, malaria vectors, Minimus Complex, mosquitoes, Vector control

Vector control is a crucial component of malaria prevention, control and elimination strategies¹, the success of which relies on correct identification of vector species². In India, there are six malaria vectors recognized as primary malaria vectors, viz., *An. culicifacies*, *An. fluviatilis*, *An. minimus*, *An. stephensi*, *An. sudaicus* and *An. dirus*³. All of them, except *An. stephensi*, are comprised of several cryptic species³. Traditionally, identification of vector species is performed on the basis of morphological characteristics that is used to differentiate mosquito species². However, there are some reports where vectors have been misidentified owing to overlapping morphological characteristics in closely related species⁴⁻⁷. Identification of vector species is further challenged by the presence of sibling species. Although considerable progress has been made

toward the development of advanced tools for the identification of sibling species of Indian malaria vectors using chromosomal, biochemical and molecular markers^{3,8-12}, but use of such techniques is limited to differentiation of members of a specific species complex only provided they are correctly identified morphologically prior to application of such techniques. It has been observed that incorrect morphological identification of mosquitoes prior to cytotaxonomy⁵ or PCR assay² can lead to misleading result. Therefore, correct morphological identification of mosquitoes prior to application of such techniques is essential².

Morphological differentiation of two closely related vector species complexes, *An. fluviatilis s.l.* and *An. minimus s.l.*, which belongs to Minimus subgroup of *Anopheles*¹³, is often challenging due to overlapping morphological characteristics. As on date, there are four reported sibling species in the Fluviatilis Complex, i.e., species S, T, U and V^{9,12} and three sibling species in the Minimus Complex, i.e., *Anopheles minimus*¹⁴, *An. harrisoni*¹⁵ and *An. yaeyamaensis*¹⁶. Instances exist in literature where *An. minimus s.l.* have been misidentified as *An. fluviatilis s.l.* due to overlapping morphological characteristics⁴⁻⁶, especially the ratio of length of the subapical pale band to dark band intervening apical and subapical pale bands which is an important morphological characteristic for differentiation of these two species. Such misidentification was attributed to hyper-melanisation of palpi of *An. minimus s.l.* mainly in the cooler season⁵, where the length of the subapical pale band is reduced substantially resembling *An. fluviatilis s.l.* However, there is no published report of misidentification of *An. fluviatilis s.l.* as *An. minimus s.l.* Here, we examined a scenario where palps of *An. fluviatilis* may resemble *An. minimus s.l.* which questions the validity of such a morphological characteristic in morphological discrimination of these two species.

Materials and Methods**Mosquito collection and processing**

Adult *An. fluviatilis s.l.* were collected from villages Ismailpur, Dargahpur and Oaspur under Laksar CHC (Community Health Centre) of district

*Correspondence:
E-mail: singh@nimr.org.in

Hardwar, India (latitude 29°N, longitude 78°E, Fig. 1), in the month of March 2014 from cattle sheds and human dwellings using a hand aspirator and flashlight torch in the morning (06:00 to 08:00 h). Presence of three sibling species of *An. fluviatilis* i.e., species T, U and V has been recorded in the past in these villages¹². The outdoor daytime temperature during the collection period was 28-32°C and 17-18°C during night. Mosquitoes were carefully transferred to a polystyrene foam box with one side opening mounted with nylon netting and transported to laboratory at Delhi. Mosquitoes were provided access to 10% glucose-soaked cotton pad during transport. Besides, some of the field mosquitoes with palpi resembling *An. minimus* s.l. (having subapical pale band equal to intervening dark band between apical and subapical pale band) were selectively pinned and preserved. Live mosquitoes were allowed to lay eggs in laboratory and reared till emergence into adult (F₁). Rearing was done at constant room temperature maintained at 25±1°C, close to the prevailing average temperature in collection sites and relative humidity of 70±5%. During rearing, larvae were fed on grinded mixture of dog biscuit and yeast in a ratio of 3:2. Upon pupation, pupae were transferred in bowl containing water and placed into

mosquito cage. Adults were provided access to 10% glucose-soaked cotton pad. Four to five day's old adult female mosquitoes were anesthetized with ether and palpal characteristics were examined under Leica M165-C stereoscopic microscope. Pinned mosquitoes were also examined.

DNA isolation

DNA was isolated from individual mosquitoes following examination of morphological characteristics. Briefly, whole mosquito was initially grinded in 1.5 mL microfuge tube with 50 µL of TE buffer (10 mM Tris-HCl, 1 mM EDTA, pH 8.0) and final volume was made up to 200 µL. This was incubated for 10 min on a heat block maintained at 96°C. Finally, the content was centrifuged at 10000 RPM for 5 min, and the supernatant transferred in another microcentrifuge tube and stored at -20°C.

Sibling species identification and DNA sequencing

For identification of sibling species of *Fluviatilis* Complex, allele-specific PCR (ASPCR) was carried out as described by Singh *et al.*¹⁰. The PCR product was visualized on 2% agarose gel (Fig. 2). For confirmation of ASPCR results, some samples were sequenced for domain D3 of 28S rDNA. For sequencing, samples were amplified using primers



Fig. 1 — Map of Hardwar district, Uttarakhand, showing study villages




Fig. 2 — Gel photograph showing allele-specific PCR product. L=100 bp DNA ladder, T= *An. fluviatilis* species T, U= species U

D3A and D3B as described by Singh *et al.*⁴. The PCR products were sequenced from both strands of DNA using BigDye Terminator v3.1 Cycle Sequencing Kit (Thermo Fisher Scientific).

Results and Discussion

Palpal ornamentation is important taxonomic characteristics for the identification of *An. fluviatilis*, *An. minimus* and *An. varuna*. Christophers (1933)¹⁷ in his monograph “The Fauna of British India, Including Ceylon and Burma” has mentioned that “Distinction (of *An. fluviatilis*) from *An. minimus*, and especially from *An. varuna*, may be difficult if the palpal ornamentation is ambiguous” and that “Specimens with typically marked palpi, which form the great majority, should offer no difficulty”. These statements underline the importance of palpal characteristics in the differentiation of the *An. fluviatilis* s.l. from *An. An. minimus* s.l. The main distinguishing characteristics of *An. fluviatilis* according to Christophers is “black band between apical and the subapical pale band usually four to five times length of the subapical pale band and at least half length of dark area between subapical and more basal pale band”¹⁰. There has been no change on this criterion since then. In our own experience, we often receive misidentified *An. fluviatilis* by field entomologist as *An. minimus*. We realized that such mistakes are due to the presence of a broad subapical band resembling *An. minimus*. We, therefore, examined the presence of such variation in field population of *An. fluviatilis* and their significance in the formal taxonomy.

A total of 53 samples were identified for sibling species by ASPCR assay following Singh *et al.*¹⁰, of which 20 samples were subjected to DNA sequencing for D3 domain of 28S-rDNA for confirmation of ASPCR-based results. DNA sequencing results of six




Fig. 3 — Proportion of mosquitoes with three different classified categories of subapical pale band in *An. fluviatilis* species T and U

Table 1 — Distribution of ratio of subapical pale band to dark band intervening apical and subapical pale bands in members of *An. fluviatilis* species

	Species T	Species U	Total
Randomly sampled (F ₁)			
Classical	17 (85%)	10 (42%)	27
Intermediate	3 (15%)	9 (37%)	12
Broader	0	5 (21%)	5
Total	20	24	44
Selectively sampled (wild caught)			
Broader	0	9 (100%)	9
Grand total			53

specimens of species T and 14 specimens of species U (having broader subapical pale band on maxillary palpi), were in agreement of ASPCR. Of the 53 samples identified for sibling species, 44 samples were randomly sampled from the F₁ generation of field collected mosquitoes and nine pinned specimens which were suspected to be *An. minimus* based on palpal characteristics. Mosquitoes were classified in three categories based on the ratio of width of dark band intervening apical and subapical pale band on the palpi to the width of subapical pale band, i.e., (i) “classical” with ratio <1/3; (ii) “intermediate” with ratio between 1/3 and 3/4; and (iii) “broader” with ratio >3/4 to 1. The distribution of these three categories of palpi in different sibling species has been shown in Table 1. Species V, which was recorded in an earlier study³, was absent in this collection. It was observed that classical subapical pale band was present in the majority of species T (85%) and “broader” category was absent. In contrast, species U showed wide variation in subapical pale band with “broader” category present in 21% of individuals (Fig. 3). All the nine pinned specimens (field collected) with subapical pale band similar to *An. minimus*, i.e., equal to the intervening dark band between apical and subapical pale band




Fig. 4 — Palpal ornamentation of an *An. fluviatilis* species U resembling *An. minimus s.l.* (A) apical pale band; (B) intervening dark area between apical pale band and subapical pale band; (C) subapical pale band

(category “broader”), were found to be species U. Thus, species U tended to have a broader subapical pale band as compared to species T. A photograph of palpal ornamentation in species U with “broader” subapical pale band has been displayed as Fig. 4.

Identification of *An. fluviatilis s.l.* has always been complicated due to overlapping morphological characteristics with *An. minimus* complex and *An. varuna*^{4,6}. Earlier record showed the widespread distribution of *An. fluviatilis* from Yemen to Taiwan¹⁸. However, later work by Harrison⁵, Chen *et al.*⁶ and Singh *et al.*⁴ have reported misidentification of *An. minimus s.l.* as *An. fluviatilis s.l.* resulting from overlapping maxillary palpal characteristics in China, Thailand and a northeast state of India (Assam). Therefore, their presence in countries eastward to India and in north-eastern states of India was very much doubted⁵. As a consequence, Harrison⁵ and Chen *et al.*⁶ have recommended removal of *An. fluviatilis* from fauna-list of Thailand and China, respectively. Interestingly, the chromosomal complements of *An. minimus s.s.* and *An. fluviatilis* species U were found to be homosequential, which was another reason for misidentification of *An. minimus* precisely as *An. fluviatilis* species U⁴. These instances related to misidentification of *An. minimus* as *An. fluviatilis* have been attributed to hyper-melanism in mosquitoes in cooler months⁵. All these reports indicate misidentification of *An. minimus* as *An. fluviatilis* but there is no any report where

An. fluviatilis have been misidentified as *An. minimus* due to hypomelanism. This study reports that such hypomelanism in palpi of *An. Fluviatilis*, especially in the species U, can lead to misidentification of *An. fluviatilis* as *An. minimus*.

Correct identification of vector species is crucial for the success of vector control programme. Due to challenge in the identification of sibling species (which may differ in epidemiologically important biological attributes, such as vectorial competence, insecticide resistance, etc), there has been significant advancement in the development of diagnostic tools for the identification of cryptic species present in various malaria vectors. Application of such tool on incorrectly-identified mosquitoes based on morphological characteristics may be seriously misleading. A recent study² carried out in South Africa reports that when 11 morphological species were subjected to standard PCR used for discrimination of *An. gambiae* complex as well as *An. funestus* group, three morphological species were incorrectly identified belonging to *An. funestus* group and four morphological species were incorrectly identified as member of *An. gambiae* complex. This report signifies importance of morphological identification of vector species in the malaria control programme, especially before applying molecular tool. Similar precaution should be taken before carrying out cytotoxic identification of sibling species. It has been reported that the chromosome complements of *An. fluviatilis* species U is identical to *An. minimus s.s.* and this has resulted in misidentification of *An. minimus s.s.* as *An. fluviatilis* species U⁴. It is thus desired that careful studies should be carried out on formal taxonomy involving morphological, chromosomal and molecular tools together.

Conclusion

The present study reports variation in the subapical pale band in *An. fluviatilis s.l.*, especially in species U, to the extent that palps of some specimens resemble *An. minimus s.l.*, inferring that palpal ornamentation may not be a reliable characteristic for identification of *An. fluviatilis s.l.*

Acknowledgment

Dr Ashish Gupta, National Institute of Malaria Research, Field Unit, Hardwar, for providing logistic support during field visit. Funding for this study was provided by Defence Research Laboratory (DRL),

Tejpur. ML was supported by University Grant Commission (UGC) Senior Research Fellowship.

Conflict of interest

The authors declare no conflict of interest.

References

- 1 WHO, *Guidelines for Malaria Vector Control*. (World Health Organization, Geneva) (2019) 161.
- 2 Erlank E, Koekemoer LL & Coetzee M, The importance of morphological identification of African anopheline mosquitoes (Diptera: Culicidae) for malaria control programmes. *Malar J*, 17 (2018) 43.
- 3 Dash AP, Adak T, Raghavendra K & Singh OP, The biology and control of malaria vectors in India. *Curr Sci*, 92 (2007) 1571.
- 4 Singh OP, Nanda N, Dev V, Bali P, Sohail M, Mehrunnisa A, Adak T & Dash AP, Molecular evidence of misidentification of *Anopheles minimus* as *Anopheles fluviatilis* in Assam (India). *Acta Tropica*, 113 (2010) 241.
- 5 Harrison BA, Medical entomology studies - XIII. The Myzomyia Series of *Anopheles (Cellia)* in Thailand, with emphasis on intra-interspecific variations (Diptera: Culicidae). *Contributions of the American Entomological Institute*, 17 (1980) 1.
- 6 Chen B, Harbach RE & Butlin RK, Molecular and morphological studies on the *Anopheles minimus* group of mosquitoes in southern China: taxonomic review, distribution and malaria vector status. *Med Vet Entomol*, 16 (2002) 253.
- 7 Van Bortel W, Harbach RE, Trung HD, Roelants P, Backeljau T & Coosemans M, Confirmation of *Anopheles varuna* in vietnam, previously misidentified and mistargeted as the malaria vector *Anopheles minimus*. *Am J Trop Med Hyg*, 65 (2001) 729.
- 8 Subbarao SK, *Anopheline species complex in South-East Asia*. (SEARO Technical Publication No. 57, WHO Regional Office for South-East Asia, New Delhi) (1998) 102.
- 9 Subbarao SK, Nanda N, Vasantha K, Dua VK, Malhotra MS, Yadav RS & Sharma VP, Cytogenetic evidence for three sibling species in *Anopheles fluviatilis* (Diptera: Culicidae). *Ann Entomol Soc Am*, 87 (1994) 116.
- 10 Singh OP, Chandra D, Nanda N, Raghavendra K, Sunil S, Sharma SK, Dua VK & Subbarao SK, Differentiation of members of the *Anopheles fluviatilis* species complex by an allele-specific polymerase chain reaction based on 28S ribosomal DNA sequences. *Am J Trop Med Hyg*, 70 (2004) 27.
- 11 Singh OP, Goswami G, Nanda N, Raghavendra K, Chandra D & Subbarao SK, An allele-specific polymerase chain reaction assay for the differentiation of members of the *Anopheles culicifacies* complex. *J Biosci*, 29 (2004) 275.
- 12 Nanda N, Singh OP, Dua VK, Pandey AC, Nagpal BN, Adak T, Dash AP & Subbarao SK, Population cytogenetic and molecular evidence for existence of a new species in *Anopheles fluviatilis* complex (Diptera: Culicidae). *Infect Genet Evol*, 13 (2013) 218.
- 13 Harbach RE, The classification of genus *Anopheles* (Diptera: Culicidae): a working hypothesis of phylogenetic relationships. *Bull Entomol Res*, 9 (2004) 537.
- 14 Harbach RE, Parkin E, Chen B & Butlin RK, *Anopheles (Cellia) minimus* Theobald (Diptera: Culicidae): Neotype designation, characterization, and systematics. *Proc Entomol Soc Washington*, 108 (2006) 198.
- 15 Harbach RE, Garros C, Manh ND & Manguin S, Formal taxonomy of species C of the *Anopheles minimus* sibling species complex (Diptera: Culicidae). *Zootaxa*, 1654 (2007) 41.
- 16 Somboon P, Rory A, Tsuda Y, Takagi M & Harbach RE, Systematics of *Anopheles (Cellia) yaeyamaensis* sp. n., alias species E of the *An. minimus* complex of southeastern Asia (Diptera: Culicidae). *Zootaxa*, 6 (2010) 43.
- 17 Christophers SR, *The Fauna of British India, Including Ceylon and Burma, Diptera, Volume IV, Family Culicidae, Tribe Anophelini*. (Taylor and Francis, Red Lion Court, Fleet Street, London) (1933) 371.
- 18 Knight KL & Stone A, *A catalog of the mosquitoes of the world (Diptera: Culicidae), Second edition* (Thomas Say Foundation, Entomological Society of America), 6 (1977) 611.