




Patients` inability to perform a pre-operative cardiopulmonary exercise test or demonstrate an anaerobic threshold is associated with inferior outcomes after major colorectal surgery



Journal:	<i>British Journal of Anaesthesia</i>
Manuscript ID:	BJA-2013-00078-PM009.R1
Manuscript Type:	Clinical Investigation
Date Submitted by the Author:	28-Mar-2013
Complete List of Authors:	Lai, Chee Wan; Plymouth Hospital NHS Trust, Department of Colorectal Surgery; Plymouth University, Peninsula Schools of Medicine and Dentistry Minto, Gary; Plymouth Hospitals NHS Trust, Department of Anaesthesia; Plymouth University, Peninsula Schools of Medicine and Dentistry Challand, Christopher; Plymouth Hospital NHS Trust, Department of Colorectal Surgery; Plymouth University, Peninsula Schools of Medicine and Dentistry Hosie, Kenneth; Plymouth Hospital NHS Trust, Department of Colorectal Surgery Sneyd, J Robert; Plymouth University, Peninsula Schools of Medicine and Dentistry; Plymouth Hospitals NHS Trust, Department of Anaesthesia Creanor, Siobhan; Plymouth University, Peninsula Schools of Medicine and Dentistry Struthers, Richard; Plymouth Hospitals NHS Trust, Department of Anaesthesia; Plymouth University, Peninsula Schools of Medicine and Dentistry
Key Words:	Surgery - abdominal, Assessment - preanaesthetic, Cardiorespiratory system - responses

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Peer Review

Figure 1. Kaplan-Meier Plot for survival after surgery stratified by CPET risk group.
Log-Rank Test $p < 0.001$

Days from Surgery	Number at Risk				
	0	200	400	600	730
Fit	174	169	165	157	155
Unfit	69	64	60	56	55
Unable	26	19	17	15	15

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

CWL, GM CPC, KBH, JRS, SC RAS, March 2013

15

1
2
3 **Patients` inability to perform a pre-operative cardiopulmonary exercise test or demonstrate**
4 **an anaerobic threshold is associated with inferior outcomes after major colorectal surgery**
5

6
7 C. W. Lai^{1 3}, G. Minto^{2 3}, C. P. Challand^{1 3}, K. B. Hosie¹, J. R. Sneyd^{2 3}, S. Creanor³, R. A. Struthers²
8
9

10 ¹Dept of Colorectal Surgery, Derriford Hospital, Plymouth, Great Britain

11 ²Dept of Anaesthesia, Derriford Hospital, Plymouth, Great Britain

12 ³Plymouth University Peninsula Schools of Medicine and Dentistry, Plymouth, Great Britain
13
14

15 Correspondence to:

16 R. A. Struthers

17 richardstruthers@nhs.net

18
19 Anaesthetic Department
20 Plymouth Hospitals NHS Trust
21 Derriford Hospital
22 Derriford Road
23 Derriford, Plymouth
24 PL6 8DH
25
26
27
28

29 Word count: abstract 246; manuscript 2223

30 Type of article: Original article
31
32
33

34 Running Title: Unable to CPET.
35
36

37 GM, RS, JRS, KH and CC conceived the idea for this audit. CWL, CC and RS performed data
38 collection and CWL, RS, GM and SC performed data analysis. CWL, RS and GM drafted the
39 manuscript and all authors were actively involved in revising the manuscript.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

CWL, GM CPC, KBH, JRS, SC RAS, March 2013

15

Abstract

Background: Surgical patients with poor functional capacity, determined by oxygen consumption at Anaerobic Threshold (AT) during Cardiopulmonary Exercise Testing (CPET), experience longer hospital stays (LOS) and higher short and medium term mortality. However previous studies excluded patients who were unable to perform a CPET or who failed to demonstrate an AT. We hypothesized that such patients are at risk of inferior outcomes after elective surgery.

Method: All patients scheduled for major Colorectal Surgery attempted CPET to assist in the planning of care. Patients were stratified by their test results into Fit ($AT \geq 11.0 \text{ mlO}_2\text{kg}^{-1}\text{min}^{-1}$), Unfit ($AT < 11.0 \text{ mlO}_2\text{kg}^{-1}\text{min}^{-1}$), or Unable to CPET groups (failed to pedal or demonstrate an AT). For each group we determined LOS and mortality.

Results: Between March 2009 and April 2010, 269 consecutive patients were screened, and proceeded to bowel resection. Median LOS was 8 days (IQR 5.1-13.4) and there were 44 deaths at 2 years (16%). 26 (9.7%) patients were categorized as Unable to CPET, 69 (25.7%) Unfit and 174 (64.7%) Fit. There were statistically significant differences between the three groups in LOS (median (IQR) 14.0 (10.5-23.8) vs. 9.9 (5.5-15) vs. 7.1 (4.9-10.8) days, $p < 0.001$) and mortality at two years (11/26 (42%) vs. 14/69 (20%) vs. 19/174 (11%) respectively ($p < 0.001$)) although the differences between Unable and Unfit were not statistically different.

Conclusion: Patients' inability to perform CPET is a marker of inferior outcome following major colorectal surgery. Future studies evaluating CPET in risk assessment for major surgery should report outcomes for this subgroup.

Key Words: Exercise Test, General Surgery, Length of Stay, Mortality, Preoperative care, Risk assessment

Introduction

In the UK, elective surgical procedures carry an in-hospital mortality of 0.4%. However, the figures conceal that a large proportion of the mortality is concentrated in a group of high risk patients undergoing major surgery.¹

Recent prominent reports have highlighted deficiencies in the identification of the high risk surgical patient and appropriate planning of their peri-operative care. Patients may be scheduled for major procedures without having a full appreciation of the risk of mortality and severe complications that they face.^{2,3}

The Royal College of Surgeons report "Peri-operative Care of the Higher Risk General Surgical Patient" recommends the use of scoring systems to estimate patients' risk of mortality and guide escalation of care for those with an expected mortality >5%.³

Major surgery triggers a strong systemic inflammatory response, which leads to an increase in oxygen demand. This is met by an increase in global oxygen delivery (DO_2), through an increase in cardiac output (CO) and increase in tissue oxygen extraction. Whilst most patients are able to mount an adequate response, there is a group who may not have the required physiological reserve. This group is at a higher risk of morbidity and mortality when undergoing major surgery.^{4,5}

Cardiopulmonary exercise testing (CPET) examines the ability of the cardio-respiratory system to deliver oxygen to tissues under stress and provides an objective determination of functional capacity and impairment. Anaerobic threshold (AT) is the point during exercise when anaerobic metabolism supplements aerobic with additional CO₂ production, creating an inflection point on a plot of pulmonary CO₂ efflux vs. O₂ uptake. Lower AT indicates lower aerobic fitness and is associated with increased post-operative mortality and severe morbidity. However, CPET studies to date have not reported the peri-operative outcomes of those unable to perform a test or in whom an anaerobic threshold (AT) was not determined.⁶⁻¹¹ Patients with heart failure who do not demonstrate an anaerobic threshold are at higher risk of major cardiac events. This may be due to short duration of tests, abnormal ventilatory patterns, which attenuate the inflection point or alterations in cardiovascular physiology, which conceal it in the early part of the test.¹² We hypothesized that

CWL, GM CPC, KBH, JRS, SC RAS, March 2013

15

patients undergoing surgery who do not demonstrate an AT may similarly be at high risk of adverse outcomes.

Methods

All patients scheduled for elective major colorectal surgery at our Colorectal Specialist Unit attempt pre-operative CPET to assist in the planning of peri-operative care. With Ethics Committee approval all patients undergoing bowel resection between March 2009 and April 2010 were screened for inclusion into a trial of intraoperative goal-directed fluid therapy and those with an $AT > 8 \text{ mlO}_2\text{kg}^{-1}\text{min}^{-1}$ were approached to be randomized. The testing methods and results of this study have already been reported.¹¹ CPET was performed on a stationary bicycle (Zan, nSpire, CO, USA) using a ramped exercise protocol with AT determined by V slope and ventilatory equivalents.

All patients tested during this period were asked for permission to audit their outcomes. Patients were stratified by their CPET results into Fit ($AT \geq 11.0 \text{ mlO}_2\text{kg}^{-1}\text{min}^{-1}$), Unfit ($AT < 11.0 \text{ mlO}_2\text{kg}^{-1}\text{min}^{-1}$), or Unable to CPET if they failed to pedal the cycle or demonstrate an AT.

Patient demographics and "end of surgery" Colorectal Possum predicted percentage mortality were recorded, along with type of surgery (laparoscopic or open), use of critical care, length of hospital stay and thirty day, ninety day and two year mortality. All patients were cared for on our Specialist Colorectal surgical unit.

Analysis

Data were collated into Microsoft Excel and statistical analysis performed using MiniTab (Minitab® Statistical software v16, USA). Statistical significance was set at the 5% level. Data are presented as mean (standard deviation) where normally distributed and median (interquartile range) where not normally distributed. Between fitness group differences in total length of hospital stay were tested using one way analysis of variance (ANOVA) following log transformation of non-parametric data. Categorical data were compared between fitness groups using Chi Squared tests. Where there was evidence of overall differences between the three fitness groups, Bonferroni-corrected follow-up

multiple comparisons were used to identify between which pairs of groups there were statistically significant differences. A survival plot, stratified by fitness groups, was performed using the Kaplan-Meier method with survival distributions compared using the log rank test. Multi-variable binary logistic regression was also undertaken to examine whether fitness group had a significant influence on the probability of death by two years post-surgery, after allowing for other important variables (age, gender, and Duke's staging of malignancy).

Results

Between March 2009 and April 2010, 287 patients attended clinic and attempted CPET; 269 proceeded to surgery with follow up, five formally declined to be followed up, and 13 did not proceed to surgery. One hundred and seventy-nine patients were recruited to the trial of intraoperative fluid management.¹¹ All consenting survivors were followed to two years. A summary of the results, stratified by CPET fitness group, is presented in Table 1.

Thirteen patients did not proceed to bowel resection – five who would have been characterized as Fit, five Unfit and three as Unable to CPET. A summary of their results and reasons for not proceeding are presented as Internet Supplementary Table 1.

For the cohort proceeding to surgery median Length of Stay was 8 days (IQR 5.1-13.4) and there were 44 deaths at 2 years (16%).

Twenty six (9.7%) patients were categorized as Unable to CPET, 69 (25.7%) Unfit and 174 (64.7%) as Fit. Of the Unable patients 15 had a measured VO₂ peak but there was no AT demonstrable on V slope or ventilatory equivalents. The other 11 did not produce a test – seven due to arthritis, three due to frailty and one due to perineal pain.

The Unable patients were, on average, older than Unfit and Fit patients and typically had higher CR-POSSUM percentage predicted mortality although there was insufficient evidence to support a statistically significant difference between Fit and Unfit groups. The Unable group had higher mortality at each time point, and in particular at two years (11/26 (42%) vs. 14/69 (20%) vs. 19/174

CWL, GM CPC, KBH, JRS, SC RAS, March 2013

15

1
2
3 (11%), $p < 0.001$), although again there was insufficient evidence to support a statistically significant
4
5 difference between the Unfit and Unable groups.

6
7 Kaplan-Meier survival curves are presented in Figure 1; there was evidence of significant
8
9 differences in the survival distributions of the three groups ($p < 0.001$).

10
11 Binary logistic regression analysis was performed looking at main effects on probability of death at
12
13 two years after surgery. Results are presented in Table 2. Age at surgery was associated with an
14
15 odds ratio of 1.09 per year (95% CI 1.04-1.14) and Colorectal POSSUM Score at end of surgery
16
17 with an odds ratio of 1.17 (1.08-1.28). per percentage increase in predicted mortality. Male Gender
18
19 and Duke's C or D cancer staging (compared to non-malignant disease) were also significantly
20
21 associated with increased probability of mortality. After allowing for these variables, an inability to
22
23 perform CPET/demonstrate an AT, compared to being in the Fit group, was associated with an
24
25 increase in probability of mortality at two years (odds ratio of 3.98 (95% CI 1.04-11.73)).
26
27 Interestingly 20% of females who could not demonstrate an AT or perform CPET had died by 2
28
29 years compared to 73% of males in the same risk group.
30
31

32 33 **Discussion**

34
35
36 Previous studies have investigated the relationship between aerobic fitness as determined by CPET
37
38 and outcomes following major surgery. However, these have typically focused on patients who were
39
40 both able to complete CPET and generate an AT. To our knowledge, no studies have reported on
41
42 the outcomes of those who were unable to do so.

43
44 Our data demonstrate that such patients typically have a longer length of hospital stay and higher
45
46 30 day mortality rate and that a substantial proportion (42%) is no longer alive two years after
47
48 surgery. This has important implications for planning peri-operative care and for the process of
49
50 informed consent, including the decision whether or not to attempt curative resection.

51
52 Current consensus opinion is that patients should be explicitly provided with an estimation of the
53
54 risk of peri-operative death from the planned procedure and that this should be documented in the
55
56 patient notes and on consent forms.³ For many procedures, data is available on unit-specific 30 day
57
58
59
60

CWL, GM CPC, KBH, JRS, SC RAS, March 2013

15

1
2
3 mortality rates for all comers, however it is likely that the risk varies greatly for subgroups. More
4 individualized risk assessment facilitates better clinical decision-making.

5
6
7 Information about medium and long term mortality after surgery conveys a more realistic picture of
8 the burden of risk faced: mortality at 90 days is consistently double that at 30 days.^{8 12 13}

9
10 Scoring systems are potentially useful for individualized risk assessment. These indices are
11 generally derived using logistic regression in a large patient cohort, with subsequent validation of
12 the score in a separate patient cohort. Possum and CR-Possum are specific examples which have
13 been validated in a surgical population.^{14 15} These can assign patients into expected population risk
14 categories although they are not necessarily indicative of risk for the individual. Interestingly, the
15 Royal College of Surgeons report suggests that patients with a predicted operative mortality greater
16 than 10% should be automatically triaged to receive post-operative critical care admission.³ The
17 mean CR-Possum predicted mortalities for the Unable group in this trial exceeded 10%, whereas
18 predicted (and observed) mortalities for those patients who could generate an AT were much lower.
19 Despite the high predicted mortality, only 10 of the 26 patients (38%) in the Unable group had
20 planned post-operative critical care admission. This suggests that the attending clinicians did not
21 appreciate the association of an inability to successfully complete a cardiopulmonary exercise test
22 with a poor clinical outcome. A primary intention of this paper is to highlight the association.

23
24
25 A higher level of planned post-operative critical care might have mitigated against mortality risk for
26 this group.^{9 10} "Prehabilitation," incorporating a program of physical exercise, smoking cessation and
27 optimization of medical conditions may improve outcome for some patients.^{16 17}

28
29
30 In contrast to risk scoring indices, which are based on a tally of demographic data, comorbidities
31 and intraoperative events, CPET objectively quantifies the aerobic fitness of an individual. Thus it is
32 possible to identify patients who are at risk of perioperative functional heart failure, many of whom
33 will not have previously had a formal diagnosis of cardiovascular or respiratory disease.⁸ We
34 speculate that the inability to generate meaningful oxygen consumption measurements during the
35 test is a marker of particularly poor functional capacity— the test becomes one not of
36 cardiorespiratory reserve but of frailty.

37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

CWL, GM CPC, KBH, JRS, SC RAS, March 2013

15

1
2
3 The study has several limitations. The combined total of patients unable to pedal the cycle or no
4 demonstrating an AT is small and prevents further modeling. Analysis of a larger database is
5 advisable to confirm these findings. In particular there was not a statistically significant difference
6 between Unfit and Unable groups in length of stay or mortality when performing Bonferroni
7 corrected comparisons – although being in the Unfit group was associated with statistically
8 significant higher probability of mortality at two years when multivariate analysis was performed.

9
10
11
12
13
14
15 Fitness is not a dichotomous measure and the use of a cutoff in AT of 11 mlO₂ kg⁻¹min⁻¹ is
16 somewhat arbitrary. The Unable patients are likely to be at the lower end of a continuum of fitness
17 and as such at high risk of being unable to meet the metabolic demands of healing after surgery. As
18 such the inability to demonstrate a statistically significant difference between outcomes in the Unfit
19 and Unable patients might not be too surprising.

20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Surgical and anaesthetic techniques were not standardized, nor was access to planned critical care. Attending clinicians were not blinded to the results of any pre-operative investigations. In the regression model we have made statistical adjustment for differences in cancer staging between groups but in any small study such as this, co-morbidity imbalances and differences in care between groups may have an effect on comparative outcome.

We did not measure the incidence nor grade the severity of post-operative complications. It is likely that the longer length of hospital stay and increased mortality observed in the Unable group is linked to an increased adverse event rate but we cannot specifically comment on this. Similarly we did not use quality of life indicators to gauge the longer term recovery from surgery. This is important in evaluating whether surgical resection represents the best treatment option for an individual. When a patient has limited life expectancy (due to frailty or co-morbidities) then it may not be in their best interest that they endure a substantial portion of their remaining days trying to regain an acceptable quality of life following a surgical procedure. Humanity and patient dignity is an important aspect in patient care highlighted in the first (N)CEPOD report in 1987¹⁸ and echoed in the most recent.² We suggest that the benefits of operative over non-surgical options must be clear in order to justify proceeding to resection in this group of patients.

CWL, GM CPC, KBH, JRS, SC RAS, March 2013

15

In summary, our results imply that poor pre-operative functional capacity is associated with inferior post-operative outcomes. Failure to report the outcomes of those unable to perform the test may distort the clinical relevance of CPET studies.

The small proportion of patients who are unable to complete a cardiopulmonary exercise test have the longest lengths of hospital stay and higher early and medium term mortality and as a consequence their care represents a substantial burden on healthcare resources. Further studies on a larger cohort are required to confirm our findings and to identify peri-operative interventions to improve their outcomes.

Declaration of Interests The authors have no declaration of interests.

Funding This work was supported by an Anaesthesia/AABGI Departmental Research Grant (Round 1, 2008) through the National Institute of Academic Anaesthesia. Service support and administrative costs were met by the National Institute of Healthcare Research.

References

1. Pearse RM, Harrison DA, James P, Watson D, Hinds C, Rhodes A, Grounds M, Bennett ED. Identification and characterization of the high-risk surgical population in the United Kingdom. Crit Care 2006; 10:R81.
2. National Confidential Enquiry Into Patient Outcome and Death- Peri-operative Care: Knowing the Risk (2011). www.ncepod.org.uk/2011poc.htm, assessed on 17 March 2012.
3. The Royal College of Surgeons of England and Department of Health. The higher risk general surgical patient: Towards improved care for a forgotten group, 2011. <http://www.RoyalCollegeofSurgeonseng.ac.uk/publications/docs/higher-risk-surgical-patient/>, accessed on 17 March 2012.
4. Shoemaker WC, Appel PL, Kram HB, Role of oxygen debt in the development of organ failure sepsis, and death in high-risk surgical patients. Chest 1992; 102: 208-15.

- 1
2
3 5. Shoemaker WC, Appel PL, Kram HB, Waxman K, Lee TS. Prospective trial of supranormal
4 values of survivors as therapeutic goals in high-risk surgical patients. *Chest* 1998; 94: 1176-
5 86.
6
7
- 8
9 6. Older P, Smith R, Courtney P and Hone R. Preoperative evaluation of cardiac failure and
10 ischemia in elderly patients by cardiopulmonary exercise testing. *Chest* 1993; 104: 701-4.
11
12
- 13 7. Older P, Hall A, Hader R. Cardiopulmonary Exercise Testing as a Screening Test for
14 Perioperative Management of Major Surgery in the Elderly. *Chest* 1999; 116: 355-362.
15
16
- 17 8. Wilson RJT, Davies S, Yates D, Redman J, Stone M. Impaired functional capacity is
18 associated with all-cause mortality after major elective intra-abdominal surgery. *Br J*
19 *Anaesth* 105(3): 297-303 (2010)
20
21
- 22 9. Swart M, and Carlisle JB. Case-controlled study of critical care or surgical ward care after
23 elective open colorectal surgery. *Br J Surg* 2012; 99: 295-299.
24
25
- 26 10. Carlisle J, Swart M, Dawe EJC, Chadwick M. Factors associated with survival after resection
27 of colorectal adenocarcinoma in 314 patients. *Br J Anaesth* 2012; 108(3): 430-435.
28
29
- 30 11. Challand C, Struthers R, Sneyd JR, Erasmus PD, Mellor N, Hosie KB, Minto G. Randomized
31 controlled trial of intraoperative goal-directed fluid therapy in aerobically fit and unfit patients
32 having major colorectal surgery. *Br J Anaesth* 2011 doi: 10.1093/bja/aer273.
33
34
- 35 12. Chase P, Arena R, Guazzi M, Myers J, Peberdy M & Bensimhon D. Prognostic usefulness
36 of the functional aerobic reserve in patients with heart failure. *Am Heart J* 2010; 160: 922-7.
37
38
- 39 13. Visser BC, Keegan H, Martin M, Wren SM. Death after colectomy: It's later than we think.
40 *Arch Surg* 2009; 144(11): 1021-1027.
41
42
- 43 14. Copeland GP, Jone D, Walters M. POSSUM: a scoring system for surgical audit. *Br J Surg*
44 1991; 73(3): 355-60.
45
46
- 47 15. Tekkis PP, Prytherch DR, Kocher HM, Senapati A, Poloniecki JD, Stamatakis JD, Windsor
48 ACJ. Development of a dedicated risk-adjustment scoring system for colorectal surgery
49 (colorectal POSSUM). *Br J Surg* 2004; 91(9): 1174-1182.
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 16. Mayo NE, Feldman L, Scott S, Zavorsky G, Kim DJ, Charlebois P, Stein B, Carli F. Impact of
4 preoperative change in physical function on postoperative recovery: Argument supporting
5 prehabilitation for colorectal surgery. *Surgery* 2011; 150: 505-14.
6
7
8
9 17. Carli F, Charlebois P, Stein B, Feldman L, Zavorsky G, Kim DJ, Scott, Mayo NE.
10 Randomized clinical trial of prehabilitation in colorectal surgery. *Br J Surg* 2010; 97: 1187-
11 1197.
12
13
14
15 18. Buck N, Devlin HB, Lunn JN. The report of a confidential Enquiry into Perioperative Deaths.
16 *NCEPOD* 1987; 1(1): 1-175.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

CWL, GM CPC, KBH, JRS, SC RAS, March 2013

15

Groups	Fit	Unfit	Unable to CPET	Overall P- value
n	174	69	26	
Age ~	63.3 (19-89) ^a	70.2 (32-89) ^b	78.0 (58-90) ^c	<0.01
Gender Male / Female	101 / 73	34 / 35	11 / 15	0.20
Cancer Diagnosis	103 (59%) ^a	52 (75%) ^b	20 (77%) ^b	<0.01
Duke's A / B / C&D	20 / 37 / 46	11 / 23 / 18	3/10/7	0.65
Predicted CR Possum % Mortality *	3.9 (5.0) ^a	5.3 (6.8) ^a	11.5 (10.3) ^b	<0.01
Anaerobic threshold (mlO₂ kg-1min-1) *	14.4 (3.1)	9.5 (1.2)	Not Available	
VO₂ Peak (mlO₂ kg-1 min-1) *	22.3 (6.1) ^a	15.1 (3.7) ^b	11.6 (3.1) ^{#b}	<0.01
Laparoscopic procedures	61 (35%)	22 (32%)	6 (23%)	0.45
HDU/ITU				
Elective admissions	8 (5%) ^a	19 (28%) ^b	10 (38%) ^b	<0.01
Unplanned admissions	14 (8%)	9 (13%)	3 (12%)	0.45
Total length of hospital stay (days) §	7.1 (4.9 – 10.8) ^a	9.9 (5.5 – 15.0) ^b	14.0 (10.5-23.8) ^b	<0.01
Mortality 30 days	3 (2%) ^a	2 (3%) ^{a,b}	4 (15%) ^b	<0.01
90 days	5 (3%) ^a	5 (7%) ^{a,b}	5 (19%) ^b	<0.01
Two years	19 (11%) ^a	14 (20%) ^{a,b}	11 (42%) ^b	<0.01

Table 1 Summary of patient characteristics, surgery and outcomes

CPET CardioPulmonary Exercise Test ~Mean (Range) * Mean (standard deviation) § Median (IQR) # n=15.

^{a,b,c} Bonferroni-corrected follow-up multiple comparisons – groups with different symbol are significantly different after correction for multiple comparisons

CWL, GM CPC, KBH, JRS, SC RAS, March 2013

15

Table 2. Binary Logistic Regression Results Table for Probability of Death at Two Years Post-Surgery.

Predictor	Coef	SE Coef	Z	P	Odds Ratio	95% CI for Odds Ratio	
						Lower	Upper
Constant	-8.545	1.732	4.93	<0.01			
Age	0.086	0.024	3.65	<0.01	1.09	1.04	1.14
Predicted CR Possum % Mortality	0.158	0.043	3.66	<0.01	1.17	1.08	1.28
Fitness Group (vs. Fit)							
Unfit	0.545	0.443	1.23	0.22	1.72	0.72	4.10
Unable	1.381	0.562	2.46	0.01	3.98	1.32	11.97
Female Gender (vs. Male)	-0.837	0.413	-2.03	0.04	0.43	0.19	0.97
Malignancy Stage (vs. Non-Malignant)							
Dukes A	-0.541	0.913	-0.59	0.55	0.58	0.10	3.48
Dukes B	0.272	0.623	0.44	0.66	1.31	0.39	4.45
Dukes C&D	1.833	0.574	3.19	<0.01	6.25	2.03	19.26