

A low H I column density filament in NGC 2403: signature of interaction or accretion

W.J.G. de Blok^{1,2,3}, Katie M. Keating⁴, D.J. Pisano^{5,6}, F. Fraternali^{7,3}, F. Walter⁸, T. Oosterloo^{1,3}, E. Brinks⁹, F. Bigiel¹⁰, A. Leroy¹¹

¹ Netherlands Institute for Radio Astronomy (ASTRON), Postbus 2, 7990 AA Dwingeloo, the Netherlands, blok@astron.nl

² Astrophysics, Cosmology and Gravity Centre, Department of Astronomy, University of Cape Town, Private Bag X3, Rondebosch 7701, South Africa

³ Kapteyn Astronomical Institute, University of Groningen, PO Box 800, 9700 AV Groningen, the Netherlands

⁴ Rincon Research Corporation, Tucson, AZ 85711, USA

⁵ WVU Dept. of Physics & Astronomy, P.O. Box 6315, Morgantown, WV 26506, USA

⁶ Adjunct Assistant Astronomer at NRAO-Green Bank, P.O. Box 2, Green Bank, WV 24944, USA

⁷ Department of Physics and Astronomy, University of Bologna, via Bertini Pichat 6/2, I-40127 Bologna, Italy

⁸ Max-Planck Institut für Astronomie, Königstuhl 17, 69117 Heidelberg, Germany

⁹ Centre for Astrophysics Research, University of Hertfordshire, College Lane, Hatfield AL10 9AB, United Kingdom

¹⁰ Institut für theoretische Astrophysik, Zentrum für Astronomie der Universität Heidelberg, Albert-Ueberle Str. 2, 69120 Heidelberg, Germany

¹¹ National Radio Astronomy Observatory, 520 Edgemont Road, Charlottesville, VA 22903, USA

Received 1 January 1900 / Accepted 1 January 1900

ABSTRACT

Observed H I accretion around nearby galaxies can only account for a fraction of the gas supply needed to sustain the currently observed star formation rates. It is possible that additional accretion happens in the form of low column density cold flows, as predicted by numerical simulations of galaxy formation. To constrain the presence and properties of such flows, we present deep H I observations obtained with the NRAO Green Bank Telescope of an area measuring $4^\circ \times 4^\circ$ around NGC 2403. These observations, with a 5σ detection limit of $2.4 \cdot 10^{18} \text{ cm}^{-2}$ over a 20 km s^{-1} linewidth, reveal the presence of a low-column density, extended cloud outside the main H I disk, about $17'$ ($\sim 16 \text{ kpc}$ or $\sim 2 R_{25}$) to the NW of the center of the galaxy. The total H I mass of the cloud is $6.3 \cdot 10^6 M_\odot$, or 0.15 percent of the total H I mass of NGC 2403. The cloud is associated with an 8-kpc anomalous-velocity H I filament in the inner disk, previously observed in deep VLA observations by Fraternali et al. (2001, 2002). We discuss several scenarios for the origin of the cloud, and conclude that it is either accreting from the intergalactic medium, or is the result of a minor interaction with a neighbouring dwarf galaxy.

Key words. galaxies: halos – galaxies: ISM – galaxies: kinematics and dynamics – galaxies: individual: NGC 2403 – galaxies: structure

1. Introduction

Accretion of gas from the intergalactic medium (IGM) is often proposed as a mechanism for galaxies to obtain sufficient gas to maintain their star formation rate over cosmic time. There is however little direct observational evidence that can constrain the existence of the accretion and the magnitude of the accretion rates. In a recent review, Sancisi et al. (2008) make an inventory of possible signatures of gas accretion in and around nearby galaxies. They find an average visible H I accretion rate of $\sim 0.2 M_\odot \text{ year}^{-1}$, which is an order of magnitude too low to sustain the current star formation rates of the galaxies studied.

Searches for accreting gas are hampered by the difficulty of disentangling external, accreting gas from gas that has been expelled from the disk by processes of star formation (the so-called “galactic fountain”; Shapiro & Field 1976). In many cases, sophisticated three-dimensional modeling of the H I data cube is required to do this. Recent examples of this kind of analysis are presented in papers from the HALOGAS survey (Hydrogen Accretion in Local GALaxieS; Heald et al. 2011), a deep H I survey of 24 nearby galaxies, looking, amongst others, for signs of ac-

cretion in these galaxies (see Gentile et al. 2013; Kamphuis et al. 2013; Zschaechner et al. 2011, 2012).

These issues were also highlighted in earlier deep surveys of two nearby galaxies, NGC 891 and NGC 2403. Deep H I observations of the edge-on galaxy NGC 891 showed the presence of H I far above the main disk (Swaters et al. 1997; Oosterloo et al. 2007). Most of this H I was found to be rotating with the disk, but lagging slightly. This slower rotation of the extra-planar gas can be explained within the context of the galactic fountain model (Fraternali & Binney 2006). A smaller fraction of the extra-planar H I in NGC 891 was found to be at velocities inconsistent with circular rotation (Oosterloo et al. 2007). This gas is thought to be accreting. A number of the prominent H I filaments in NGC 891 are also difficult to explain with the galactic fountain model.

A second galaxy where prominent, lagging extra-planar H I is detected is NGC 2403 (Fraternali et al. 2001, 2002). This galaxy is observed at an intermediate inclination of 63° , which has the advantage that signs of in- or outflow of the gas can be detected directly. Modeling by Fraternali et al. (2001, 2002) and Frater-

nali & Binney (2008) shows that the kinematics of the extraplanar gas in NGC 2403 is consistent with inflow and this could therefore be a sign of accretion.

The analyses described above were done using data obtained with the Very Large Array (VLA) or the Westerbork Synthesis Radio Telescope (WSRT), i.e., radio interferometric telescopes. While these have excellent spatial resolution, a large observing effort is required to reach the low column density levels where one expects to see signs of accretion. For this reason, we have started a survey of a number of nearby galaxies (mainly taken from the THINGS survey; Walter et al. 2008) using the Green Bank Telescope (GBT), with the goal to obtain the deepest maps of these galaxies obtained so far, and quantify the presence of low-column density, and possibly accreting, features around these galaxies. A full description and first results of this survey will be presented elsewhere (Pisano 2014, Pisano et al., in prep.).

Here we report on the discovery, using GBT observations, of an extended, low-column density H I feature close to and possibly connected with NGC 2403. We describe the observations in Sect. 2. In Sect. 3, we argue that the feature is associated with an anomalous-velocity H I complex discovered in earlier high-resolution VLA observations (Fraternali et al. 2001, 2002). In Sect. 4 we describe various evolution scenarios for the feature. We summarize our results in Sect. 5.

2. Observations

NGC 2403 is a nearby ($D = 3.2$ Mpc; $1' = 0.93$ kpc) late-type spiral, and part of the M81 group. It has an absolute luminosity $M_B = -19.4$, an extended H I disk, and an inclination of 63° . It has a well-defined, symmetrical rotation curve, with a maximum rotation velocity of 135 km s^{-1} . Its heliocentric systemic velocity is 132.8 km s^{-1} . See Walter et al. (2008); de Blok et al. (2008) and Trachternach et al. (2008) for more detailed information regarding general properties of NGC 2403.

2.1. GBT observations

Observations of NGC 2403 were carried out with the 100m Robert C. Byrd Green Bank Telescope (GBT) of the NRAO¹ in 21 sessions between 29 May and 30 September 2009. We observed a $4^\circ \times 4^\circ$ area centered on NGC 2403. Strips of constant right ascension and declination, spaced by $3'$, were observed to form a ‘basket weave’ pattern over the region. The spacing and integration times ensure Nyquist-sampling. The total integration time for the entire map was approximately 127 hours, corresponding to an integration time per beam of ~ 11 min. The GBT spectrometer was used with a bandwidth of 12.5 MHz. This corresponds to a velocity range from $-885 \text{ km s}^{-1} < v_{\text{LSR}} < 1750 \text{ km s}^{-1}$. The typical system temperature for each channel of the dual-polarization receiver was ~ 20 K.

A reference spectrum for each of the sessions was made by observing an emission-free region about 4.75° away from the galaxy. The reference spectrum was then used to perform a (signal–reference)/reference calibration of each pixel. The calibrated spectra were scaled by the system temperature, corrected for atmospheric opacity and GBT efficiency. We adopted the GBT efficiency equation (1) from Langston & Turner (2007)

¹ The National Radio Astronomy Observatory (NRAO) is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

with a zenith atmospheric opacity $\tau_0 = 0.009$. Data reduction was done using custom routines developed using GBTIDL².

The frequency range observed was relatively free of radio-frequency interference (RFI), with $\sim 5\%$ of all spectra adversely affected. Spectra that showed high noise values across many channels were flagged and removed. After amplitude calibration and gridding, a first-order polynomial was fit to line-free regions of the spectra and subtracted from the gridded spectra.

Data from all observing sessions were converted to heliocentric velocities and combined into a single data cube. In order to match the typical linewidths of neutral hydrogen features observed around galaxies, spectra were smoothed to a velocity resolution of 5.2 km s^{-1} . The effective angular resolution is $8.7'$, and we use a pixel size of $1.75'$. The calibration from K to Jy was derived by observing 3C286 in the same way as the H I maps. The final scale factor from K to Jy is 0.43 ± 0.03 . Due to remaining low-level RFI and residual continuum emission, the noise level varies by $\sim 20\%$ throughout the cube. The average RMS noise in the final data cube is $6.0 \text{ mJy beam}^{-1}$ or 13.6 mK per 5.2 km s^{-1} channel. This translates to a 1σ column density sensitivity of $1.3 \cdot 10^{17} \text{ cm}^{-2}$ per channel, or a 5σ sensitivity of $2.4 \cdot 10^{18} \text{ cm}^{-2}$ for a 20 km s^{-1} full width at half maximum H I line.

2.2. Deep VLA observations

In this paper we compare the GBT data with deep VLA observations obtained by Fraternali et al. (2001, 2002). They observed NGC 2403 for 40 hours with the VLA in its CS configuration. Here we use the $30''$ data presented in Fraternali et al. (2002). The data cube was created using a robustness parameter of 0 and a taper of $27''$. The channel spacing is 5.15 km s^{-1} . The data were Hanning-smoothed, leading to an effective velocity resolution of 10.3 km s^{-1} . The synthesized beam size of this cube is $29.7'' \times 29.3''$ and the column density sensitivity limit for a 5σ detection over one velocity resolution element is $2.0 \cdot 10^{19} \text{ cm}^{-2}$. See Fraternali et al. (2002) for more details.

In addition, we also smoothed the $30''$ cube spatially to a resolution of $60''$ and $90''$, as well as to the GBT resolution of $8.7'$ (using a simple Gaussian smoothing function). We use these cubes later to determine how much of the H I detected with the GBT was also detected by the deep VLA observations.

3. Analysis

3.1. GBT channel maps and moment maps

To derive the total H I mass and the global velocity profile, we clipped the GBT data cube at 2.5σ , also removing by hand a small number of spurious noise peaks (caused by low-level residual RFI). The rotation and systemic velocity of NGC 2403 mean that for radial velocities below $v_{\text{hel}} \sim 10 \text{ km s}^{-1}$ galaxy emission overlaps with Galactic emission. Based on symmetry considerations the velocity range where we cannot recover the NGC 2403 H I distribution is however small.

Figure 1 shows the global H I profile. We find a total flux of $1677.1 \text{ Jy km s}^{-1}$. This does not include any emission below $v_{\text{hel}} = 10 \text{ km s}^{-1}$, but it is clear from Fig. 1 that this will be only a small fraction of the total flux. Assuming a distance of 3.2 Mpc, we find an H I mass of $4.03 \cdot 10^9 M_\odot$. This is higher than was found in previous observations. Single-dish data by Rots (1980) yield a total flux of $1448.1 \text{ Jy km s}^{-1}$ or an H I mass of $3.50 \cdot 10^9$

² <http://gbtidl.nrao.edu>




Fig. 1. Global H I profile of NGC 2403 derived from the GBT data. The dotted line indicates $v_{\text{hel}} = 10 \text{ km s}^{-1}$. Below this velocity, Galactic emission affects the data and emission from NGC 2403 could not be unambiguously identified.

M_{\odot} . Interferometric data by Begeman (1987) gave an H I mass of $3.0 \cdot 10^9 M_{\odot}$, observations by Sicking (1997) $3.13 \cdot 10^9 M_{\odot}$, and deep observations by Fraternali et al. (2002) $3.28 \cdot 10^9 M_{\odot}$ (all values corrected to $D = 3.2 \text{ Mpc}$).

Uncertainties in interferometrically measured H I masses are usually dominated by calibration uncertainties, bandpass corrections, selection of masks, etc., rather than by the formal statistical uncertainties. They are therefore difficult to quantify. An empirically motivated estimate of the uncertainties is $\sim 10\%$. This value is higher than the formal statistical uncertainty, which tends to be a few percent at most for bright sources such as considered here (cf. the values quoted in Begeman 1987 and Sicking 1997). Comparing the above measurements with the GBT observations, we see that the latter detect 20% more flux, i.e., a significantly higher value.

Part of this additional flux will be due to the larger area observed by the GBT compared to the single interferometer pointings. However, we also detect more flux than found in the Rots (1980) single-dish data. Rots (1980) surveyed a similar area as our observations, so the difference cannot be due to a more limited field of view. A comparison between the two single-dish global profiles, shows that our observations detect more flux over the entire velocity range of the galaxy. Although this could be a calibration issue, it is likely that our new GBT observations, which are approximately 50–100 times more sensitive in terms of column density, detect more low column density H I. For example, the low-column density kinematically anomalous H I detected by Fraternali et al. (2001, 2002) has a total H I mass of $\sim 3 \cdot 10^8 M_{\odot}$, which could explain a large part of the difference in the derived single-dish masses.

Selected channel maps from the GBT data cube are shown in Fig. 2. The emission from the main disk of NGC 2403 is symmetric and regular, though the velocity field (see below) indicates that the outer disk is slightly warped. In the channel maps between $v = 150.5$ and 104.1 km s^{-1} we see a faint extension to the NW of the main emission component, indicated by the arrow

in the $v = 140.2 \text{ km s}^{-1}$ panel. This “cloud” seems to connect up with the main disk at lower velocities, but looks separate from it towards higher velocities. All velocities mentioned here and below are heliocentric velocities.

Using the clipped data cube described above we create moment maps showing the integrated emission (zeroth moment) and the intensity-weighted average velocity (first moment). These are shown in the left-hand panels in Fig. 3. In this Figure, we also show for comparison (in the right-hand panels) the integrated H I map and velocity field derived from the 30'' VLA data set from Fraternali et al. (2002). The H I distribution in the GBT data, even when taking the different beam sizes into account, extends further out than in the VLA data. Part of this is due to the size of the primary beam of the VLA. Figure 4 shows the 50 percent and 10 percent sensitivity VLA primary beam sizes overlaid on the VLA integrated H I map. The NGC 2403 emission as observed by the VLA extends out to the edge of the 50 percent sensitivity primary beam. Beyond this radius the primary beam sensitivity drops rapidly, hampering the detection any emission that may be present there. This emission is however detected by the GBT observations which do not suffer from primary beam effects.

The velocity field derived from the GBT data cube looks regular, and shows signs of a slight warp in the outer parts, as indicated by a small variation of the position angle of the kinematical major axis.

In addition to the full data cube used to create the moment maps discussed above, we also created a cube containing only the emission from the cloud. This was done by applying an intensity cut to remove the very bright emission from the main disk. We created an integrated H I map of the cloud, which is shown in the top panels of Fig. 3 overlaid on the GBT and VLA maps. The cloud clearly overlaps with the main disk as seen at the GBT resolution. The high contrast between cloud and disk emission, combined with the relatively large size of the beam, means we can only identify cloud emission where (in the channel maps) it does not overlap with the main disk (see, e.g., Fig. 2 at $v = 124.8 \text{ km s}^{-1}$ where the cloud emission merges in projection with the main disk emission). It is therefore likely that the cloud extends further toward the SE than Fig. 2 suggests, and that the SE edge is artificial. Due to the confusion with Galactic emission it is not possible to establish whether other low column density features are present in the channels with $v < 10 \text{ km s}^{-1}$. If present, they would however not be related to the cloud because of the different spatial location of the main disk H I at these velocities.

The total H I mass of the cloud is $6.3 \cdot 10^6 M_{\odot}$, or about 0.15 percent of the total H I mass of NGC 2403. The radial distance of the highest column density part of the cloud to the center of NGC 2403 is $\sim 17'$ (corresponding to $\sim 16 \text{ kpc}$ or ~ 2 times R_{25}). The H I mass is a lower limit because, as noted above, we can only identify cloud emission when it is not projected against the main disk.

3.2. Comparison with deep VLA observations

We compare the GBT data with the deep, single-pointing VLA H I study of NGC 2403 by Fraternali et al. (2001, 2002) as described in Sect. 2.2.

One of the main results to come out of that study is the detection of an extended, kinematically anomalous H I component in NGC 2403, which Fraternali et al. (2001, 2002) interpret as a thicker disk which rotates slower than the thin, main, cold H I disk. The velocity field of the thick component shows evidence for inflow motion towards the center. In addition, Fraternali et al.




Fig. 2. Selected channel maps of the GBT data cube. Contours levels are -2σ (gray dashed contours) and $(2, 4, 8, 16, 32, 64, 128, 256, 512, 1024) \cdot \sigma$ (full contours) where $\sigma = 6 \text{ mJy beam}^{-1}$. This corresponds within a single channel to a column density of $1.25 \cdot 10^{17} \text{ cm}^{-2}$. Heliocentric velocities of the channels in km s^{-1} are shown in the top-right of each channel. From 150.5 to 104.1 km s^{-1} every channel is shown. Outside this range only every third channel is shown. The GBT beam is shown in the bottom-left corner of the top-left panel. Channels in the bottom-row are marginally affected by Galactic emission. In the 140.2 km s^{-1} panel the arrow indicates cloud emission.

(2001, 2002) report the presence of gas at “forbidden” velocities, that is, having velocities that make it impossible for it to be part of the regular rotation of the disk.

In addition, NGC 2403 was found to contain a number of kinematically anomalous H I complexes that are not at forbidden velocities, but nevertheless deviate from regular, circular rotation. Fraternali et al. (2001, 2002) interpret the anomalous gas as a sign of accretion.

One of the most prominent of these anomalous H I complexes is a filament with an H I mass of $\sim 1 \cdot 10^7 M_{\odot}$, located in the inner disk, with a velocity that differs 60–100 km s^{-1} from

the local rotation, and which is referred to as the “8-kpc filament” by Fraternali et al. (2002).

Fraternali et al. (2002) fit Gaussians to the peaks of the H I profiles and subtract these from the data to isolate emission with anomalous velocities. We also use this method to separate out the emission from the 8-kpc filament in the VLA data cube. We show an integrated H I map of this filament superimposed on that of the main disk in the top panels of Fig. 3 (see also Figs. 7 and 8 in Sancisi et al. 2008). These figures immediately show that the 8-kpc filament points towards the cloud seen by the GBT.

We make a more detailed comparison in Fig. 5, where we show again the GBT channel maps from Fig. 2, but this time su-




Fig. 3. H I moment maps of NGC 2403. **Top-left:** Integrated H I map derived from the GBT data. Thick dark-gray contours show the H I distribution of the entire galaxy. Contour levels are $5 \cdot 10^{17}$, $5 \cdot 10^{18}$, $5 \cdot 10^{19}$, and $5 \cdot 10^{20} \text{ cm}^{-2}$. The gray-scale runs from $5 \cdot 10^{17}$ (white) to $1.5 \cdot 10^{21} \text{ cm}^{-2}$ (black). Thin, black contours show the H I distribution of the cloud. Contour levels are $(6.25, 12.5, 25, 62.5) \cdot 10^{17} \text{ cm}^{-2}$. White contours indicate the location of the 8-kpc filament. The GBT beam is shown in the bottom-left. **Top-right:** Integrated H I map derived from the 30'' VLA data from Fraternali et al. (2002). Grayscale runs from $2 \cdot 10^{19}$ (white) to $2 \cdot 10^{21} \text{ cm}^{-2}$ (black). Contours are as in top-left panel. The VLA beam is indicated in the bottom-right corner. **Bottom-left:** Intensity-weighted mean velocity field derived from the GBT data. The thick black contour indicates $v = 130 \text{ km s}^{-1}$. White contours decrease with respect to this value in steps of 20 km s^{-1} . Black contours increase in steps of 20 km s^{-1} . **Bottom-right:** Intensity-weighted mean velocity field derived from the 30'' VLA data. Contours are as in bottom-left panel.

perimposed on the corresponding deep VLA data channel maps from Fraternali et al. (2002). Because of the large area shown, the VLA data are not corrected for primary beam effects.

In the VLA data in Fig. 5, note the “finger” of emission pointing to the NW away from the main emission between 150.5 and 109.3 km s^{-1} . This is the 8-kpc filament. There is a clear correlation between the appearance of this “finger” in the VLA data and the cloud emission in the GBT data, which strongly hints at the cloud and filament being associated, and likely forming one single complex. As mentioned earlier, the spatial separation between the filament and cloud features is probably artificial, due to our inability to detect faint cloud emission projected against the bright main disk. We show two channel maps in more detail in the bottom panels of Fig. 6, where we show the GBT emission superimposed on the VLA data smoothed to $60''$. The higher sensitivity that we achieve due to the spatial smoothing allows us to trace the “finger” further out, to the point where it touches the cloud emission detected by the GBT. Apart from the

spatial continuity, there is also a clear velocity continuity which we illustrate using the position-velocity (pV) diagram shown in the top panel of Fig. 6. The pV diagram was created by taking a slice through the VLA and GBT cubes, going through both the center of the cloud and the filament, as indicated by the line in Fig. 4. Due to the large difference in resolution between the two datasets, we chose different thicknesses for the slices. For the GBT data we extracted a slice with a thickness of one GBT beam width. To maximize the signal in the VLA data, we extracted a slice with a thickness of $100''$, corresponding to the average width of the filament. The comparison between the slices in Fig. 6 clearly shows that the filament and the cloud occur at the same velocities (and *only* those velocities). The clear spatial and velocity association strongly suggests that the cloud and the filament form one single complex.

The link with the 8-kpc filament definitively rules out the possibility that the cloud is a Galactic foreground object not related to NGC 2403. Such objects are known to exist in this part




Fig. 4. A view of the cloud-filament complex. Grayscale shows the integrated H I distribution derived from the 30'' VLA data and run from $2 \cdot 10^{19}$ (white) to $2 \cdot 10^{21}$ cm^{-2} (black). Thin, black contours show the H I distribution of the cloud derived from the GBT data. Contour levels are $(6.25, 12.5, 25, 62.5) \cdot 10^{17}$ cm^{-2} . White contours show the 8-kpc filament, derived from the 30'' VLA data. The contour level is $2 \cdot 10^{19}$ cm^{-2} . The highest column density found in the filament is $6.5 \cdot 10^{19}$ cm^{-2} . The GBT and 30'' VLA beams are indicated in the bottom-left corner. The thick gray line shows the location of the position-velocity slice shown in Fig. 6. The long-dashed circle indicates the 50 percent sensitivity level of the VLA primary beam. The short-dashed circle shows the 10 percent level.

of the sky (Chynoweth et al. 2009), but they are typically found at velocities of ~ -150 km s^{-1} and are therefore likely to be part of the Milky Way high-velocity cloud population.

We investigated whether the filament could be traced even further out towards the cloud by convolving the 30'' VLA data to a beam size of 90''. We created a moment map of all emission between 119.6 and 150.4 km s^{-1} (cf. Fig. 6). No clipping or masking was applied. The resulting map is shown in Fig. 7. Compared to the 30'' data, the emission extends twice as far and the tip of the filament overlaps with the SE part of the cloud. Figure 7 shows hints that at the $\sim 1.5\sigma$ to $\sim 2\sigma$ level the filament extends even further towards the center of the cloud. However, at this level, the data are also affected by deconvolution artefacts (present as faint features in the bottom-left and top-right of the Figure) which partly overlap with the possible faint extension. In the absence of deeper high-resolution data it is therefore difficult to say more about this low-level extension. Nevertheless, in Fig. 7, the main filament feature is present at high significance levels and just overlaps with the (likely artificial) SE of the cloud, further suggesting that we are dealing with a single cloud/filament complex.

But is the cloud as seen by the GBT indeed a new feature which has not been seen before, or is it simply the already known 8-kpc filament as observed with the VLA, but smoothed to a lower resolution? To answer this, we use the VLA data cube smoothed to the GBT resolution, as described in Sect. 2.2. We compare corresponding channel maps and a position-velocity slice of this smoothed VLA data set and the GBT data in Figs. 5 and 6, respectively. From these Figures it is clear that the cloud

emission was not detected in the single-pointing VLA observations, and that the GBT observations are showing additional H I. An explanation for this was already mentioned: the single-pointing VLA observation has a limited field of view, and as can be seen in Fig. 4, the cloud is located in the region of the primary beam where the VLA sensitivity has dropped to between 50 and 10 percent of the central value.

Can we say anything about the column densities of the cloud compared with that of the filament? The H I mass of the cloud is $6.3 \cdot 10^6 M_{\odot}$. If we take the dimensions of the cloud to be $16' \times 12'$ (i.e., the angular size of the cloud as shown in Fig. 4 corrected for the GBT beam size), then the average H I column density of the cloud is $\sim 6 \cdot 10^{18}$ cm^{-2} assuming a uniform distribution. This would not be detected by the deep VLA observations.

Due to the GBT beam size we can, however, not distinguish between smooth emission which fills the beam, or a more clumpy, higher column density distribution with a smaller beam filling factor. To take the other extreme, if we assume that the H I in the cloud has a size and distribution identical to that of the 8-kpc filament, then the inferred average column density increases to $\sim 6 \cdot 10^{19}$ cm^{-2} . The column densities in the actual observed 8-kpc filament are in the range ~ 3 to $\sim 6 \cdot 10^{19}$ cm^{-2} , so consistent with this value.

It would however still be difficult to detect this more compact emission in the existing 30'' VLA data. Putting the observed 8-kpc filament at the position of the cloud means applying a primary beam loss of sensitivity of a factor of ~ 0.4 . The resulting apparent column density would thus be ~ 1.5 to $2.5 \cdot 10^{19}$ cm^{-2} . Assuming that the emission has a velocity width of ~ 50 km s^{-1} ,




Fig. 5. Selected channel maps of the GBT data cube superimposed on corresponding channel maps from the deep VLA data. The GBT data are shown in black contours. Black contours levels are $(2.5, 5 [thick\ contour], 10, 50) \cdot 10^{17} \text{ cm}^{-2}$. Gray, dashed contours denote $-2.5 \cdot 10^{17} \text{ cm}^{-2}$ in the GBT data. The gray scale shows the H I emission in the deep $30''$ cube from Fraternali et al. (2002). The thin, red contour surrounding this emission shows the $5 \cdot 10^{18} \text{ cm}^{-2}$ in these VLA data. The thick, red contours show the $5 \cdot 10^{17} \text{ cm}^{-2}$ level in the VLA data when smoothed to the GBT resolution of $8.7''$. This is the same column density as shown by the thick black contour. None of the VLA contours or grayscales are corrected for primary beam effects. The panel in the bottom-right is affected by Galactic emission. The GBT beam is shown in the bottom-left of the top-left panel. The two small features to the north of the main emission, as indicated by the thick red contour at $v = 258.7$ and 243.3 km s^{-1} are data artefacts due to the severe smoothing used.

implies that in the $30''$ VLA data, an 8-kpc filament at the position of the cloud would show at the $\sim 1\sigma$ level in each channel (cf. the sensitivities listed in Sec. 2.2).

Figure 7 shows that the $90''$ VLA data has detected more H I associated with the filament than the $30''$ data. The “extra” emission detected in the former data set (compared to the $30''$ maps) has a corresponding H I mass between $\sim 3 \cdot 10^5 M_{\odot}$ and $\sim 8 \cdot 10^5 M_{\odot}$, depending on how exactly the extent of the $30''$ filament is defined. This is only between 5 and 13 percent of the H I cloud mass detected by the GBT, so it is clear that even

with the extra spatial smoothing applied here, the VLA does not detect all of the cloud emission.

In an attempt to further constrain the nature of the cloud/filament complex, we observed the cloud with the WSRT in Aug and Dec 2013 for a total of 21 hours. The telescope was pointed at the center of the GBT cloud. The data were reduced using standard methods, and an image cube was produced using a robustness parameter of 0 and a taper of $30''$. The resulting beam size was $42'' \times 47''$, with a noise level of $0.9 \text{ mJy beam}^{-1}$.




Fig. 6. Comparison of GBT and 30'' VLA data. **Top panel:** pV slice through centers of cloud and 8-kpc filament. Position and orientation of the slice is indicated in Fig. 4. The thickness of the slice through the VLA data is 100''; the thickness of the slice through the GBT data is one GBT beam. The grayscale shows the VLA data with $(2.5, 5) \cdot 10^{18} \text{ cm}^{-2}$ contours overplotted in dark-gray. The black contours show the GBT data at $(5, 10, 25, 50, 100) \cdot \sigma$, where σ corresponds to $3 \cdot 10^{17} \text{ cm}^{-2}$. The cloud is detected at the $> 10\sigma$ level. The red contour shows the VLA data smoothed to the GBT resolution at the $5 \cdot 10^{17} \text{ cm}^{-2}$ level. It is clear the emission detected by the GBT is more extended. The VLA contours and grayscales are not corrected for primary beam effects. The hatched area at the top indicates velocities affected by Galactic emission. The cloud and 8-kpc filament are indicated by arrows. The two horizontal lines indicate the velocities of the channel maps shown in the bottom panels. **Bottom panels:** Channel maps at $v = 135.1 \text{ km s}^{-1}$ (left) and $v = 124.8 \text{ km s}^{-1}$ (right). In both panels the grayscale shows the VLA data at 60''. Thin dark-gray contours show VLA column density levels at $(1, 5) \cdot 10^{18} \text{ cm}^{-2}$. Thin white contours show VLA column density levels at $(25, 50, 250) \cdot 10^{18} \text{ cm}^{-2}$. The lowest contour of $1 \cdot 10^{18} \text{ cm}^{-2}$ corresponds to $\sim 2.5\sigma$. Features in the extreme top-right and bottom-left of the channel maps are deconvolution artefacts. The thick black contours show GBT column density levels and are as in Fig. 5. The GBT beam and the 60'' VLA beam are shown in the bottom-left of the left channel map.

This corresponds to a 5σ , 10 km s^{-1} column density sensitivity of $2.5 \cdot 10^{19} \text{ cm}^{-2}$.

The WSRT data confirm all features seen in the deep VLA data, but do not detect significant emission at the position of the cloud. The similar limits derived from both the WSRT and the VLA data thus indicate that the cloud H I must have column den-

sities below $\sim 10^{19} \text{ cm}^{-2}$ (when observed at 30'' and integrated over the velocity width of the cloud) for it not be detected in the deep VLA, or the more shallow but targeted WSRT observations. The combination of this column density limit, and the total H I mass of the cloud, implies that the true spatial distribution of the H I must be fairly uniform, on scales comparable to the extent




Fig. 7. Integrated H I map of the emission between 119.6 and 150.4 km s^{-1} derived from the deep VLA data at 90'' resolution (grayscale). Light-gray contours show $(5, 10, 20, 50, 100, 200) \cdot 10^{18} \text{ cm}^{-2}$ column density levels. The noise in the map corresponds to $2 \cdot 10^{18} \text{ cm}^{-2}$ and the lowest VLA contour corresponds to $\sim 2.5\sigma$. Thin black contours show the GBT cloud, thick white contours the 8-kpc filament at 30'' resolution. Contours as in Fig. 4. The 90'' VLA beam is indicated in the bottom-right corner, the GBT beam in the bottom-left corner.

of the GBT-detected emission, as more clumpy H I distributions with higher local column densities would have been detected.

4. Interpretation

4.1. Comparison with optical data

We investigate whether any signatures or counterparts of the cloud are visible at other wavelengths. Inspection of GALEX images (Gil de Paz et al. 2007) shows no sign of an overdensity of near-ultraviolet (NUV) or far-ultraviolet (FUV) sources in the area of the cloud, so there is no indication of a concentration of recent star formation at that location.

A study by Barker et al. (2012) looks at the resolved stellar population in and around NGC 2403, using wide-field observations with Suprime-Cam on the Subaru telescope. They used colour-magnitude diagrams to identify different stellar populations and found that the stellar distribution of NGC 2403 extends to at least ~ 40 kpc, at a surface brightness level of $\mu_V \sim 32$ mag arcsec $^{-2}$. The field of view of these observations thus comfortably encompasses the location of the cloud, which is at a radius of $\sim 17'$ or 16 kpc.

Barker et al. (2012) present the spatial distribution of the stellar populations in their Fig. 10. Their plot of the distribution of RGB and AGB stars shows diffuse overdensities of stars to the NW and SE of the main disk, with the NW overdensity being most prominent. In Fig. 8 we show the distribution of RGB stars in NGC 2403 from Barker et al. (2012), and overlay the cloud and the filament. The center of the cloud is offset $\sim 5'$ to the NE from the approximate center of the stellar overdensity, but still has a substantial overlap.

The optical overdensity shows little internal structure and is likely part of the faint outer stellar disk, possibly tracing the warp as seen in H I (cf. Fig. 3). As noted above, according to Barker et al. (2012), stellar emission can be traced out to well beyond the




Fig. 8. The cloud and filament (contour levels as in Fig. 3) overlaid on the distribution of RGB stars from Barker et al. (2012) (grayscale). The ellipses are spaced 10 kpc in deprojected radius. The absence of stars in the center is due to crowding. The GBT beam is shown in the lower left of the panel. The distribution of RGB stars is copied from Fig. 10 of Barker et al. (2012).

cloud. The (radially averaged) surface brightness at the radius of the cloud is $\mu_V \sim 29$ mag arcsec $^{-2}$.

The location of the cloud near the optical overdensity may thus be a chance superposition and the cloud/filament complex is possibly the equivalent of a large high velocity cloud (HVC). The H I mass of the complex is similar to that of the Complex C HVC in our Galaxy. This raises the question of whether the cloud/filament complex could be caused by galactic fountain processes, i.e., could it be gas blown out of the main disk due to star formation and forming a HVC? As discussed above, Fraternali et al. (2001, 2002) show the presence of a thick, lagging H I disk in NGC 2403. Fraternali & Binney (2006, 2008) model this extra-planar gas component using particles that are blown out of the disk on ballistic orbits. Fraternali & Binney (2008) conclude that the general distribution of the gas can indeed be modelled in this way, but that smaller substructures and large filaments (like the 8-kpc filament) are not easily produced in the fountain model. With the addition of the GBT data, the 8-kpc filament is now twice as massive and extends much further out than originally assumed, making it less likely to be purely the result of the galactic fountain process.

4.2. Comparison with NGC 891: a fly-by scenario

We already noted that another galaxy in which extraplanar gas has been observed and extensively studied is NGC 891. A paper by Mapelli et al. (2008), which focusses on the origin of the lopsided disk in NGC 891, offers an interesting alternative explanation for the presence of some of the extra-planar gas observed in that galaxy.

Mapelli et al. (2008) mention that the main extra-planar filament in NGC 891 is pointing at nearby LSB galaxy UGC 1807 (as first pointed out in Oosterloo et al. 2007; cf. their Fig. 8), and using simulations, show that a fly-by interaction with an

intruder dwarf galaxy can create this filament. Note, incidentally, that these types of fly-bys are also a common phenomenon in cosmological cold dark matter simulations (Sinha & Holley-Bockelmann 2012).

For the specific case of NGC 891, Mapelli et al. (2008) find the best match with an interaction that happened 300 Myr ago, at a fly-by velocity of 260 km s^{-1} . They show that the filament must consist mostly of gas stripped off the UGC 1807 intruder.

The Mapelli et al. (2008) simulations indicate the presence of gas between the two galaxies. Deep GBT H I observations of the area around NGC 891 do, however, not detect any connection between NGC 891 and UGC 1807 (D.J. Pisano, priv. com.) implying that if the fly-by hypothesis holds, then this gas must be of very low column density and/or ionized, and only close to the NGC 891 disk dense enough to be detected.

Could the cloud-complex in NGC 2403 be caused by a similar fly-by of a neighboring galaxy? If we extend the major axis of the VLA 8-kpc filament³ (using a major axis position angle of -17° ; cf. Fig. 4) we find that it passes close to the dwarf spheroidal galaxy DDO 44 (at about 1.5° or $\sim 85 \text{ kpc}$ from NGC 2403) and also the late-type dwarf galaxy NGC 2366 (at $\sim 3.5^\circ$ or $\sim 190 \text{ kpc}$). The velocities of the H I in NGC 2366 completely overlap with those of the H I in NGC 2403. In NGC 2403, H I is found from ~ -5 to $\sim 270 \text{ km s}^{-1}$. In NGC 2366, this ranges from ~ 40 to $\sim 170 \text{ km s}^{-1}$ (Walter et al. 2008). DDO 44 has a single radial velocity determination that puts it at 213 km s^{-1} (Karachentsev et al. 2011), which is inside the range of velocities found in NGC 2403. See Fig. 9 for an overview.

Arbitrarily assuming the same fly-by velocity of 260 km s^{-1} as given in Mapelli et al. (2008) (and assuming that this velocity is in the plane of the sky, ignoring any radial velocities), we find the timescale for an interaction with DDO 44 to be $\sim 300 \text{ Myr}$. For NGC 2366 we find $\sim 750 \text{ Myr}$. The DDO 44 timescale is comparable to the one Mapelli et al. (2008) found for NGC 891.

Given the orbital time of NGC 2403, which is $\sim 700 \text{ Myr}$ at the approximate position of the cloud, it is not clear which part of the stellar disk would have been affected most. However, the effects of the interaction on anything other than the local gas distribution must have been minor. In a study of the star formation histories at various locations in the NGC 2403 disk, Williams et al. (2013) find no evidence for any major temporal variations in the star formation rate. The locations studied were all on the receding side of the galaxy (i.e., the side of the galaxy *opposite* to that of the cloud). Given the regularity and the symmetry of the stellar disk of NGC 2403, there is, however, no reason to assume that the star formation history of the approaching side of the galaxy is dramatically different. Williams et al. (2013) suggest that as NGC 2403 is so undisturbed, it is unlikely to have had major interactions in the past.

Do any of the two potential intruder galaxies show any signs of interactions or disturbance? NGC 2366 is a gas-rich dwarf-galaxy which is known to have a disturbed H I velocity field and an outer gas component that does not follow regular rotation (de Blok et al. 2008; Oh et al. 2008). It is undergoing a star-burst and shows evidence for major gas outflows (van Eymeren et al. 2009). McQuinn et al. (2010), from modeling of the stellar population, find that the star formation burst started and peaked $\sim 450 \text{ Myr}$ ago.

The second, closer, galaxy DDO 44 is a dwarf-spheroidal galaxy. In a search for H α clumps in the M81 group, Karachentsev et al. (2011) detected an H II region in this dSph galaxy, containing a number of blue, late-type B-stars. Karachentsev et

al. (2011) note that, despite the absence of early-type B- or O-stars, the FUV luminosity of $\sim 1 \cdot 10^{36} \text{ erg s}^{-1}$ of the detected stars is sufficient to ionise the H II region. These stars are also seen in GALEX data. The implied star formation rate (SFR) is $\sim 10^{-5} M_\odot \text{ yr}^{-1}$. Despite this small SFR value, it is surprising to find recent star formation in such a gas-poor system. No H I has so far been detected in DDO 44, and we also find no evidence in our deep GBT cube (which also covers the position of DDO 44). Assuming, like Karachentsev et al. (2011), that the velocity width of DDO 44 is $W \leq 30 \text{ km s}^{-1}$, puts the 5σ H I mass upper limit derived from our GBT data at $M_{\text{HI}} \sim 8 \cdot 10^4 M_\odot$.

In their paper, Karachentsev et al. (2011) speculate that the star formation is caused by accretion of gas from the intergalactic medium. If the fly-by scenario is valid, this might therefore be star formation due to gas dragged out of NGC 2403's immediate environment. The life-time of a late-type B-star (a few hundred Myr) is consistent with the timescale for the fly-by derived above.

We did search the GBT data cube for evidence of an H I connection between NGC 2403 and DDO 44, but did not find any emission beyond what was already detected in the filament, though one may expect that gas to be of such low column density that it is ionised. In fact, we do not see any evidence throughout the entire GBT cube for emission not associated with NGC 2403 or our Milky Way at the 3σ level ($3.9 \cdot 10^{17} \text{ cm}^{-1}$ per 5.2 km s^{-1} channel). Nevertheless, the simulations presented in Mapelli et al. (2008) suggest that in the fly-by scenario, the intruder and the target galaxy should be connected by a stream of gas. Given the absence of this stream in emission in the GBT data, it will be interesting to see if signs of it can be detected in absorption. Alternatively, careful stacking of spectra in the supposed area of the supposed stream could yield a result, though the unknown velocities of the stream (needed to shift the spectra to a common velocity prior to stacking) could hamper this exercise.

4.3. Accretion

Alternatively, the cloud could be a sign of accretion happening. This could be accretion (or rather infall) of a dwarf galaxy, or bona-fide accretion from the IGM or cosmic web.

To start with the first possibility, the cloud and filament complex could represent the tail left behind by a dwarf galaxy as it plunges into NGC 2403 and is being stripped. Unfortunately, as we are seeing the tail projected against the bright stellar disk of NGC 2403, it will be difficult to directly detect the stellar population belonging to the dwarf galaxy. A careful study of the resolved stellar population in the region where the 8-kpc filament connects up with the main disk is probably the only way to directly detect this stellar population, if present.

The second possibility is accretion of gas directly from the IGM. Simulations show that a fraction of gas falling from the cosmic web into a galaxy halo potential does not get heated to the virial temperature of the halo (Kereš et al. 2005). These filaments of cooler gas can penetrate from the IGM through the hot gas in the halo of a galaxy and directly deliver gas to the disk. This process is commonly known as “cold accretion”. However, the gas involved still has a temperature of $\sim 10^5 \text{ K}$, and is therefore not directly comparable with the cold gas we are observing in the H I line. The cold accretion process appears to be most efficient at high redshifts ($z > 2$). At low redshifts it seems (according to the simulations) to mainly occur in low-mass galaxies with $M_{\text{halo}} \lesssim 5 \cdot 10^{11} M_\odot$ (Joung et al. 2012), potentially including NGC 2403. As noted, the “cold” accreting gas would have temperatures too high to be observed in the H I line. However,

³ We do not include the GBT cloud as its morphology is too uncertain.

further simulations (Joung et al. 2012) show that in the innermost regions of the halo ($R \lesssim 100$ kpc) some of the gas can cool further and form H I clouds. For a Milky-Way-type galaxy, simulations show that at $z = 0$ the amount of cold gas present in the halo is $\sim 10^8 M_\odot$ (Fernández et al. 2012). This is an order of magnitude higher than the H I mass of the NGC 2403 complex, but the final stages of accretion are complex and difficult to model, and it is not clear whether this difference is significant. See also Nelson et al. (2013) for a comparison of simulations done with the smoothed particle hydrodynamics and with the moving mesh techniques. In the latter the amount of cold gas and corresponding cold accretion rates are much lower.

Qualitatively, the density distribution of the filament can be consistent with an accretion scenario. The VLA observations show a dense filament close to the disk, with the GBT data showing it with a lower density further out. However, whereas the current observations are consistent with the infall of an accreting cloud, they do not conclusively prove that accretion from the IGM is indeed happening.

5. Conclusions

Deep H I observations obtained with the GBT show the presence of a $6.3 \cdot 10^6 M_\odot$ cloud near NGC 2403. Comparison with deep VLA observations by Fraternali et al. (2001, 2002) suggest that this cloud is part of a larger complex that also includes the $\sim 1 \cdot 10^7 M_\odot$ “8-kpc” anomalous-velocity H I filament in the inner disk of NGC 2403.

Deep optical observations by Barker et al. (2012) show the presence of a stellar overdensity near the position of the cloud. It is not clear however whether the overdensity and the cloud are associated. We suggest that the cloud/filament complex could be a direct observation of accretion, or, that we could be seeing the after-effects of a minor interaction (fly-by) with a neighboring galaxy.

If we are seeing accretion happening, assuming NGC 2403 is not a unique galaxy, future observations of other galaxies to similar column densities should reveal more of these events. Should these not turn up in significant numbers, then this would be a strong indication that features as now seen in NGC 2403 are interaction events. It would imply that significant *observable* accretion from the cosmic web is not happening or at least rare at low redshifts. Future, deeper observations with instruments such as the SKA, or its precursors, are required to be able to routinely probe the environment around galaxies and quantify the importance of accretion from and the relation with the cosmic web.

Acknowledgements. We thank the anonymous referee for constructive comments. We thank the staff at the GBT for their assistance with the observations. WJGdB was supported by the European Commission (grant FP7-PEOPLE-2012-CIG #333939). The work of DJP was partially supported by NSF CAREER grant AST-1149491. FF acknowledges financial support from PRIN MIUR 2010-2011, prot. 2010LY5N2T.

References

Barker, M. K., Ferguson, A. M. N., Irwin, M. J., Arimoto, N., & Jablonka, P. 2012, MNRAS, 419, 1489
 Begeman, K. G. 1987, Ph.D. Thesis, University of Groningen
 Chynoweth, K. M., Langston, G. I., Holley-Bockelmann, K., & Lockman, F. J. 2009, AJ, 138, 287
 de Blok, W. J. G., Walter, F., Brinks, E., et al. 2008, AJ, 136, 2648
 Fernández, X., Joung, M. R., & Putman, M. E. 2012, ApJ, 749, 181
 Fraternali, F. 2013, IAU Symp. 298 (arXiv:1310.2956)
 Fraternali, F., & Binney, J. J. 2006, MNRAS, 366, 449
 Fraternali, F., & Binney, J. J. 2008, MNRAS, 386, 935

Fraternali, F., Oosterloo, T., Sancisi, R., & van Moorsel, G. 2001, ApJ, 562, L47
 Fraternali, F., van Moorsel, G., Sancisi, R., & Oosterloo, T. 2002, AJ, 123, 3124
 Gentile, G., Józsa, G. I. G., Serra, P., et al. 2013, A&A, 554, A125
 Gil de Paz, A., Boissier, S., Madore, B. F., et al. 2007, ApJS, 173, 185
 Heald, G., Józsa, G., Serra, P., et al. 2011, A&A, 526, A118
 Joung, M. R., Putman, M. E., Bryan, G. L., Fernández, X., & Peek, J. E. G. 2012, ApJ, 759, 137
 Kamphuis, P., Rand, R. J., Józsa, G. I. G., et al. 2013, MNRAS, 434, 2069
 Karachentsev, I., Kaisina, E., Kaisin, S., & Makarova, L. 2011, MNRAS, 415, L31
 Kereš, D., Katz, N., Weinberg, D. H., & Davé, R. 2005, MNRAS, 363, 2
 Langston, G., & Turner, B. 2007, ApJ, 658, 455
 Mapelli, M., Moore, B., & Bland-Hawthorn, J. 2008, MNRAS, 388, 697
 McQuinn, K. B. W., Skillman, E. D., Cannon, J. M., et al. 2010, ApJ, 721, 297
 Nelson, D., Vogelsberger, M., Genel, S., et al. 2013, MNRAS, 429, 3353
 Oh, S.-H., de Blok, W. J. G., Walter, F., Brinks, E., & Kennicutt, R. C., Jr. 2008, AJ, 136, 2761
 Oosterloo, T., Fraternali, F., & Sancisi, R. 2007, AJ, 134, 1019
 Pisano, D. J. 2014, AJ, 147, 48
 Putman, M. E., Peek, J. E. G., & Joung, M. R. 2012, ARA&A, 50, 491
 Richter, O.-G., & Sancisi, R. 1994, A&A, 290, L9
 Rots, A. H. 1980, A&AS, 41, 189
 Sancisi, R., Fraternali, F., Oosterloo, T., & van der Hulst, T. 2008, A&A Rev., 15, 189
 Shapiro, P. R., & Field, G. B. 1976, ApJ, 205, 762
 Sicking, F. J. 1997, Ph.D. Thesis,
 Sinha, M., & Holley-Bockelmann, K. 2012, ApJ, 751, 17
 Swaters, R. A., Sancisi, R., & van der Hulst, J. M. 1997, ApJ, 491, 140
 Trachternach, C., de Blok, W. J. G., Walter, F., Brinks, E., & Kennicutt, R. C., Jr. 2008, AJ, 136, 2720
 van Eymeren, J., Marcelin, M., Koribalski, B., et al. 2009, A&A, 493, 511
 Walter, F., Brinks, E., de Blok, W. J. G., et al. 2008, AJ, 136, 2563
 Williams, B. F., Dalcanton, J. J., Stilp, A., et al. 2013, ApJ, 765, 120
 Zschaechner, L. K., Rand, R. J., Heald, G. H., Gentile, G., & Kamphuis, P. 2011, ApJ, 740, 35
 Zschaechner, L. K., Rand, R. J., Heald, G. H., Gentile, G., & Józsa, G. 2012, ApJ, 760, 37




Fig. 9. The field around NGC 2403. Contours and grayscales of NGC 2403 (bottom of the plot) are as in Fig. 3. The circle indicates the position and approximate size of DDO 44. In the top of the panel the H I distribution of NGC 2366 from THINGS (Walter et al. 2008) is shown. All distances and sizes are to scale.