

Role of COX-1 and COX-2 in the release of prostanoids in murine lung and isolated lung fibroblasts

William R Wright¹, Nicholas Kirkby^{1,2}, Louise Harrington¹, Martina Lundberg¹, Mark J Paul-Clark¹, Timothy D Warner² and Jane A Mitchell¹

¹ Cardiothoracic Pharmacology, National Heart and Lung Institute, Imperial College London, SW3 6LY, United Kingdom

² William Harvey Research Institute, Barts and the London School of Medicine and Dentistry, John Vane Science Centre, Charterhouse Square, London, EC1M 6BQ, United Kingdom

Correspondence: william.wright09@imperial.ac.uk; jane.a.mitchell@imperial.ac.uk

Cyclooxygenase (COX) is the first enzyme in the conversion of arachidonic acid to prostanoids. There are two isoforms of COX; COX-1, which is constitutively expressed with a homeostatic role in most tissues, and COX-2, which while constitutively expressed in some discreet sites is generally inducible by growth factors and during inflammation. In the current study, we have used tissues and cells from knock-out mice to investigate the relative contributions of COX-1 and COX-2 to PGE₂ production by lung tissue *ex vivo* and by proliferating lung fibroblasts *in vitro*.

Lung tissues from WT (C57Bl6), COX-1^{-/-} and COX-2^{-/-} mice were immediately dissected (<15 min after death) and incubated (37 °C) for 30 min in DMEM containing 50 μM calcium ionophore (A23187). Release of PGE₂ was determined by competitive immunoassay. In parallel studies, murine lung fibroblasts from COX-1^{-/-} and COX-2^{-/-} mice were explanted and cultured before being seeded in 96-well plates at sub-confluence (5000-8000/well) and incubated for 24-48 hours in the presence of 10% FCS. Accumulated release of PGE₂ was then measured as above.

Over 30 min PGE₂ was released by lung pieces from wild type (1117 ± 55 pg/ml) and COX-2^{-/-} (2013 ± 255 pg/ml) but not from COX-1^{-/-} (<61pg/ml) mice (n=4). In contrast, proliferating lung fibroblasts from COX-1^{-/-} (4978.9 ± 1392 pg/ml) mice released higher levels of PGE₂ than cells from COX-2^{-/-} (1194 ± 617 ng/ml) mice (n=4 using cells from 2-3 separate mice for each genotype).

These results show that COX-1 activity underpins the stimulated release of PGE₂ in healthy mouse lung tissue. Conversely, COX-2 activity predominates in proliferating lung fibroblasts, which may be important as COX-derived PGE₂ mediates proliferation of lung fibroblasts (Trends Immunol.2004;25(1):40-6). Our results suggest a switch in COX isoform in lung cells during proliferation which could be relevant to our understanding of conditions such as idiopathic pulmonary fibrosis.