



Asian Journal of Chemistry; Vol. 23, No. 11 (2011), 5161-5162

# ASIAN JOURNAL OF CHEMISTRY

[www.asianjournalofchemistry.co.in](http://www.asianjournalofchemistry.co.in)


## NOTE

### An Insight into Fatty Acid Composition of *Calliergonella cuspidata*

BORIS PEJIN<sup>1,\*</sup>, LJUBODRAG VUJISIC<sup>2</sup>, MARKO SABOVLJEVIC<sup>3</sup>, VELE TESEVIC<sup>2</sup> and VLATKA VAJS<sup>2</sup><sup>1</sup>Department of Organic Chemistry, Faculty of Chemistry, University of Belgrade, 11000 Belgrade, Serbia<sup>2</sup>Center of Chemistry, Institute of Chemistry, Technology and Metallurgy, University of Belgrade, 11000 Belgrade, Serbia<sup>3</sup>Institute of Botany and Garden, Faculty of Biology, University of Belgrade, 11000 Belgrade, Serbia\*Corresponding author: E-mail: [borispejin@yahoo.com](mailto:borispejin@yahoo.com)

(Received: 15 October 2010;

Accepted: 30 July 2011)

AJC-10236

The fatty acid composition of the moss *Calliergonella cuspidata* (Hedw.) Loeske (Amblystegiaceae) was preliminary analyzed by gas chromatography (GC FID) and gas chromatography-mass spectrometry (GC-MS). Six fatty acids were identified in its chloroform-methanol extract 1/1: palmitic acid (66.83 %), stearic acid (11.73 %), oleic acid (8.21 %), linolenic acid (6.01 %),  $\alpha$ -linolenic acid (3.95 %) and arachidic acid (3.26 %). Arachidonic acid and *cis*-5,8,11,14,17-eicosapentaenoic acid, typical long-chain polyunsaturated fatty acids for many mosses, were not detected in *Calliergonella cuspidata*.

**Key Words:** Moss, Amblystegiaceae, Phytochemistry, GC FID, GC-MS.

Bryophytes, the second biggest group of terrestrial plants, are known to be a reservoir of interesting and useful chemicals<sup>1</sup>. Many new compounds are described from bryophytes, mainly liverworts, which possess oil-bodies rich in terpenes and terpene-like substances. Mosses, another group of bryophytes, remain less chemically studied since its members do not possess oil-bodies in their cells<sup>2</sup>. Fatty acids are known to be present within mosses but there are many assumptions and generalizations from a small number of species studied<sup>3-5</sup>. Therefore, in this study the moss *Calliergonella cuspidata* (Hedw.) Loeske (Amblystegiaceae) has been screened on those chemical constituents and found to be rich in saturated fatty acids. According to our best of knowledge, there is no previous report on fatty acid chemistry of this moss species.

The moss was collected in the surrounding of Bonn (Germany) in December, 2007. Voucher specimen has been deposited in the Herbarium of the Institute of Botany, University of Belgrade, Serbia (bryophyte collection-BEOU No. 4704).

*C. cuspidata* was carefully selected and cleaned from soil and other contaminants. The gametophyte tips were used for the extraction. Air-dried parts of the sample were ground (1 g) and extracted three times with the chloroform-methanol 1/1 for 1 h at room temperature. The extract was evaporated to dryness and further *trans*-esterified with 5 % H<sub>2</sub>SO<sub>4</sub> in MeOH (v/v) for 4 h at 80 °C. The resulting methyl esters of fatty

acids were analyzed by comparing its GC FID chromatogram with that of standard mixture (Supelco 37) obtained under the same conditions and/or by analysis of GC-MS data using NIST 5 and Wiley 7 libraries.

GC analysis was performed on Agilent 7890A GC system equipped with 5975C MSD and FID detector, using DB-23 column (30 m × 0.25 mm × 0.25 μm). Injection volume was 1 μL and injector temperature was 220 °C with 10:1 split ratio. Carrier gas (He) flow rate was 0.9 mL/min while column temperature was linearly programmed in a range of 150-240 °C at a rate of 4 °C/min and hold at 240 °C for 10 min. Transfer line was heated at 240 °C. The FID detector temperature was 300 °C. EI mass spectra (70 eV) were acquired in *m/z* range 40-500.

Six fatty acids were identified in its chloroform-methanol extract 1/1: palmitic acid (66.83 %), stearic acid (11.73 %), oleic acid (8.21 %), linolenic acid (6.01 %),  $\alpha$ -linolenic acid (3.95 %) and arachidic acid (3.26 %). Arachidonic acid and *cis*-5,8,11,14,17-eicosapentaenoic acid, typical long-chain polyunsaturated fatty acids for many moss species<sup>6,7</sup>, were not detected in the examined species. This may be the consequence of the period when the moss has been collected (temperature, water regime or both, *etc.*), so further studies in seasonal *i.e.* environmental variation of these chemical constituents in *C. cuspidata* are needed.

**ACKNOWLEDGEMENTS**

This study was supported by the Ministry of Science and Technological Development of the Republic of Serbia (Research grants No. 142053 and 143015).

**REFERENCES**

1. S. Sabovljevic and M. Sabovljevic, in ed.: J.N. Govil, Bryophytes, A Source of Bioactive New Compounds, Phytopharmacology and Therapeutic Values IV, Studium Press, Houston, Texas, pp. 9-25 (2008).
2. V. Herout, in eds.: H.D. Zinsmeister and R. Mues, Diterpenes and Higher Terpenes from Bryophytes, Bryophytes, their Chemistry and Chemical Taxonomy, Clarendon Press, Oxford, pp. 83-102 (1990).
3. S. Huneck, in ed.: R.M. Shuster, Chemistry and Biochemistry of Bryophytes, New Manual of Bryology, The Hatory Botanical Laboratory, Nichian, Miyazaki, Vol. 1, pp. 1-116 (1983).
4. P. Karunen, in eds.: H.D. Zinsmeister and R. Mues, The acyl lipids of Bryophytes, Bryophytes, their Chemistry and Chemical Taxonomy, Clarendon Press, Oxford, pp. 121-141 (1990).
5. V.M. Dembitsky, *Prog. Lipid Res.*, **32**, 281 (1993).
6. P. Karunen, *J. Hattori Bot. Lab.*, **53**, 255 (1982).
7. I.A. Guschina and J.L. Hardwood, *J. Exp. Bot.*, **53**, 455 (2002).