

Supplementary data for article :

Šovrc, L.; Borovska, K.; Cinková, K.; Stanković, D.; Planková, A. Advanced Electrochemical Platform for Determination of Cytostatic Drug Flutamide in Various Matrices Using a Boron-Doped Diamond Electrode. *Electrochimica Acta* **2017**, *251*, 621–630.

<https://doi.org/10.1016/j.electacta.2017.08.077>

SUPPLEMENTARY INFORMATION

Advanced electrochemical platform for determination of cytostatic drug flutamide in various matrices using a boron-doped diamond electrode

Lubomír Švorc^{a*}, Katarína Borovská^a, Kristína Cinková^a, Dalibor M. Stanković^{b,c},
Alexandra Planková^d


^aInstitute of Analytical Chemistry, Faculty of Chemical and Food Technology, Slovak University of Technology in Bratislava, Radlinského 9, Bratislava, SK-812 37, Slovak Republic

^bDepartment of Analytical Chemistry, Innovation Center of the Faculty of Chemistry, University of Belgrade, Studentski trg 12-16, Belgrade, 11000, Serbia

^cInstitute of Nuclear Sciences “Vinča”, University of Belgrade, P. O. Box 522, Belgrade, 11000, Serbia

^dDepartment of Pharmaceutical Analysis and Nuclear Pharmacy, Faculty of Pharmacy, Comenius University in Bratislava, Odbojárov 10, Bratislava, SK-832 32, Slovak Republic

Fig. S1



The linear dependences (left: $\log I_p$ against $\log f$, right: E_p against $\log f$) evaluated from SWV records of 1 mM FLU for the set of the frequencies (f): (a) 5, (b) 10, (c) 25, (d) 50, (e) 100 and (f) 150 Hz in 0.1 M H_2SO_4 on the BDD electrode (Fig. 2). The other SWV parameters: amplitude of 50 mV and step potential of 1 mV.