

Supplementary data for the article:

Stojsavljević, A.; Trifković, J.; Rasić-Milutinović, Z.; Jovanović, D.; Bogdanović, G.; Mutić, J.; Manojlović, D. Determination of Toxic and Essential Trace Elements in Serum of Healthy and Hypothyroid Respondents by ICP-MS: A Chemometric Approach for Discrimination of Hypothyroidism. *Journal of Trace Elements in Medicine and Biology* **2018**, *48*, 134–140. <https://doi.org/10.1016/j.jtemb.2018.03.020>

Supplementary material

Determination of toxic and essential trace elements in serum of healthy and hypothyroid respondents by ICP-MS: A chemometric approach for discrimination of hypothyroidism

Aleksandar Stojsavljević^a, Jelena Trifković^a, Zorica Rasić-Milutinović^b, Dragana Jovanović^c, Gradimir Bogdanović^d, Jelena Mutić^a, Dragan Manojlović^a

^a*University of Belgrade - Faculty of Chemistry, Studentski trg 12-16, 11100 Belgrade, Serbia*

^b*Department of Endocrinology, Faculty of Medicine, University Hospital Zemun, Belgrade, Serbia*

^c*Institute of Public Health of Serbia "Dr Milan Jovanovic Batut", Belgrade, Serbia*

^d*Blood Transfusion Institute, Belgrade, Serbia*

Content

Figure S1. Plots of residuals for linear regression model versus predicted values

Table S1. The *F*-value of the Lack-of-fit (**LOF**) test and coefficient of determination (**R²**) for linear (**LRM**) and quadratic regression model (**QRM**)




Figure S1. Plots of residuals for linear regression model versus predicted values

Table S1. The F -value of the Lack-of-fit (LOF) test and coefficient of determination (R^2) for linear (LRM) and quadratic regression model (QRM) obtained for investigated calibration curves

	LRM		QRM	
	LOF^a	R^2	LOF^b	R^2
Cr	3.995	0.9980	0.321	0.9994
Mn	1.528	0.9968	0.938	0.9978
Co	0.561	0.9985	0.181	0.9998
Ni	0.649	0.9972	0.034	0.9999
As	0.297	0.9989	0.021	0.9994
Cd	0.138	0.9983	0.089	0.9989
Cu	1.029	0.9949	0.379	0.9999
Zn	1.342	0.9874	0.173	0.9988
Se	0.313	0.9975	0.207	0.9995

^a $F_{cr, 95\%} = 4.534$

^b $F_{cr, 95\%} = 4.757$