

Supplementary data for article:

Stanković, I. M.; Božinovski, D. M.; Brothers, E. N.; Belić, M. R.; Hall, M. B.; Zarić, S. D.
Interactions of Aromatic Residues in Amyloids: A Survey of Protein Data Bank Crystallographic
Data. *Crystal Growth and Design* **2017**, *17* (12), 6353–6362.
<https://doi.org/10.1021/acs.cgd.7b01035>

Supporting Information for the Manuscript

Interactions of aromatic residues in amyloids: A Survey of PDB Crystallographic Data

Authors:

Ivana M. Stanković¹, Dragana M. Božinovski², Edward N. Brothers³, Milivoj R. Belić³, Michael B. Hall⁴ and Snežana. D. Zarić^{2,3}*

¹ University of Belgrade, Institute of Chemistry, Technology and Metallurgy, Njegoševa 12, 11000 Belgrade, Serbia. ² Department of Chemistry, University of Belgrade, Studentski trg 12-16, 11000 Belgrade, Serbia. ³ Science Program, Texas A&M University at Qatar, Texas A&M Engineering Building, Education City, Doha, Qatar. ⁴ Department of Chemistry, Texas A&M University, College Station, TX 77843-3255, USA

* Corresponding author:

Prof. Dr. Snežana Zarić




Fig. S1. Turn and coil aromatic amino acids (in red). PDBids: 2e8d, 2lmo.




Fig. S2. The structure with Trp-Trp intrasheet interactions, PDBid 2nnt.




Fig. S3. The 20-framed NMR structure, PDBid 2m5n. An example of the high-offset interactions as a consequence of geometry, rings in red. $P_1/P_2 = 36.09^\circ$, $R = 1.78 \text{ \AA}$, $r = 5.91 \text{ \AA}$, $d = 6.17 \text{ \AA}$. Surrounding interactions are intrasheet aromatic-aromatic interactions.