

Supplementary data for article:

Kekez, B. D.; Gojgic-Cvijovic, G. D.; Jakovljevic, D. M.; Stefanovic Kojic, J. R.;
Markovic, M. D.; Beskoski, V. P.; Vrvic, M. M. High Levan Production by *Bacillus*
Licheniformis NS032 Using Ammonium Chloride as the Sole Nitrogen Source. *Applied*
Biochemistry and Biotechnology **2015**, *175* (6), 3068–3083. <https://doi.org/10.1007/s12010-015-1475-8>

Supplementary Material

High levan production by *Bacillus licheniformis* NS032 using ammonium chloride as the sole nitrogen source

B. D. Kekez^a, G. D. Gojgic-Cvijovic^{b,*}, D. M. Jakovljevic^b, J. R. Stefanovic Kojic^b, M. D. Markovic^b, V. P. Beskoski^a, M. M. Vrvic^{a,b}


^aFaculty of Chemistry, University of Belgrade, Studentski trg 12-16, P.O. Box 51, 11000 Belgrade, Serbia,

^bInstitute of Chemistry, Technology and Metallurgy, Department of Chemistry, University of Belgrade, Njegoseva 12, P.O. Box 473, 11000 Belgrade, Serbia

* Corresponding author: Gordana Gojgic-Cvijovic, Institute of Chemistry, Technology and Metallurgy, Department of Chemistry, University of Belgrade, Njegoseva 12, P.O. Box 473, 11000 Belgrade, Serbia Tel.:+381 11 2637273; Fax: +381 11 2636061, Email: ggojgic@chem.bg.ac.rs


Supplementary Material, Figure S1

ATR-FTIR spectrum of the polysaccharide produced by *B. licheniformis* NS032



Supplementary Material, Figure S2

^{13}C NMR spectrum of the polysaccharide produced by *B. licheniformis* NS032



Supplementary Material, Table S3

Change of pH values during levan production by *B. licheniformis* NS032 in the base medium with different nitrogen sources

N source	Time, hours			
	24	48	72	96
YE/BE	6.08	5.78	5.33	5.49
YE	6.21	5.48	5.31	5.32
BE	6.35	5.65	5.37	5.42
Peptone	6.42	5.91	5.42	5.59
(NH ₄) ₂ SO ₄	6.02	5.43	4.73	4.75
NH ₄ Cl	5.94	5.30	4.91	4.57
NaNO ₃	6.80	6.59	6.21	6.26

Medium: BM with 100 g /L sucrose, initial pH 7.0, YE yeast extract, BE- beef extract; see Material and methods for details and Fig. 1d