

Supplementary data for article:

Božinović, N.; Ajdačić, V.; Lazic, J.; Lecerf, M.; Daventure, V.; Nikodinovic-Runic, J.; Opsenica, I. M.; Dimitrov, J. D. Aromatic Guanylhydrazones for the Control of Heme-Induced Antibody Polyreactivity. *ACS Omega* **2019**, *4* (24), 20450–20458.

<https://doi.org/10.1021/acsomega.9b01548>

Supporting Information

Aromatic guanylhyazones for the control of heme-induced antibody polyreactivity

Nina Božinović*,§, Vladimir Ajdačić#, Jelena Lazic#, Maxime Lecerf§, Victoria Daventure§, Jasmina Nikodinovic-Runic#,* , Igor M. Opsenica# and Jordan D. Dimitrov* ,§

§Centre de Recherche des Cordeliers, INSERM, Sorbonne Université, USPC, Université Paris Descartes, Université Paris Diderot, F-75006 Paris, France. #University of Belgrade – Faculty of Chemistry, Studentski trg 16, P.O. Box 51, 11158 Belgrade, Serbia. *Institute of Molecular Genetics and Genetic Engineering, University of Belgrade, Vojvode Stepe 444a, 11000 Belgrade, Serbia

Figure S1: Spectroscopic study of 20 μM hemin with 100 μM guanyldiazones. Differential spectra obtained by subtraction of the individual spectra of hemin and the compound from the spectrum of their mixture.

