

# Impact of short-term storage on the quantity of extended-spectrum beta-lactamase–producing *Escherichia coli* in broiler litter under practical conditions

Paul Siller,<sup>\*,1</sup> Katrin Daehre,<sup>\*,†</sup> Nadine Thiel,<sup>‡</sup> Ulrich Nübel,<sup>‡,§,#</sup> and Uwe Roesler<sup>\*</sup>

<sup>\*</sup>Institute for Animal Hygiene and Environmental Health, Freie Universität Berlin, Berlin, Germany; <sup>†</sup>Department of Food, Feed and Commodities, Federal Office of Consumer Protection and Food Safety, Berlin, Germany; <sup>‡</sup>Department of Microbial Genome Research, Leibniz Institute DSMZ-German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany; <sup>§</sup>Braunschweig Integrated Center of Systems Biology (BRICS), Technical University, Braunschweig, Germany; and <sup>#</sup>German Center for Infection Research (DZIF), Partner site Hannover-Braunschweig, Germany

**ABSTRACT** Applying broiler litter containing extended-spectrum beta-lactamase (ESBL)–producing *Escherichia coli* (*E. coli*) to arable land poses a potential risk for humans to get colonized by contact with contaminated soil or vegetables. Therefore, an inactivation of these bacteria before land application of litter is crucial. We performed 2 short-term litter storage trials (one in summer and winter, respectively), each covering a time span of 5 D to investigate the effectiveness of this method for inactivation of ESBL-producing *E. coli* in chicken litter. Surface and deep litter samples were taken from a stacked, ESBL-positive chicken litter heap in triplicates in close sampling intervals at the beginning and daily for the last 3 D of the experiments. Samples were analyzed quantitatively and qualitatively for ESBL-producing *E. coli*, total *E. coli*, and enterococci. Selected isolates were further characterized by whole-genome sequencing (WGS). In the depth of the heap ESBL-producing *E. coli* were detected quantitatively

until 72 h and qualitatively until the end of the trial in winter. In summer detection was possible quantitatively up to 36 h and qualitatively until 72 h. For surface litter samples a qualitative detection of ESBL-producing *E. coli* was possible in all samples taken in both trials. In the deep samples a significant decrease in the bacterial counts of over 2 Log<sub>10</sub> was observed for total *E. coli* in the winter and for total *E. coli* and enterococci in the summer. Genetic differences of the isolates analyzed by WGS did not correlate with survival advantage. In conclusion, short-term storage of chicken litter stacked in heaps is a useful tool for the reduction of bacterial counts including ESBL-producing *E. coli*. However, incomplete inactivation was observed at the surface of the heap and at low ambient temperatures. Therefore, an extension of the storage period in winter as well as turning of the heap to provide aerobic composting conditions should be considered if working and storage capacities are available on the farms.

**Key words:** antibiotic resistance, ESBL, *E. coli*, broiler litter, environment

2020 Poultry Science 99:2125–2135

<https://doi.org/10.1016/j.psj.2019.11.043>

## INTRODUCTION

Extended-spectrum beta-lactamases (ESBL) are enzymes occurring in Enterobacteriaceae. Their ability to hydrolyze the  $\beta$ -lactam ring of a variety of  $\beta$ -lactam antibiotics including extended-spectrum cephalosporins of the third and fourth generation leads to an inactivation

of antibiotic properties. Cephalosporins of the third and fourth generation have a broad-spectrum activity against Gram-positive and Gram-negative bacteria and are often used for the treatment of infections in intensive care units. The emergence of resistance against these drugs limits therapeutic options (Remschmidt et al., 2017).

ESBL-producing *Escherichia coli* (*E. coli*) are commonly found in broiler production with a prevalence of up to 100% in fattening farms (Dierikx et al., 2010, 2013; Laube et al., 2013; Blaak et al., 2015; Hering et al., 2016; Daehre et al., 2018). Furthermore, ESBL-producing *E. coli* have been detected in the vicinity of broiler barns, and an airborne and waterborne

© 2019 Published by Elsevier Inc. on behalf of Poultry Science Association Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Received September 30, 2019.

Accepted November 14, 2019.

<sup>1</sup>Corresponding author: [paul.siller@fu-berlin.de](mailto:paul.siller@fu-berlin.de)

dissemination have been described (Laube et al., 2014; Blaak et al., 2015). In Germany, 600 million broiler chickens were slaughtered in 2017 (Statistisches Bundesamt, 2017) contributing to the 1.1 million metric tons of poultry litter that are spread to arable land in Germany annually (Statistisches Bundesamt, 2016). This presents a possible important emission source of resistant bacteria from the barns to the environment.

Blaak et al. (2015) reported that ESBL-producing *E. coli* were found in the soil at a distance of 1–5 m of litter storage areas with up to  $2.0 \times 10^4$  cfu/kg. Additionally, it was shown that ESBL-producing *E. coli* can be transferred from animal husbandry to soil and are able to survive on the fields for at least 1 y (Hartmann et al., 2012). A cross-transmission of ESBL-producing *E. coli* and mobile genetic elements encoding for the production of ESBL between animals, including chickens, humans, and the environment is hypothesized (Leverstein-van Hall et al., 2011; Huijbers et al., 2014). The spread of litter containing ESBL-producing Enterobacteriaceae to the environment poses a potential risk for humans to be colonized with these bacteria after contact with contaminated soils or via contaminated vegetables. Hence, inactivation of these resistant bacteria before land application is crucial.

Storing the litter in piles after removal from the barns could be a useful and cost efficient tool for the reduction of resistant bacteria in litter. Studies that investigated the reduction of nonresistant *E. coli* in chicken litter by storage (anaerobic conditions) and composting (active aeration) under practical conditions were performed previously (Erickson et al., 2010; Wilkinson et al., 2011). Considering practicability and economic sustainability short-term storage of litter presents the most advantageous method of litter treatment. The decline of ESBL-producing *E. coli* in chicken litter under field conditions has not been investigated so far. For a more detailed assessment of bacterial inactivation in short-term chicken litter storage, concentrations of nonresistant *E. coli* and enterococci were additionally monitored in this study. Enterococci are approved gram-positive indicator microorganisms present in feces and have a higher tenacity compared to *E. coli*. Two short-term storage trials each covering a time span of 5 D were performed. One trial was performed in the summer (summer trial) and one in the winter (winter trial) to explore climatic influences on the decline of these bacteria.

## MATERIALS AND METHODS

### Experimental Design

An initial screening of 40 different barns of a large broiler farm in Germany was performed to select barns with high quantities of ESBL-producing *E. coli* in the litter. Screening consisted of one boot swab and one composite litter sample per barn and was done for both the summer trial and the winter trial. The winter trial was carried out in early March, and the summer trial in early

June. The time span between the screening and the litter storage trial was 2 wk for the winter trial and 1 wk for the summer trial.

In all barns, 1.5 kg/m<sup>2</sup> of wood pellets were used as bedding material. The litter (approximately 15 metric tons for each trial) was removed from the barns with a front-end loader and piled up on a concrete surface behind the barn directly after the chickens were housed out.

Storing the litter behind the barns for several days is not unusual in broiler production. Although on the farm where the trials were performed, the litter is removed as fast as possible if working capacities are available. Under suitable conditions, it is used directly for fertilization or otherwise transported to further storage areas.

The first samples were taken immediately after the litter heap was stacked. For both trials, the litter heaps were sampled at 6 points in time: 12 h, 24 h, 36 h, 48 h, 72 h, and 96 h after storage begin. We expected a faster reduction of the bacterial counts in the summer because of the higher ambient temperatures. Hence, in the summer trial, additional samplings were performed at 1 h, 3 h, and 6 h after the begin of storage. At each point in time, 3 surface and 3 deep samples were taken from the litter heap.

For the surface litter samples, approximately 50 g of litter from the heap's surface were collected in sterile 120 mL specimen containers (VWR, Radnor, PA)

For the deep litter samples, custom-built steel sample containers were used. These containers are cylindrical, 9.9 cm long, have a diameter of 4.4 cm, and drill holes with a diameter of 7 mm, ensuring the same environmental conditions in the sampling container and the surrounding litter heap. These sterilized containers were filled with litter from the heap and were placed in the litter heap at a depth of 50 to 55 cm at the start of the experiment. Wires were attached to the containers allowing quick retraction and sample collection at each point in time.

The ALMEMO 2490 device (AHLBORN, ZA9020-FS and FH A696-GF1 Holzkirchen, Germany) was used to record temperature and moisture at each sampling spot immediately after sampling.

The weather data for the trial periods were obtained from the closest weather station located approximately 20 km from the sampling site (Archive of the German Meteorological Office)

### pH Value Analysis

The pH value was measured for all litter samples. Samples were diluted with purified water at a ratio of 1:10 and homogenized for 30 s with a vortex mixer. The pH value was measured with the handheld measurement instrument AL10 (AQUALYTIC, Dortmund, Germany).

### Microbiological Analyses

All boot swabs and litter samples from the screenings were analyzed quantitatively and qualitatively for ESBL-producing *E. coli*. Litter samples from the litter

storage trials were analyzed quantitatively and qualitatively for total *E. coli* and ESBL-producing *E. coli*. Additionally, enterococci were analyzed quantitatively.

All litter samples were mixed with Luria/Miller-broth (LB) (Roth, Karlsruhe, Germany) in stomacher bags at a ratio of 1:10. Boot swabs were put in stomacher bags, and 200 ml of LB medium was added. The samples were homogenized using a Stomacher 400 Circulator (Seward Limited, West Sussex, UK) for 2 min at 200 rpm.

For the quantitative analyses aliquots of the suspensions were taken, and triplicates of 100 µL were streaked on specific agar plates after serial dilution. For *E. coli*, MacConkey agar No. 3 (Oxoid, Wesel, Germany) was used. To detect ESBL-producing *E. coli*, 1 mg/L cefotaxime (AppliChem, Darmstadt, Germany) was added as suggested by the EFSA (2011). For enterococci, we used Bile Aesculin Azide agar (Merck, Darmstadt, Germany). We did not set a minimum number of cfu per plate for the evaluation of the undiluted samples, resulting in a quantitative detection limit of  $3.3 \times 10^1$  cfu/g of litter.

For qualitative testing, the homogenized samples were incubated in LB medium for 20 to 24 h at 37°C. Subsequently, 10 µL were streaked on MacConkey agar with and without the addition of cefotaxime, respectively, with an inoculation loop.

MALDI-TOF Mass Spectrometry (MALDI Microflex LT and Biotyper database, Bruker Daltonics, Bremen, Germany) was used for species confirmation of colonies which were phenotypically suspected to be *E. coli* or enterococci.

### Real-Time PCR and Sanger Sequencing

Real-time qPCR as described by Roschanski et al. (2014) was used to detect the most important beta-lactamase genes *bla*<sub>SHV</sub>, *bla*<sub>TEM</sub>, *bla*<sub>CTX-M</sub>, and the CIT-type AmpC *bla*<sub>CMY-2</sub> in isolates of all samples.

For both trials, the ESBL-gene of 8 isolates were sequenced by Sanger sequencing to identify the present ESBL-variants. Two isolates from the litter and one isolate from a surface and deep litter sample for day 1, 3, and 5 were chosen for sequencing for each trial, respectively. All isolates that showed a *bla*<sub>TEM</sub> resistance gene in addition to the predominant resistance gene were also chosen for sequencing.

The DNA was isolated, and PCR was performed as published previously by Projahn et al. (2017). The purified PCR products were sent to LGC Genomics (Berlin, Germany) who provided the sequences. Nucleotide sequences were analyzed using DNASTAR Lasergene (Madison, WI) and compared with the reference sequences of GenBank (<https://www.ncbi.nlm.nih.gov/genbank/>) according to the accession numbers of the lahey database (<https://www.lahey.org/studies/>).

### Phylotyping

All isolates were analyzed for their phylogenetic group as published by Clermont et al. (2013) with modified PCR conditions according to Projahn et al. (2017).

Isolates that could not be assigned to a phylogroup because of unspecific band patterns were declared as a combined phylogroup.

### Whole Genome Sequencing

Forty-four ESBL-producing *E. coli* isolates were selected for whole genome sequencing and recultivated on Brain-Heart-Infusion agar (Roth, Karlsruhe, Germany). DNA was isolated using the Qiagen Blood and Tissue Kit (Qiagen, Hilden, Germany), and sequencing libraries were prepared using the Nextera XT protocol with modifications (Baym et al., 2015; Steglich et al., 2018). The libraries were sequenced on a NextSeq machine with a NextSeq 500/550 mid output v2 kit (Illumina, San Diego) to  $\geq 50$ -fold sequencing coverage.

Short-read sequencing data were uploaded to the online platform Enterobase (<https://enterobase.warwick.ac.uk>), where they were assembled. Resulting contigs were quality-controlled and subjected to classification by 7-gene multilocus sequence typing (MLST), core-genome multilocus sequence typing (cgMLST), and cgMLST-based hierarchical clustering. Clusters at the level HC1100 (Hierarchical Cluster 1,100, that is, chains of genomes differing pairwise by maximally 1,100 cgMLST alleles) represent major genetic populations within the species *E. coli*, largely congruent with sequence-type complexes based on legacy 7-gene MLST (Zhou et al., 2019). In addition, Enterobase used genome sequence information to predict phylogroups according to Clermont typing (Zhou et al., 2019) based on algorithms by Beghain et al. (2018) and Waters et al. (2018).

Genome sequences were screened for antibiotic resistance genes by using the tools Resfinder (Zankari et al., 2012), AMRFinder (Feldgarden et al., 2019), and CARD (Jia et al., 2017), as implemented in ABRicate (<https://github.com/tseemann/abricate>). Genome sequencing data were submitted to the European Nucleotide Archive (<https://www.ebi.ac.uk/ena>) under study number PRJEB34161.

### Statistical Analysis

The software SPSS, version 25, (SPSS, Inc., Chicago, IL) was used for statistical analysis. The data on microbial counts had no normal distribution. We used log transformation to achieve log normal distribution, and geometric means were calculated as proposed by Bland and Altman (1996). The upper and lower 95% confidence intervals were calculated. The winter trial and summer trial were analyzed separately.

## RESULTS

### Environmental Conditions During the Litter Storage Trials Conducted in Summer and Winter

The relevant weather data for the period of both trials are summarized in Table 1.

**Table 1.** Environmental conditions during the short-term litter storage trials as provided by the weather station closest to the trial site (German meteorological office).

Trial	Day	Daily air temperature (°C)			RH (%) <sup>1</sup>	Sunshine (h)	Rainfall (mm)
		Minimum	Maximum	Mean			
Winter	1	-13.1 °C	-2.0 °C	-7.2 °C	58.8%	9.5 h	0.0 mm
	2	-12.8 °C	-0.6 °C	-6.1 °C	56.5%	9.4 h	0.0 mm
	3	-7.7 °C	7.4 °C	-0.5 °C	54.7%	5.1 h	1.1 mm
	4	-3.8 °C	11.9 °C	4.4 °C	74.5%	7.9 h	0.0 mm
	5	-5.5 °C	5.2 °C	1.0 °C	82.6%	0.9 h	6.9 mm
Summer	1	13.6 °C	28.8 °C	21.6 °C	75.6%	10.0 h	20.4 mm
	2	13.6 °C	30.4 °C	23.5 °C	69.9%	12.3 h	0.0 mm
	3	14.9 °C	27.0 °C	21.6 °C	78.8%	6.5 h	0.2 mm
	4	14.4 °C	21.5 °C	19.2 °C	86.1%	0.3 h	0.1 mm
	5	13.7 °C	25.2 °C	20.5 °C	76.0%	5.4 h	0.0 mm

<sup>1</sup>Relative humidity.

### Temperature, Moisture, and pH Value of the Litter

In both trials, the temperature of the litter increased continuously at a depth of 50 to 55 cm compared with the surface of the litter heap. In the winter trial, 50.4°C were reached at the end of the trial period (96 h). In the summer trial, temperatures over 50°C were already reached after 36 h, and the maximum temperature measured was 58.5°C at the end of the trial (96 h). The temperature in the surface samples was lower for both trials ranging from 16.8°C to 24.2°C in the winter trial and 29.3°C to 42.7°C in the summer trial.

In the winter trial, moisture of the deep samples ranged from 16.2 to 23.2%. In the summer trial, moisture of the deep samples increased from approximately 9.0% at the beginning of the experiment to 27.9% at the end of the experiment. Surface samples from both trials showed lower moisture levels with values ranging from 5.8 to 8.6% in the winter trial and 5.9 to 8.8% in the summer trial. Temperature and moisture development for both trials is shown in Figure 1.

The pH values measured in the chicken litter directly after the removal from the barn was 8.0 in the winter trial and 8.1 in the summer trial.

For both trials, the pH value increased in the surface samples over 5 consecutive day up to a maximum pH value of 8.8 in the winter trial and 8.6 in the summer

trial. In the deep samples, the pH value decreased in both trials to a minimum of 5.6 in the winter trial and 6.5 in the summer trial. The mean pH values for both trials are depicted in Figure 2.


### Microbiological Status of the Barns in the Initial Screenings


For both trials, the barn with the highest concentration of ESBL-producing *E. coli* in the litter samples of the initial screenings was chosen for the litter storage trial. The bacterial counts of the investigated microorganisms in the barns are shown in Table 2 for the boot swabs and composite litter samples.

### Quantitative and Qualitative Detection of ESBL-Producing *E. coli* in the Litter

Surface litter samples were firstly taken immediately after the litter was removed from the barns (0 h). The mean number of ESBL-producing *E. coli* per g of litter was  $5.2 \times 10^4$  cfu in the winter trial and  $9.5 \times 10^2$  cfu in the summer trial.

In the winter trial, the mean number of ESBL-producing *E. coli* was  $1.3 \times 10^4$  cfu/g of litter at the 12 h sampling point for the surface samples, decreased until 36 h and increased again to  $5 \times 10^3$  cfu/g after

**Figure 1.** Development of mean sample temperatures and moistures in the winter trial (A) and summer trial (B).



**Figure 2.** Mean sample pH values in the winter trial (A) and summer trial (B) for each point in time.

72 h. At the end of the trial (96 h), ESBL-producing *E. coli* were not quantitatively detectable.

The deep litter samples in the winter trial showed an ESBL-producing *E. coli* concentration of  $3.6 \times 10^3$  cfu/g at 12 h and gradually decreased in the following measurements. For the 72 h and 96 h samples, ESBL-producing *E. coli* could not be detected quantitatively. A qualitative detection was possible for 33% ( $n = 1/3$ ) of the samples after 72 h and for 66% ( $n = 2/3$ ) of the samples after 96 h.

In the summer trial, the number of ESBL-producing *E. coli* in the surface samples decreased slightly in the first 12 hours (mean at 1 h =  $2 \times 10^2$  cfu/g, mean at 12 h =  $9.3 \times 10^1$  cfu/g). At 24 h and 72 h, the concentration of ESBL-producing *E. coli* was higher than the initial count with up to  $6.7 \times 10^3$  cfu/g. At 36 h, the number of ESBL-producing *E. coli* was below the detection limit.

ESBL-producing *E. coli* were not quantitatively detectable in the deep samples of the summer trial for the 6 h point in time. Additionally, after 36 h for the last 4 points in time (36 h, 48 h, 72 h, and 96 h), ESBL-producing *E. coli* were constantly under the detection limit. The qualitative analysis was only positive for 67% ( $n = 2/3$ ) of the samples after 48 h and for 0% ( $n = 0/3$ ) after 72 h and 96 h, respectively.

A qualitative detection of ESBL-producing *E. coli* was possible for all surface samples in both trials. The data on ESBL-producing *E. coli* are shown in Figure 3.

### Quantitative and Qualitative Detection of Total *E. coli* in the Litter

In the litter samples taken directly after the litter was removed from the barns (0 h), the mean number of *E.*

*coli* per g of litter was  $5.6 \times 10^6$  cfu for the winter trial and  $5.8 \times 10^4$  cfu for the summer trial.

In the winter trial, the number of *E. coli* dropped in both, the surface and the deep samples. After 96 h, the mean amount of *E. coli* was  $1.1 \times 10^4$  cfu/g of litter in the surface and  $1.1 \times 10^3$  cfu/g of litter in the deep samples. Qualitative *E. coli* detection was possible in 66% ( $n = 2/3$ ) of the deep samples at 72 h and 96 h.

In the summer trial, on the other hand, the mean number of *E. coli* in surface litter samples increased from  $5.8 \times 10^4$  cfu/g of litter at 0 h to a maximum of  $2.8 \times 10^6$  cfu/g at 72 h and  $2.4 \times 10^6$  cfu/g at 96 h. Deep litter samples showed a constant decrease in *E. coli* concentrations with a drop below the detection limit at 72 h. A qualitative detection was possible for 67% ( $n = 2/3$ ) of samples at 48 h and for 0% ( $n = 0/3$ ) of samples at 72 h and 96 h. The data for total *E. coli* are shown in Figure 4.

### Quantitative Detection of Enterococci in the Litter


The number of enterococci for the 0 h point in time was  $4.8 \times 10^7$  cfu/g of litter for the winter trial and  $5.5 \times 10^6$  cfu/g for the summer trial, respectively.

In the winter trial, the quantity of enterococci was comparatively stable for both, surface and deep samples, ranging from  $3.2 \times 10^6$  to  $5.6 \times 10^7$  cfu/g in all samples taken.

In the summer trial, the quantity of enterococci increased in surface samples in the sampling period, reaching a maximum of  $2.7 \times 10^8$  cfu/g after 48 h. The quantity at the end of the sampling period (96 h) was  $1.6 \times 10^8$  cfu/g of litter.

**Table 2.** Bacterial counts for the initial screening of the barns in cfu/boot swab and cfu/g of litter.

Trial	Type of sample	ESBL-producing <i>E. coli</i> in cfu/boot swab or cfu/g	Total <i>E. coli</i> in cfu/boot swab or cfu/g	Enterococci in cfu/boot swab or cfu/g
Winter	Boot swab	$1.1 \times 10^6$ cfu/boot swab	$2.3 \times 10^7$ cfu/boot swab	$2.7 \times 10^8$ cfu/boot swab
	Litter sample	$3.9 \times 10^4$ cfu/g	$4 \times 10^9$ cfu/g	$2.9 \times 10^7$ cfu/g
Summer	Boot swab	$3.6 \times 10^6$ cfu/boot swab	$3.6 \times 10^8$ cfu/boot swab	$5.9 \times 10^8$ cfu/boot swab
	Litter sample	$6.3 \times 10^5$ cfu/g	$1.7 \times 10^7$ cfu/g	$7.2 \times 10^7$ cfu/g



**Figure 3.** Results of the quantitative and qualitative analysis of ESBL-producing *E. coli* in cfu/g of litter for the winter trial samples (A) and the summer trial samples (B). The geometric mean of 3 samples is shown for each point in time. The error bars indicate the upper and lower 95% confidence intervals. The graphs were shifted to improve the visibility of the error bars. ESBL, extended-spectrum beta-lactamase.

In contrast, in the summer trial, deep litter samples showed a significant decrease in enterococci concentration. A mean of  $1.3 \times 10^4$  and  $3.4 \times 10^4$  cfu/g were detected after 72 h and 96 h, respectively. The data for enterococci are shown in Figure 4.

## Molecular Characterization of ESBL-Producing *E. coli*

**Phylogroups and Sequence Types** In the winter trial, the phylogenetic group was determined for 48 ESBL-producing *E. coli* isolates using classical gel-based PCR. Forty-seven isolates were allocated to phylogroup F, and one isolate was allocated to phylogroup A/C.


In the summer trial, the number of ESBL-producing *E. coli* isolates available for phylogenetic analysis was 54 of which 17 isolates belonged to phylogroup B1. One isolate belonged to the groups A and F, respectively, and 35 isolates were allocated to the combined phylogroup D/E.

In Table 3, a comparison of the Clermont phylogroups determined by PCR and the by whole genome sequenced isolates is shown. It also provides information on the 7-gene MLST, cgMLST, and HC1100 clustering from the Enterobase analyses and the predicted O and H antigens for all isolates.

**Resistance Genes** Real time qPCR revealed that all isolates from the winter trial ( $n = 48$ ) harbored a resistance gene belonging to the *bla*<sub>SHV</sub> gene family. The resistance gene was sequenced in 8 isolates, identifying it as *bla*<sub>SHV-12</sub> in all chosen isolates. In the summer trial, all isolates ( $n = 54$ ) harbored a resistance gene belonging to the *bla*<sub>CTX-M</sub> gene family. All 8 *bla*<sub>CTX-M</sub> genes sequenced were identified as *bla*<sub>CTX-M-1</sub>.

One winter trial isolate and 4 summer trial isolates showed an additional *bla*<sub>TEM</sub> gene. All 5 *bla*<sub>TEM</sub> genes were identified as broad spectrum beta-lactamase resistance gene *bla*<sub>TEM-1</sub>.

For all genome-sequenced isolates from the winter trial ( $n = 19$ ), genome sequencing confirmed the presence of *bla*<sub>SHV-12</sub>. In addition, genome sequencing detected the plasmid-mediated quinolone-resistance



**Figure 4.** Results of the quantitative and qualitative analysis of total *E. coli* in cfu/g of litter, and the quantitative analysis of enterococci in cfu/g of litter for the winter trial samples (A) and the summer trial samples (B). The geometric mean of 3 samples is shown for each point in time. The error bars indicate the upper and lower 95% confidence intervals. The graphs were shifted to improve the visibility of the error bars.

**Table 3.** Phylogroups determined with gel-based PCR and phylogroups and sequence types determined via Enterobase.

Sample ID	Trial	Timepoint (h)	Sampling site	Phylogroup FU <sup>1</sup>	Phylogroup Enterobase	ST 7 gene MLST <sup>2</sup>	ST cGMLST <sup>3</sup>	HC1100 cgST <sup>4</sup>	H-Antigen	O-Antigen
7-1EP06	Summer	0 h	Barn	D/E	D	2,309	86,589	5,033	H6	O15
7-1EP05	Summer	0 h	Barn	D/E	D	2,309	86,626	5,033	H6	O15
7-1MP03	Summer	0 h	Surface	D/E	D	2,309	86,616	5,033	H6	O15
7-1MP05	Summer	1 h	Depth	D/E	D	2,309	86,618	5,033	H6	O15
7-1MP11	Summer	3 h	Depth	B1	B1	162	86,592	138	H10	O9
7-1MP10	Summer	3 h	Surface	D/E	D	2,309	86,619	5,033	H6	O15
7-1MP16	Summer	6 h	Surface	B1	B1	1,304	86,593	152	H7	O91
7-1MP17	Summer	6 h	Depth	D/E	D	2,309	86,608	5,033	H6	O15
7-1MP23	Summer	12 h	Depth	B1	D	2,309	86,590	5,033	H6	O15
7-1MP22	Summer	12 h	Surface	B1	B1	162	86,622	138	H10	O88
7-1MP31	Summer	24 h	Depth	D/E	D	2,309	86,580	5,033	H6	O15
7-1MP39	Summer	36 h	Depth	D/E	D	2,309	86,577	5,033	H6	O15
7-1MP34	Summer	36 h	Surface	B1	B1	1,304	86,593	152	H7	O91
7-1MP38	Summer	36 h	Surface	D/E	D	2,309	86,614	5,033	H6	O15
7-1MP35	Summer	36 h	Depth	D/E	D	2,309	86,620	5,033	H6	O15
7-1MP37	Summer	36 h	Depth	D/E	D	2,309	86,617	5,033	H6	O15
7-1MP41	Summer	48 h	Depth	D/E	D	2,309	86,580	5,033	H6	O15
7-1MP40	Summer	48 h	Surface	D/E	D	2,309	86,601	5,033	H6	O15
7-1MP42	Summer	48 h	Surface	D/E	D	2,309	86,623	5,033	H6	O15
7-1MP43	Summer	48 h	Depth	D/E	D	2,309	86,623	5,033	H6	O15
7-1MP44	Summer	48 h	Surface	D/E	D	2,309	86,580	5,033	H6	O15
7-1MP46	Summer	72 h	Surface	B1	B1	162	86,615	138	H10	O88
7-1MP48	Summer	72 h	Surface	D/E	D	2,309	86,625	5,033	H6	O15
7-1MP56	Summer	96 h	Surface	B1	B1	1,304	86,593	152	H7	O91
7-1MP54	Summer	96 h	Surface	D/E	D	2,309	86,624	5,033	H6	O15
5-2EP02	Winter	0 h	Barn	F	F	117	86,591	50	H4	O8
5-2EP01	Winter	0 h	Barn	F	F	117	86,578	50	H4	O8
5-2MP32	Winter	24 h	Surface	A/C	A	10	86,613	13	H48	O12
5-2MP33	Winter	24 h	Depth	F	F	117	86,579	50	H4	O8
5-2MP35	Winter	36 h	Depth	F	F	117	86,599	50	H4	O8
5-2MP34	Winter	36 h	Surface	F	F	117	86,653	50	H4	O8
5-2MP41	Winter	48 h	Depth	F	F	117	86,647	50	H4	O8
5-2MP43	Winter	48 h	Depth	F	F	117	86,600	50	H4	O8
5-2MP44	Winter	48 h	Surface	F	F	117	86,621	50	H4	O8
5-2MP42	Winter	48 h	Surface	F	F	117	86,632	50	H4	O8
5-2MP49	Winter	72 h	Depth	F	F	117	86,578	50	H4	O8
5-2MP50	Winter	72 h	Surface	F	F	117	86,578	50	H4	O8
5-2MP48	Winter	72 h	Surface	F	F	117	86,578	50	H4	O8
5-2MP46	Winter	72 h	Surface	F	F	117	86,578	50	H4	O8
5-2MP55	Winter	96 h	Depth	F	F	117	86,578	50	H4	O8
5-2MP54	Winter	96 h	Surface	F	F	117	86,627	50	H4	O8
5-2MP57	Winter	96 h	Depth	F	F	117	86,579	50	H4	O8
5-2MP56	Winter	96 h	Surface	F	F	117	86,578	50	H4	O8
5-2MP52	Winter	96 h	Surface	F	F	117	86,612	50	H4	O8

<sup>1</sup>Freie Universität Berlin.<sup>2</sup>Sequence type 7 gene multilocus sequence type.<sup>3</sup>Sequence type core genome multilocus sequence type.<sup>4</sup>Hierarchical cluster 1,100 core genome sequence type.

determinant *qnrS1* and *mdfA* genes in all 19 isolates, and the broad-spectrum beta-lactamase resistance gene *bla*<sub>TEM-1</sub> and the oxytetracycline resistance determinant Tet 34 in one isolate.

In the summer trial, the ESBL-resistance gene *bla*<sub>CTX-M-1</sub> was confirmed or found in all sequenced isolates ( $n = 25$ ). In addition, the *mdfA* gene was detected in all isolates, and Tet 34 was detected in 84% ( $n = 21/25$ ) of the isolates. *bla*<sub>TEM-1</sub> was detected in 4 isolates and the *sulIII* gene in 3 isolates.

### Genome-Based Phylogeny

Phylogenetic trees were calculated in Enterobase and can be found in the [Supplemental Figures 1 and 2](#).

While there are genomic differences between isolates from the summer and winter experiments, isolates within each experiment showed little variation with

the summer experiment appearing to be slightly more diverse.

Within the 2 main clusters (one for each season), the different sampling sites are distributed equally as well as the time points for sampling. The genetic differences between the 2 experiments is also reflected in the distribution of resistance genes.

## DISCUSSION

The aim of our study was to evaluate whether short-term litter storage provides sufficient inactivation of ESBL-producing *E. coli* present in chicken litter under field conditions in winter and summer.

The most important findings of the study concerning the amount of cultivable ESBL-producing *E. coli* were that regardless of the season, the inactivation occurs faster in a depth of about 50 cm compared with the

surface of the litter pile. Additionally, a seasonal influence concerning the required time span and the effectiveness of the inactivation of ESBL-producing *E. coli* was shown. Extended-spectrum beta-lactamases-producing *E. coli* were not detected quantitatively in the deep samples in the winter trial after 72 h and already after 36 h in the summer trial.

Initial concentrations of nonresistant *E. coli* in the chicken litter were significantly higher than ESBL-producing *E. coli* for both trials. The proportion of the ESBL-producing subpopulation on the total amount of *E. coli* was 0.94% in the winter trial and 1.64% in the summer trial for the samples taken at storage begin. A similar proportion of 1.1% ESBL-producing isolates was recently reported by Friese et al. (2019) for turkey-rearing flocks. This is a possible explanation for the extended time in which nonresistant *E. coli* were detected in the litter compared with the ESBL-producing subpopulation. In the study performed by Erickson et al. (2010), nonresistant *E. coli* naturally occurring in the litter were not detected quantitatively and qualitatively in surface and deep samples after 4 D in static piles of chicken litter in summer, fall, and winter.

### **Abiotic Factors Influencing Microbial Counts in the Litter Storage Trials**

Litter piles are microbiologically highly heterogeneous and local conditions are influenced by a variety of biological, physical, and chemical factors. In our study, we explored the influence of temperature, pH value, and moisture content of the chicken litter on the bacterial cell count.

In the study performed by Erickson et al. (2010) in the USA, temperatures within static piles of chicken litter were measured at different intervals. The highest mean temperatures reported in a depth of 30 cm from the heaps surface were 54.4°C in the summer after 4 D of storage and 51.8°C in the winter after 3 D of storage. This is in accordance with our findings where a maximum temperature of 58.5°C was measured in the summer and 50.4°C in the winter. Temperatures over 65°C were reported by Wilkinson et al. (2011) for static piles of poultry litter in the first weeks of aging.

In a recently performed laboratory scale anaerobic digestion experiment by Thomas et al. (2019), ESBL-/AmpC-producing *E. coli* were added in a concentration of over  $10^7$  cfu/ml to a mix of chicken litter and an inoculum from a biogas plant. They showed that at a constant temperature of 55°C, ESBL-producing *E. coli* were quantitatively undetectable by direct count after 2 h of incubation. In our summer trial, temperatures reached levels constantly above 53°C at 48 h. In the subsequent samples (72 h and 96 h), we did not detect ESBL-producing *E. coli* and nonresistant *E. coli* quantitatively and qualitatively. We therefore assume that a temperature of 53°C is sufficient for inactivation of all ESBL-producing and nonresistant *E. coli* under practical conditions of an anaerobic litter storage.

A laboratory scale study by Laport et al. (2003) showed that a 2 h incubation period at 55°C will result in >90% reduction of *Enterococcus faecium*. Accordingly, at the end of our summer trial enterococci concentrations had decreased by > 99% compared with the initial concentrations in the deep samples.

In the winter trial, a temperature of over 50°C was observed in the deep samples for the last point in time after 96 h only. Extended-spectrum beta-lactamases-producing *E. coli* were qualitative, and nonresistant *E. coli* were quantitatively detectable until the end of the trial. The number of enterococci was stable in both the surface and deep samples in the winter trial, ranging from  $3.2 \times 10^6$  to  $5.6 \times 10^7$  cfu/g in all samples. Therefore, we assume that the increase in temperature to a maximum of 50.4°C in the winter trial was insufficient for a distinct reduction of the monitored bacteria in chicken litter.

A further factor, which might influence the bacterial counts in the chicken litter, is the pH value. The pH value of the litter at the beginning of the experiment (0 h) was in the alkaline range, a frequent finding in chicken litter (Huang et al., 2017). The pH value drop in the depth of the litter heap was presumably because of anaerobic fermentation and formation of organic acids like propionic acid, butyric acid, and acetic acid (Cornell Waste Management Institute, 1996). The minimum pH value of 5.6 measured in the winter trial and 6.5 in the summer trial is not sufficient to inactivate *E. coli*. It was shown that *E. coli* has a high probability of surviving pH values of 1.5 to 4.0 (Takumi et al., 2000).

### **Insights From Whole Genome Sequencing**

Whole genome sequencing revealed that the analyzed isolates for both trials harbored additional resistance genes besides ESBL resistance genes. Identified genes included *qnrS1*, which may mediate resistance to quinolones (Cerquetti et al., 2009), *mdfA*, which substantially increases resistance to amphoteric lipophilic compounds (e.g., ethidium bromide, benzalkonium, and tetracycline) (Edgar and Bibi, 1997) and the Tet 34 and *sulIII* resistance genes, which mediate for resistance against oxytetracycline (Nonaka and Suzuki, 2002) and sulfonamides (Radstrom and Swedberg, 1988).

The results also indicate that one strain of ESBL-producing *E. coli* was predominant in the chicken barns for each of the trials. This finding is especially prominent in the winter trial, where 94.7% (n = 18/19) of the isolates belonged to the same 7 gene MLST (ST117) and HC1100cgST (ST50).

Results of the phylogenetic analyses suggest that genetic differences do not equip the isolates of a certain cluster with a survival advantage.

### **Advantages and Limitations of the Study**

In our study, we did not artificially add ESBL-producing *E. coli* to the litter. Instead, chicken litter naturally contaminated with these resistant bacteria



was used, mirroring true field conditions. [Hutchison et al. \(2005\)](#) pointed out that this is crucial because bacteria are already adapted to their environment, and therefore, bacterial stress is minimized ([Wesche et al., 2009](#)). A further advantage over lab-scale studies is that the litter heaps were exposed to environmental conditions which are typical for the winter and summer season in central Europe. For on-farm waste management, similar environmental conditions are likely to appear; therefore, better transferability of the results can be expected when compared with laboratory studies.

A limitation of the study is that the measurements of temperature, humidity, and the litter samples taken for the microbiological analyses cannot cover the conditions in the entire storage mass. It was shown previously that because of increased deposition of fecal droppings on the litter surface, the number of coliform bacteria is significantly higher in the top layer of a chicken litter bed compared with the bottom layers ([Barker et al., 2010](#)). Using a front-end loader to remove and stack the litter at the end of the fattening period will not assure complete mixing of the litter, resulting in an inhomogeneous distribution of coliform bacteria. This might explain inconsistencies we saw in the course of the bacterial counts for some points in time. However, through our sampling scheme that covered both surface and deep samples with 3 samples evenly distributed over the litter heap for each point in time, we achieved representative results.

### **Evaluation of Short-Term Litter Storage as an On-Farm Strategy to Prevent the Spread of ESBL-Producing *E. coli* to the Environment**

It was shown by [Merchant et al. \(2012\)](#) that resistant *E. coli* were detectable in soil fertilized with chicken litter for at least 7 mo. In that study performed in Canada of 295 *E. coli* isolated from soil, 139 carried either a *bla*<sub>SHV</sub>, *bla*<sub>TEM</sub>, or *bla*<sub>CMY-2</sub> resistance gene. This highlights the importance of sufficient inactivation of resistant *E. coli* in litter before land application.

Very low quantities of resistant microorganisms might be able to horizontally transfer mobile genetic elements to microorganisms in the environment, thus potentially contributing to a spread of antibiotic resistance. As a result, the qualitative detection of ESBL-producing *E. coli* in litter is of particular importance. A recent study by [Pornasukarom and Thakur \(2017\)](#) demonstrated that the application of manure containing Enterobacteriaceae which carry plasmids mediating for antibiotic resistance enriches the environmental resistome. Our results indicate that a storage period of 5 D is sufficient to reduce the amount of ESBL-producing *E. coli* in the depth of a chicken litter heap in the summer below the detection limit. For very low ambient temperatures, as present in our winter trial, an extension of the storage period should be considered because we observed an incomplete inactivation of ESBL-producing *E. coli* in a 5-day storage period.

Even if cultivation-based methods are unable to detect resistant bacteria, a transfer of plasmids carrying resistance genes might occur. In a study performed by [Le Devendec et al. \(2016\)](#), chicken manure was stored for 6 wk. After this time span, *E. coli* were not detected in the manure by cultivation without enrichment. Plasmid capture assays with the stored chicken manure revealed an uptake of plasmids encoding resistance to sulfonamides, aminoglycosides, and streptomycin in recipient strains. This indicates that even if there are no cultivable bacteria left in the litter, the possibility of spread of resistance cannot be discounted because viable plasmids could still be present. In contrast, [Guan et al. \(2007\)](#) stated that composting of chicken manure at high temperatures could help prevent the spread of antibiotic-resistant genes via plasmids in the environment. In their study, neither viable *E. coli* nor their plasmids could be detected in compost microcosms, which reached temperatures of over 50°C.

The outer edges of litter piles may present a reservoir for bacteria, and turning the litter pile may therefore lead to a recontamination of the interior parts ([Pereira-Neto et al., 1986](#)). This is in accordance with our observation in the summer trial, where the number of *E. coli* and enterococci significantly increased on the surface of the litter heap.

The increased quantity of these bacteria over the course of the trial may be caused by beneficial environmental factors such as rainfall, which influences moisture levels in litter piles and can promote regrowth of enteric bacteria ([Gibbs et al., 1997](#)). In the summer trial of our study, rainfall at the end of the first trial day led to increased moisture in the litter heap. Corresponding temperatures on the surface of the litter pile ranged from 29.2°C to 42.7°C, which are known to be sufficiently high for bacterial regrowth ([Kumar and Libchaber, 2013](#)).

The survival time of *E. coli* in manure is significantly longer under anaerobic than under aerobic conditions ([Semenov et al., 2011](#)). Additionally, it was shown in the trial by [Wilkinson et al. \(2011\)](#) that temperatures in composted chicken litter piles are higher than in stored piles. As previous research and this study indicate a faster inactivation of ESBL-producing *E. coli* can be achieved at higher temperatures. It appears therefore that composting litter under aerobic conditions could lead to a faster inactivation of ESBL-producing *E. coli* compared with storing it in anaerobic conditions. However, increased working and litter-storage capacities are required and not available on all farms.

In conclusion, short-term litter storage is a useful, easily realizable tool leading to an effective reduction of the amount of ESBL-producing *E. coli* in chicken litter. However, we did not observe a complete inactivation of ESBL-producing *E. coli* in the depth of the heap in the winter and on the surface of the heap for both trials. An extension of the storage period for low ambient temperatures and stirring the pile one time or composting the litter instead of storing it could increase the effectiveness of chicken litter hygienization.

## ACKNOWLEDGMENTS

The authors would like to thank the staff of the broiler farm for the excellent collaboration. They thank Kerstin Rosen and Benjamin Reichelt (Institute for Animal Hygiene and Environmental Health, Berlin, Germany) for assistance with taking and processing the samples. They also thank Heike Rose, Susann Sellenthin, Sabrina Hansen, Michael Kühl (Institute for Animal Hygiene and Environmental Health, Berlin, Germany), and Vera Junker (DSMZ, Braunschweig, Germany) for excellent technical advice and assistance. They thank Alexander Bartel (Institute for Veterinary Epidemiology and Biostatistics, Berlin, Germany) for statistical advice and Samira Schlesinger (Clinic of Animal Reproduction, Berlin, Germany) for proofreading the manuscript.

The study was funded by the Leibniz Association (grant number: SAW-2017-DSMZ-2)

## SUPPLEMENTARY DATA

Supplementary data associated with this article can be found in the online version at <https://doi.org/10.1016/j.psj.2019.11.043>.

## REFERENCES

- Barker, K. J., J. L. Purswell, J. D. Davis, H. M. Parker, M. T. Kidd, C. D. McDaniel, and A. S. Kiess. 2010. Distribution of bacteria at different poultry litter depths. *Int. J. Poultry Sci.* 9:10–13.
- Baym, M., S. Kryazhimskiy, T. D. Lieberman, H. Chung, M. M. Desai, and R. Kishony. 2015. Inexpensive Multiplexed library Preparation for Megabase-Sized genomes. *PLoS One* 10:1–15.
- Beghain, J., A. Bridier-Nahmias, H. Le Nagard, E. Denamur, and O. Clermont. 2018. ClermontTyping: an easy-to-use and accurate in silico method for *Escherichia* genus strain phylotyping. *Microb. Genomics* 4:1–8.
- Blaak, H., A. H. A. M. Van Hoek, R. A. Hamidjaja, R. Q. J. Van Der Plaats, L. Kerkhof-De Heer, A. M. De Roda Husman, and F. M. Schets. 2015. Distribution, numbers, and diversity of ESBL-producing *E. coli* in the poultry farm environment. *PLoS One* 10:1–23.
- Bland, J. M., and D. G. Altman. 1996. Statistics notes: Transformations, means, and confidence intervals. *BMJ* 312:1079.
- Cerquetti, M., A. García-ferna, M. Giufre, D. Fortini, M. Accogli, C. Graziani, I. Luzzi, A. Caprioli, and A. Carattoli. 2009. First Report of plasmid-mediated quinolone resistance determinant qnrS1 in an *Escherichia coli* strain of animal Origin in Italy. *Antimicrob. Agents Chemother.* 53:3112–3114.
- Clermont, O., J. K. Christenson, E. Denamur, and D. M. Gordon. 2013. The Clermont *Escherichia coli* phylo-typing method revisited: Improvement of specificity and detection of new phylo-groups. *Environ. Microbiol. Rep.* 5:58–65.
- Cornell Waste Management Institute, C. U. 1996. [science @ compost.css.cornell.edu](http://compost.css.cornell.edu/science.html). Accessed Feb. 2020. <http://compost.css.cornell.edu/science.html>.
- Dahre, K., M. Projahn, T. Semmler, U. Roesler, and A. Friese. 2018. Extended-spectrum beta-lactamase-/AmpC beta-lactamase-producing enterobacteriaceae in broiler farms: transmission dynamics at farm level. *Microb. Drug Resist* 24:511–518.
- Le Devendec, L., G. Mourand, S. Bougeard, J. Léaustic, E. Jouy, A. Keita, W. Couet, N. Rousset, and I. Kempf. 2016. Impact of colistin sulfate treatment of broilers on the presence of resistant bacteria and resistance genes in stored or composted manure. *Vet. Microbiol.* 194:98–106.
- Dierikx, C., A. van Essen-Zandbergen, K. Veldman, H. Smith, and D. Mevius. 2010. Increased detection of extended spectrum beta-lactamase producing *Salmonella enterica* and *Escherichia coli* isolates from poultry. *Vet. Microbiol.* 145:273–278.
- Dierikx, C., J. van der Goot, T. Fabri, A. van Essen-Zandbergen, H. Smith, and D. Mevius. 2013. Extended-spectrum- $\beta$ -lactamase- and AmpC- $\beta$ -lactamase-producing *Escherichia coli* in Dutch broilers and broiler farmers. *J. Antimicrob. Chemother.* 68:60–67.
- Edgar, R., and E. Bibi. 1997. MdfA, an *Escherichia coli* Multidrug resistance Protein with an Extraordinarily broad spectrum of drug Recognition. *J. Bacteriol.* 179:2274–2280.
- EFSA. 2011. Scientific Opinion on the public health risks of bacterial strains producing extended-spectrum  $\beta$ -lactamases and/or AmpC  $\beta$ -lactamases in food and food producing strains. *EFSA J.* 9:1–95.
- Erickson, M. C., J. Liao, G. Boyhan, C. Smith, L. Ma, X. Jiang, and M. P. Doyle. 2010. Fate of manure-borne pathogen surrogates in static composting piles of chicken litter and peanut hulls. *Bioresour. Technol.* 101:1014–1020.
- Feldgarden, M., B. Vyacheslav, D. H. Hafta, A. B. Prasada, D. J. Slotta, I. Tolstoya, G. H. Tysonb, S. Zhaob, C.-H. Hsub, P. F. McDermottb, D. A. Tadesseb, C. Morales, M. Simmons, G. Tillman, J. Wasilenko, J. P. Folsterd, and W. Klimke. 2019. Using the NCBI AMRFinder tool to determine antimicrobial resistance genotype-phenotype correlations within a collection of NARMS isolates. Accessed Feb. 2020. <https://www.biorxiv.org/content/10.1101/550707v1.article-info>.
- Friese, A., H. M. Lu, R. Merle, and U. Roesler. 2019. Extended-spectrum beta-lactamase and AmpC beta-lactamase-producing Enterobacteriaceae in Turkey fattening farms: a cross-sectional study. *Berl. Munch. Tierarztl. Wochenschr.* 132:352–359.
- Gibbs, R. A., C. J. Hu, G. Ho, and I. Unkovich. 1997. Regrowth of faecal coliforms and salmonellae in stored biosolids and soil amended with biosolids. *Water Sci. Technol.* 35:269–275.
- Guan, J., A. Wasty, C. Grenier, and M. Chan. 2007. Influence of temperature on survival and conjugative transfer of multiple antibiotic-resistant plasmids in chicken manure and compost microcosms. *Poult. Sci.* 86:610–613.
- Hartmann, A., A. Locatelli, L. Amoureux, G. Depret, C. Jolivet, E. Gueneau, and C. Neuwirth. 2012. Occurrence of CTX-M producing *Escherichia coli* in soils, Cattle, and farm environment in France (Burgundy region). *Front. Microbiol.* 3:83.
- Hering, J., C. Frömke, C. von Münchhausen, M. Hartmann, B. Schneider, A. Friese, U. Rösler, L. Kreienbrock, and K. Hille. 2016. Cefotaxime-resistant *Escherichia coli* in broiler farms-A cross-sectional investigation in Germany. *Prev. Vet. Med.* 125:154–157.
- Huang, J., Z. Yu, H. Gao, X. Yan, J. Chang, C. Wang, J. Hu, and L. Zhang. 2017. Chemical structures and characteristics of animal manures and composts during composting and assessment of maturity indices. *PLoS One* 12:1–16.
- Huijbers, P. M. C., E. A. M. Graat, A. P. J. Haenen, M. G. van Santen, A. van Essen-Zandbergen, D. J. Mevius, E. van Duijkeren, and A. H. A. M. van Hoek. 2014. Extended-spectrum and AmpC  $\beta$ -lactamase-producing *Escherichia coli* in broilers and people living and/or working on broiler farms: prevalence, risk factors and molecular characteristics. *J. Antimicrob. Chemother.* 69:2669–2675.
- Hutchison, M. L., L. D. Walters, S. M. Avery, and A. Moore. 2005. Decline of zoonotic agents in livestock waste and bedding heaps. *J. Appl. Microbiol.* 99:354–362.
- Jia, B., A. R. Raphenya, B. Alcock, N. Waglehner, P. Guo, K. K. Tsang, B. A. Lago, B. M. Dave, S. Pereira, A. N. Sharma, S. Doshi, M. Courtot, R. Lo, L. E. Williams, J. G. Frye, T. Elsayegh, D. Sardar, E. L. Westman, A. C. Pawlowski, T. A. Johnson, F. S. L. Brinkman, G. D. Wright, and A. G. McArthur. 2017. Card 2017: Expansion and model-centric curation of the comprehensive antibiotic resistance database. *Nucleic Acids Res.* 45:D566–D573.
- Kumar, P., and A. Libchaber. 2013. Pressure and temperature Dependence of Growth and Morphology of *Escherichia coli*: experiments and Stochastic model. *Biophys. J.* 105:783–793.
- Laport, M. S., M. Ramos, C. C. Silva, C. De Freire, and M. Giambiagi-deMarval. 2003. Heat-resistance and Heat-Shock Response in the Nosocomial pathogen *Enterococcus faecium*. *Curr. Microbiol.* 46:313–317.
- Laube, H., A. Friese, C. von Salviati, B. Guerra, A. Käsbohrer, L. Kreienbrock, and U. Roesler. 2013. Longitudinal monitoring of extended-spectrum-beta-lactamase/ampC-producing *Escherichia*

- coli at German broiler chicken fattening farms. *Appl. Environ. Microbiol.* 79:4815–4820.
- Laube, H., A. Friese, C. von Salviati, B. Guerra, and U. Rösler. 2014. Transmission of ESBL/AmpC-producing *Escherichia coli* from broiler chicken farms to surrounding areas. *Vet. Microbiol.* 172:519–527.
- Leverstein-van Hall, M. A., C. M. Dierikx, J. C. Stuart, G. M. Voets, M. P. van den Munckhof, A. van Essen-Zandbergen, T. Platteel, A. C. Fluit, N. van de Sande-Bruinsma, J. Scharinga, M. J. M. Bonten, and D. J. Mevius. 2011. Dutch patients, retail chicken meat and poultry share the same ESBL genes, plasmids and strains. *Clin. Microbiol. Infect.* 17:873–880.
- Merchant, L. E., H. Rempel, T. Forge, T. Kannangara, S. Bittman, P. Delaquis, E. Topp, K. A. Ziebell, and Moussa S. Diarra. 2012. Characterization of antibiotic-resistant and potentially pathogenic *Escherichia coli* from soil fertilized with litter of broiler chickens fed antimicrobial-supplemented diets. *Can. J. Microbiol.* 58:1084–1098.
- Nonaka, L., and S. Suzuki. 2002. New Mg<sup>2+</sup>-Dependent oxytetracycline resistance determinant Tet 34 in *Vibrio* isolates from Marine fish Intestinal contents. *Antimicrob. Agents Chemother.* 46:1550–1552.
- Pereira-Neto, J., E. I. Stentiford, and D. V. Smith. 1986. Survival of faecal indicator micro-organisms in refuse/sludge composting using the aerated static pile system. *Waste Manag. Res.* 4:397–406.
- Pornsukarom, S., and S. Thakur. 2017. Cross horizontal dissemination of antimicrobial resistance determinants in multiple *Salmonella* Serotypes following isolation from the Commercial swine Operation environment after manure. *Appl. Environ. Microbiol.* 83:1–14.
- Projahn, M., K. Daehre, U. Roesler, and A. Friese. 2017. Extended-spectrum-beta-lactamase and plasmid-encoded cephalosporinase-producing enterobacteria in the broiler hatchery as a potential mode of pseudo-vertical transmission. *Appl. Environ. Microbiol.* 83.
- Radstrom, P., and G. Swedberg. 1988. RSF1010 and a conjugative plasmid Contain sulII, one of two known genes for plasmid-borne sulfonamide resistance Dihydropteroate Synthase. *Antimicrob. Agents Chemother.* 32:1684–1692.
- Remschmidt, C., S. Schneider, E. Meyer, B. Schroeren-Boersch, P. Gastmeier, and F. Schwab. 2017. Surveillance of antibiotic use and resistance in intensive care units (SARI) a 15-year cohort study. *Dtsch. Arztebl. Int.* 114:858–865.
- Roschanski, N., J. Fischer, B. Guerra, and U. Roesler. 2014. Development of a multiplex real-time PCR for the rapid detection of the predominant beta-lactamase genes CTX-M, SHV, TEM and CIT-type AmpCs in enterobacteriaceae. *PLoS One* 9:e100956.
- Semenov, A. V., L. Van Overbeek, A. J. Termorshuizen, and A. H. C. Van Bruggen. 2011. Influence of aerobic and anaerobic conditions on survival of *Escherichia coli* O157:H7 and *Salmonella enterica* serovar Typhimurium in Luria-Bertani broth, farm-yard manure and slurry. *J. Environ. Manage.* 92:780–787.
- Statistisches Bundesamt. 2016. Wirtschaftsdünger Tierischer Herkunft in Landwirtschaftlichen Betrieben/Agrarstrukturerhebung. Fachserie 3 R. 2.2.2 2030222169, pp. 2–74.
- Statistisches Bundesamt. 2017. Bundesamt (Destatis), 2017. Land- und Forstwirtschaft, Fischerei – Wirtschaftsdünger tierischer Herkunft in landwirtschaftlichen Betrieben/Agrarstrukturerhebung. Accessed Feb. 2020. <https://www.destatis.de/DE/Themen/Branchen-Unternehmen/Landwirtschaft-Forstwirtschaft-Fischerei/Produktionsmethoden/Publikationen/Downloads-Produktionsmethoden/wirtschaftsduenger-2030222169005.html>.
- Steglich, M., J. D. Hofmann, J. Helmecke, J. Sikorski, C. Spröer, T. Riedel, B. Bunk, and J. Overmann. 2018. Convergent Loss of ABC transporter genes from *Clostridioides difficile* genomes is Associated with Impaired Tyrosine uptake and p-Cresol production. *Front. Microbiol.* 9:1–12.
- Takumi, K., R. De Jonge, and A. Havelaar. 2000. Modelling inactivation of *Escherichia coli* by low pH: application to passage through the stomach of young and elderly people. *J. Appl. Microbiol.* 89:935–943.
- Thomas, C., C. Idler, C. Ammon, C. Herrmann, and T. Amon. 2019. Inactivation of ESBL-/AmpC-producing *Escherichia coli* during mesophilic and thermophilic anaerobic digestion of chicken manure. *Waste Manag.* 84:74–82.
- Waters, N., F. Brennan, A. Holmes, F. Abram, and L. Pritchard. 2018. Easily Phylotyping *E. coli* via the EzClermont Web App and Command-Line Tool. Accessed Feb. 2020. <https://www.biorxiv.org/content/10.1101/317610v1.article-info>.
- Wesche, A. M., J. B. Gurtler, B. P. Marks, and E. T. Ryser. 2009. Stress, Sublethal Injury, Resuscitation, and Virulence of bacterial Foodborne Pathogenst. *J. Food Prot.* 72:1121–1138.
- Wilkinson, K. G., E. Tee, R. B. Tomkins, G. Hepworth, and R. Premier. 2011. Effect of heating and aging of poultry litter on the persistence of enteric bacteria. *Poult. Sci.* 90:10–18.
- Zankari, E., H. Hasman, S. Cosentino, M. Vestergaard, S. Rasmussen, O. Lund, F. M. Aarestrup, and M. V. Larsen. 2012. Identification of acquired antimicrobial resistance genes. *J. Antimicrob. Chemother.* 67:2640–2644.
- Zhou, Z., N. Alikhan, K. Mohamed, S. Group, M. Achtman, G. H. Road, U. Kingdom, M. Chattaway, R. Delahay, S. Hutton, H. Graz, L. Petrovska, P. H. Agency, W. Tyne, Y. Fan, N. Williams, and P. Health. 2019. The User’s Guide to Comparative Genomics with Enterobase. Three Case Studies: Micro-clades within *Salmonella enterica* Serovar Agama, Ancient and Modern Populations of *Yersinia pestis*, and Core Genomic Diversity of All *Escherichia*. Accessed Feb. 2020. <https://www.biorxiv.org/content/10.1101/613554v1.article-info>.