

Journal of Molecular and Genetic Medicine

Minamizaki et al., J Mol Genet Med 2016, 10:4 DOI: 10.4172/1747-0862.1000231

Commentary OMICS Internationa

Nano-Architecture of Osteocyte Lacunae

Minamizaki T, Ahmed F and Yoshiko Y*

Department of Calcified Tissue Biology, Oral Health Science Major, Hiroshima University Institute of Biomedical and Health Sciences, 1-2-3 Kasumi, Minami-ku, Hiroshima 7348553, Japan

Introduction

During bone formation, 20 percent of bone-forming osteoblasts can be trapped in the newly formed bone matrix and become osteocytes. Osteocytes are most commonly found in bone, while their differentiation processes are largely unknown. Osteocytes reside inside spaces in the bone matrix "lacunae," and extend their cell processes through tiny canals called canaliculi, resulting in the formation of an extensive cellular network throughout the bone matrix [1]. Although osteocytes appear to be morphologically inert, recent findings indicate that the cells act as master regulators of bone and mineral homeostasis through the mechanisms underlying the mechanosensing system [2,3]. For example, sclerostin and FGF23 are known to be exclusively produced by osteocytes [4]. The former acts an inhibitor of the Wnt pathway and the resulting decrease in bone formation, and the latter regulates renal phosphate transport to maintain serum phosphate levels that impact bone mineralization. It is also well known that RANKL derived from osteocytes stimulates osteoclast formation and activate bone resorption [4].

Cross-sectional imaging including X-ray computed tomography (CT) and its 3D geometries and properties are used in the clinical routine with a maximum spatial resolution from hundred to dozen μm . The nano-architectures exceeding this resolution currently cannot be visualized [5]. Nano-CT provides high-resolution cross-sectional imaging and significantly exceeds the resolution capacity of established CT systems. Nano-CT technology can achieve a superior spatial resolution at hundred or less nm levels, and hence is anticipated to be utilized for analysis of the nano-architecture in various fields including medical sciences. In general, osteocytes and their lacunae are less than 30 μm in size (long axis length), and accordingly little attention had

Lacunar volume	(mm³)
Lacunar volume/bone volume	(%)
Osteocyte lacunar surface	(mm²)
Lacunar surface/lacunar volume ratio	(1/mm)
Lacunar surface/bone volume	(1/mm)
Lacunar surface convexity index	(1/mm)
Lacunar diameter	(mm)
Lacunar linear density	(1/mm)
Lacunar separation	(mm)
Degree of anisotropy (non-random alignment of osteocyte lacunae)	Scalar value between zero and the degree
Fractal dimension (complexity of osteocyte lacunar surface)	D=ratio of the numbers
Lacunae number	n
Lacunar number/bone volume	(1/mm³)
Average lacunar length	(mm)
SD lacunar thickness	(mm)
SD lacunar separation	(mm)
Lacunar diameter distribution	(mm)
Lacunar separation distribution	(mm)
Eigenvalue (vertical direction, lateral direction, longitudinal direction)	Special set of scalars
(Provided by Brukar micro CT, Kontich, Belgium).	

 Table 1: Osteocyte lacunar parameters.




Figure 1: Nano-CT images of osteocyte lacunae in the mouse femur. The femur from the 6-week-old *kl/+* male mouse was used for analysis. Nano-CT scan was performed using a high-resolution nano-CT system (voxel size, 600 nm; Sky Scan 2211, SkyScan NV, Kontich, Belgium). The energy of the beam was set to 45 kV, and an aluminum filter (0.5 mm) was used. After standardized reconstruction by using NRecon software (Brukar micro CT, Kontich, Belgium), 3D modeling of the samples was carried out by using CTvox software (Brukar) and the data sets for each bone section were analyzed using CTan software (Brukar). Nano-CT analysis was performed on 401 slices of 600 nm thickness in the femoral diaphysis. (a) A reconstructed slice of the overview scan. (b-e) 3D images of osteocyte lacunae. Two different view directions at low magnification (b and c), and higher magnification views of b (d and e). Light blue, cortical bone; Brown, osteocyte lacunae.

been paid to the contribution of conventional CT systems to assessing osteocyte properties.

Osteocytes change activities or die, accompanied by their morphologies, throughout life as well as in pathological situations [2-4]. Functional and morphological anomalies of osteocytes are reflected in osteocyte lacunar density, volume, shape, and anisotropy which was evaluated by Syncrotron Radiation micro-CT and nano-CT [6-8].

*Corresponding authors: Dr. Yuji Yoshiko, Department of Calcified Tissue Biology, Oral Health Science Major, Hiroshima University Institute of Biomedical and Health Sciences, 1-2-3 Kasumi, Minami-ku, Hiroshima 7348553, Japan, Tel: 81 82 257 5620, Fax: 81 82 257 5621; E-mail: yyuji@hiroshima-u.ac.jp

Received November 03, 2016; Accepted November 17, 2016; Published November 21, 2016

Citation: Minamizaki T, Ahmed F, Yoshiko Y (2016) Nano-Architecture of Osteocyte Lacunae. J Mol Genet Med 10: 231 doi:10.4172/1747-0862.1000231

Copyright: © 2016 Minamizaki T, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Thus, criteria for osteocyte lacunae may be beneficial for the diagnosis and management of bone development and health. Currently, multiple parameters are available to define the nano-architectures of osteocyte lacunae by nano-CT (Table 1).

Three-dimensional rendering of nano-CT images shows multiple types of osteocyte lacunae in the mouse cortical bone (Figure 1), suggesting the remarkable ability of nano-CT. The total number of osteocytes in the adult skeleton is reported to be ~ 42 billion. As for osteocyte dendritic projections (cell networks), the total number reaches 100 times more than that of osteocytes [1]. These data allow us to better understand osteocyte properties by evaluating as many as possible of osteocyte lacunae in the site of interest. In terms of covering a broad range of osteocyte lacunae with high resolution, nano-CT exhibits its high performance.

References

1. Buenzli PR, Sims NA (2015) Quantifying the osteocyte network in the human skeleton. Bone 75: 144-150.

- 2. Prideaux M, Findlay DM, Atkins GJ (2016) Osteocytes: The master cells in bone remodelling. Curr Opin Pharmacol 28: 24-30.
- 3. Van Hove RP, Nolte PA, Vatsa A, Semeins CM, Salmon PL, et al. (2009) Osteocyte morphology in human tibiae of different bone pathologies with different bone mineral density-Is there a role for mechanosensing? Bone 45: 321-329.
- 4. Dallas SL, Prideaux M, Bonewald LF (2013) The osteocyte: an endocrine cell and more. Endocr Rev 34: 658-690.
- $\label{eq:lambda} \text{Kampschulte M, Langheinirch AC, Sender J, Litzlbauer HD, Althöhn U, et al.}$ (2016) Nano-computed tomography: Technique and applications. Rofo 188: 146-154.
- 6. Bach-Gansmo FL, Weaver JC, Jensen MH, Leemreize H, Mader KS, et al. (2015) Steocyte lacunar properties in rat cortical bone: Differences between lamellar and central bone. J Struct Biol 191: 59-67.
- 7. Matsuo K, Kuroda Y, Nango N, Shimoda K, Kubota Y, et al. (2015) Osteogenic capillaries orchestrate growth plate-independent ossification of the malleus. Development 142: 3912-3920.
- Peyrin F, Dong P, Pacureanu A, Langer M (2014) Micro and nano-CT for the study of bone ultrastructure. Curr Osteoporos Rep 12: 465-474.

OMICS International: Open Access Publication Benefits & **Features**

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50.000+ editorial team Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: http://www.omicsonline.org/submission//

Citation: Minamizaki T. Ahmed F. Yoshiko Y (2016) Nano-Architecture of Osteocyte Lacunae, J Mol Genet Med 10: 231 doi:10.4172/1747-0862.1000231