



저작자표시-비영리-변경금지 2.0 대한민국

이용자는 아래의 조건을 따르는 경우에 한하여 자유롭게

- 이 저작물을 복제, 배포, 전송, 전시, 공연 및 방송할 수 있습니다.


다음과 같은 조건을 따라야 합니다:



저작자표시. 귀하는 원저작자를 표시하여야 합니다.



비영리. 귀하는 이 저작물을 영리 목적으로 이용할 수 없습니다.




변경금지. 귀하는 이 저작물을 개작, 변형 또는 가공할 수 없습니다.

- 귀하는, 이 저작물의 재이용이나 배포의 경우, 이 저작물에 적용된 이용허락조건을 명확하게 나타내어야 합니다.
- 저작권자로부터 별도의 허가를 받으면 이러한 조건들은 적용되지 않습니다.

저작권법에 따른 이용자의 권리는 위의 내용에 의하여 영향을 받지 않습니다.

이것은 [이용허락규약\(Legal Code\)](#)을 이해하기 쉽게 요약한 것입니다.

[Disclaimer](#) 

A Thesis for the Degree of Master of Science in Pharmacy

**Isolation and Structure Determination of
Bioactive Constituents from *Poncirus trifoliata* Raf.**

지실의 성분연구

August 2018

by
Won-Hee Lee

Natural Products Science Major, College of Pharmacy
Master Course in the Graduate School
Seoul National University

Abstract

**Isolation and Structure Determination of
Bioactive Constituents from *Poncirus trifoliata* Raf.**

Won-Hee Lee
Natural Products Science Major
College of Pharmacy
Master Course in the Graduate School
Seoul National University

Sortase A (SrtA), a gram-positive transpeptidase, is an important target in reducing bacterial infection. The extract from the fruits of trifoliolate orange, *Poncirus trifoliata* Rafinesque, showed significant Srt A inhibition.

Poncirus trifoliata Raf. is a member of the family Rutaceae and it has been widely used in Korean traditional medicine for the treatment of diverse pathogenic symptoms.

Bioactivity-guided chromatographic separation afforded the isolation of in total twelve compounds, eight coumarins and four flavanone glycosides.

The structures of four novel coumarins were determined by a combination of spectroscopic analyses such as NMR and LC-MS.

The absolute configurations of the new coumarins and known

coumarins, previously unassigned, were determined by an application of the 2-methoxy-2-(trifluoromethyl)phenylacetic acid (MTPA) method.

Key words: herbal medicine, *Poncirus trifoliata* Rafinesque, coumarins, flavanone glycosides, sortase A inhibition

Student number: 2016-29626

List of Contents

Abstract in English	I
List of Contents	III
List of Scheme and Tables	V
List of Figures	VII
Introduction	1
Experimental Section	3
1. General Experimental Procedures	3
2. Plant Material	3
3. Extraction and Isolation	4
4. Preparation of the (<i>S</i>)- and (<i>R</i>)- α -Methoxy- α - trifluoromethylphenylacetic chloride (MTPA-Cl) Esterification of Compound 1-6	6
Results	12
1. Compound 1	12
2. Compound 2	12
3. Compound 3	13
4. Compound 4	13
5. Compound 5	14

6. Compound 6	14
7. Compound 7	15
8. Compound 8	15
9. Compound 9	16
10. Compound 10	16
11. Compound 11	16
12. Compound 12	17
Discussion	40
References	41
Abstract in Korean	75

List of Scheme and Tables

Scheme 1. Isolation of compounds from <i>Poncirus trifoliata</i> Raf.	5
Table 1. ¹³ C and ¹ H NMR assignment for compound 1	18
Table 2. ¹³ C and ¹ H NMR assignment for compound 2	20
Table 3. ¹³ C and ¹ H NMR assignment for compound 3	22
Table 4. ¹³ C and ¹ H NMR assignment for compound 4	24
Table 5. ¹³ C and ¹ H NMR assignment for compound 5	26
Table 6. ¹³ C and ¹ H NMR assignment for compound 6	28
Table 7. ¹³ C and ¹ H NMR assignment for compound 7	30
Table 8. ¹³ C and ¹ H NMR assignment for compound 8	31
Table 9. ¹³ C and ¹ H NMR assignment for the aglycone moiety of compound 9	32
Table 10. ¹³ C NMR assignment for the sugar moieties of compound 9	33
Table 11. ¹³ C and ¹ H NMR assignment for the aglycone moiety of compound 10	34
Table 12. ¹³ C NMR assignment for the sugar moieties of compound 10	35
Table 13. ¹³ C and ¹ H NMR assignment for the aglycone moiety of compound 11	36

Table 14. ^{13}C NMR assignment for the sugar moieties of	
compound 11	37
Table 15. ^{13}C and ^1H NMR assignment for the aglycone moiety of	
compound 12	38
Table 16. ^{13}C NMR assignment for the sugar moieties of	
compound 12	39

List of Figures

Figure 1. ^1H and ^{13}C NMR spectrum of compound 1	44
Figure 2. COSY spectrum of compound 1	45
Figure 3. HMBC spectrum of compound 1	46
Figure 4. HSQC spectrum of compound 1	47
Figure 5. UV and CD spectrum of compound 1	48
Figure 6. ^1H and ^{13}C NMR spectrum of compound 2	49
Figure 7. COSY spectrum of compound 2	50
Figure 8. HMBC spectrum of compound 2	51
Figure 9. HSQC spectrum of compound 2	52
Figure 10. UV and CD spectrum of compound 2	53
Figure 11. ^1H and ^{13}C NMR spectrum of compound 3	54
Figure 12. COSY spectrum of compound 3	55
Figure 13. HMBC spectrum of compound 3	56
Figure 14. HSQC spectrum of compound 3	57
Figure 15. UV and CD spectrum of compound 3	58
Figure 16. ^1H and ^{13}C NMR spectrum of compound 4	59
Figure 17. COSY spectrum of compound 4	60
Figure 18. HMBC spectrum of compound 4	61
Figure 19. HSQC spectrum of compound 4	62

Figure 20. UV and CD spectrum of compound 4	63
Figure 21. ¹ H and ¹³ C NMR spectrum of compound 5	64
Figure 22. UV and CD spectrum of compound 5	65
Figure 23. ¹ H and ¹³ C NMR spectrum of compound 6	66
Figure 24. UV and CD spectrum of compound 6	67
Figure 25. ¹ H and ¹³ C NMR spectrum of compound 7	68
Figure 26. ¹ H and ¹³ C NMR spectrum of compound 8	69
Figure 27. ¹ H and ¹³ C NMR spectrum of compound 9	70
Figure 28. ¹ H and ¹³ C NMR spectrum of compound 10	71
Figure 29. ¹ H and ¹³ C NMR spectrum of compound 11	72
Figure 30. ¹ H and ¹³ C NMR spectrum of compound 12	73
Figure 31. $\Delta\delta$ values (= $\delta_S - \delta_R$; in ppm) obtained for (<i>S</i>)- and (<i>R</i>)-MTPA esters of compounds 1-6	74

Introduction

Sortase A (SrtA), a gram-positive transpeptidase, plays a key role in the pathogenesis of gram-positive bacteria by adhesion to the host cell via the covalent anchoring of virulence-associated proteins to cell wall peptidoglycans.¹ Accordingly, inhibitors of SrtA might be promising candidates for the treatment and prevention of gram-positive bacterial infection.²

In the preparatory study on the SrtA inhibition test from Korean traditional medicine, approximately two-hundreds of plant extracts were tested. Among them, the extract from the fruits of trifoliolate orange, *Poncirus trifoliata* Rafinesque, showed significant Srt A inhibition.

Poncirus trifoliata Rafinesque, a member of the family Rutaceae, is a deciduous or semi-deciduous shrub native to China and Korea.³ *Poncirus trifoliata* Raf. are widely used in oriental traditional medicine as a remedy for digestive ulcers, gastritis, emesis, inflammation, allergy, and dysentery.⁴ Also, diverse bio-activity was reported.⁵⁻¹⁰

The organic extracts from *Poncirus trifoliata* Raf. were separated by employing solvent-partitioning. Bioactivity-guided chromatographic separation afforded the isolation of in total twelve compounds, eight coumarins and four flavanone glycosides. Among the isolated compounds,

four coumarins were defined to be new natural products. The structures of four novel coumarins were determined by a combination of spectroscopic analyses.

The absolute configurations of the new coumarins (**1-4**) and known coumarins (**5** and **6**), previously unassigned, were determined by an application of the 2-methoxy-2-(trifluoromethyl)phenylacetic acid (MTPA) method.¹¹

Experimental section

1. General Experimental Procedures

Optical rotations were measured on a JASCO P-1020 polarimeter and using a 1 cm cell. CD data were obtained on a JASCO J-715 spectropolarimeter in MEOD solutions. IR spectra were obtained on a JASCO FT/IR-300E spectrophotometer. UV spectra were recorded on a Hitachi U-3010 spectrophotometer. NMR spectra were recorded in CDCl₃, DMSO, and CD₃OD solutions, on a Bruker AMX-500 and 125 MHz, respectively. Mass spectra were provided by the Korea Basic Science Institute, Daegu Branch, Korea. All solvents used were spectral grade or were distilled from glass prior to use.

2. Plant Material

Poncirus trifoliata Raf. were purchased from the Kyungdong-Market, Seoul, Korea, in April, 2016. A voucher specimen is on deposit at the Natural Products Research Institute, College of Pharmacy, Seoul National University.


3. Extraction and Isolation

Poncirus trifolata Raf. was extracted with Methanol (10 L X 3) and CH₂Cl₂ (10L X 3). The combined crude extract (113.8 g) were partitioned between H₂O (31.7 g) and n-butanol (81.7 g). The latter fraction was repartitioned between n-hexane (8.8 g) and 15% aqueous methanol (56.5 g). 15% aqueous methanol layer from solvent partitioning was subjected to C₁₈ reversed-phase vacuum flash chromatography sequential mixtures of H₂O and Methanol (elution order: 50%, 40%, 30%, 20%, 10% aqueous methanol, 100% methanol), 100% acetone, and finally 100% ethyl acetate.

Based on the result of ¹H NMR and bioactivity test, the fractions eluted with 50% aqueous methanol (5.6 g), 40% aqueous methanol (1.0 g), 30% aqueous methanol (1.1 g) were chosen for separation. The fraction eluted with 50% aqueous methanol was separated by semi-preparative reversed-phase HPLC (YMC-ODS column, 10mm X 250mm; 70% aqueous methanol) to afford 16.0, 70.0, 7.5, 4.0, 295.0, and 176.4 mg of Compounds **5**, **6**, **7**, **8**, **9**, and **10**, respectively.

The fraction eluted with 40% aqueous methanol was separated by semi-preparative reversed-phase HPLC (YMC-ODS column, 10mm X 250mm; 53% aqueous methanol) to afford 10.0, 3.0, 10.0, 302.6, and 105.5 mg of Compounds **1**, **2**, **3**, **11**, and **12**, respectively.

The fraction eluted with 30% aqueous methanol was separated by semi-preparative reversed-phase HPLC (YMC-ODS column, 10mm X 250mm; 43% aqueous methanol) to afford 4.5 mg of Compounds **4**.



Scheme 1. Isolation of compounds from *Poncirus trifoliata* Raf.

4. Preparation of the (S)- and (R)- α -Methoxy- α -trifluoromethylphenylacetic chloride (MTPA-Cl) Esterification of Compound 1-6

Reactions were performed as follows. To duplicate solution of **1** (0.9 mg, 2 μ mol each) and DMAP (0.5 mg each) in 0.8 mL of anhydrous pyridine were added (R)- or (S)-MTPA chloride (8 mg, 0.03 mmol each). The mixture was allowed to stand under N₂ at room temperature for 2 h. After reaction, samples were dried under vacuum and separated by HPLC (YMS-ODS column, 4.6mm X 250 mm; 25% aqueous methanol) to afford the pure (S)- and (R)-MTPA esters derivatives of **1**. The (S)- and (R)-MTPA esters of **2-6** were also prepared by similar method.

¹H NMR (Chloroform-*d*, 600 MHz) data of **1S**: δ_{H} 7.5579 (1H, d, $J = 9.7$ Hz, H-4), 7.54-7.17 (10H, m, MTPA-Ar), 7.2897 (1H, d, $J = 8.2$ Hz, H-5), 6.7043 (1H, d, $J = 8.7$ Hz, H-6), 6.1848 (1H, d, $J = 9.4$ Hz, H-3), 5.5459 (1H, dd, $J = 2.6, 7.7$ Hz, H-2'), 5.4605 (1H, dd, $J = 2.3, 11.5$ Hz, H-2''), 4.1992 (2H, m, H-1'), 3.5223 (3H, s, MTPA-OMe), 3.2726 (3H, s, MTPA-OMe), 3.2278 (1H, dd, $J = 11.9, 14.0$ Hz, H-1''a), 3.1831 (3H, s, OMe), 2.9657 (1H, dd, $J = 2.5, 14.3$ Hz, H-1''b), 1.3393 (3H, s, H-4''), 1.2582 (3H, s, H-5''), 1.0995 (3H, s, H-4'), 1.0803 (3H, s, H-5'); LRESIMS m/z 835.1 [M + Na]⁺ (calcd for C₄₀H₄₂F₆O₁₁, 812.3).

¹H NMR (Chloroform-*d*, 600 MHz) data of **1R**: δ_{H} 7.5976 (1H, d, $J = 9.7$ Hz, H-4), 7.55-7.15 (10H, m, MTPA-Ar), 7.2912 (1H, d, $J = 8.2$ Hz, H-5),

6.7522 (1H, d, $J = 8.7$ Hz, H-6), 6.2389 (1H, d, $J = 9.4$ Hz, H-3), 5.5914 (1H, dd, $J = 2.6, 7.7$ Hz, H-2'), 5.4443 (1H, dd, $J = 2.3, 11.5$ Hz, H-2''), 4.1945 (2H, m, H-1'), 3.5054 (3H, s, MTPA-OMe), 3.4186 (3H, s, MTPA-OMe), 3.3904 (1H, dd, $J = 11.9, 14.0$ Hz, H-1''a), 3.2123 (3H, s, OMe), 3.1119 (1H, dd, $J = 2.5, 14.3$ Hz, H-1''b), 1.3086 (3H, s, H-4''), 1.2179 (3H, s, H-5''), 1.1948 (3H, s, H-4'), 1.1883 (3H, s, H-5'); LRESIMS m/z 835.1 $[M + Na]^+$ (calcd for $C_{40}H_{42}F_6O_{11}$, 812.3).

1H NMR (Methanol- d_4 , 600 MHz) data of **2S**: δ_H 7.8684 (1H, d, $J = 9.1$ Hz, H-4), 7.54-7.17 (10H, m, MTPA-Ar), 7.5188 (1H, d, $J = 8.7$ Hz, H-5), 6.9172 (1H, d, $J = 8.7$ Hz, H-6), 6.2450 (1H, d, $J = 9.2$ Hz, H-3), 5.8427 (1H, dd, $J = 2.8, 11.4$ Hz, H-2'), 5.4390 (1H, dd, $J = 2.3, 7.9$ Hz, H-2''), 4.2344 (2H, m, H-1''), 3.5424 (3H, s, MTPA-OMe), 3.2945 (3H, s, OMe), 3.2353 (3H, s, MTPA-OMe), 3.2142 (1H, dd, $J = 11.3, 14.0$ Hz, H-1'a), 2.7834 (1H, dd, $J = 3.1, 14.1$ Hz, H-1'b), 1.3268 (3H, s, H-4''), 1.3268 (3H, s, H-5''), 1.1349 (3H, s, H-5'), 1.1402 (3H, s, H-4'); LRESIMS m/z 857.2 $[M - H + HCO_2H]^-$ (calcd for $C_{40}H_{42}F_6O_{11}$, 812.3).

1H NMR (Methanol- d_4 , 600 MHz) data of **2R**: δ_H 7.8971 (1H, d, $J = 9.1$ Hz, H-4), 7.56-7.04 (10H, m, MTPA-Ar), 7.5533 (1H, d, $J = 8.7$ Hz, H-5), 6.9830 (1H, d, $J = 8.7$ Hz, H-6), 6.2785 (1H, d, $J = 9.2$ Hz, H-3), 5.8257 (1H, dd, $J = 2.8, 11.4$ Hz, H-2'), 5.5402 (1H, dd, $J = 2.3, 7.9$ Hz, H-2''), 4.2781 (2H, m, H-1''), 3.5713 (3H, s, MTPA-OMe), 3.4219 (1H, dd, $J = 11.3, 14.0$ Hz, H-1'a), 3.3685 (3H, s, MTPA-OMe), 3.2677 (3H, s, OMe),

2.9286 (1H, dd, $J = 3.1, 14.1$ Hz, H-1'b), 1.3076 (3H, s, H-4''), 1.2993 (3H, s, H-5''), 1.2770 (3H, s, H-5'), 1.2686 (3H, s, H-4'); LRESIMS m/z 857.2 [M - H + HCO₂H]⁻ (calcd for C₄₀H₄₂F₆O₁₁, 812.3).

¹H NMR (Methanol-*d*₄, 600 MHz) data of **3S**: δ_H 7.9072 (1H, d, $J = 9.6$ Hz, H-4), 7.5747 (1H, d, $J = 8.7$ Hz, H-5), 7.55-7.27 (5H, m, MTPA-Ar), 7.0787 (1H, d, $J = 8.7$ Hz, H-6), 6.2597 (1H, d, $J = 9.6$ Hz, H-3), 5.5363 (1H, dd, $J = 1.8, 8.8$ Hz, H-2'), 4.4349 (1H, dd, $J = 1.8, 10.5$ Hz, H-1'a), 4.3364 (1H, dd, $J = 8.8, 10.8$ Hz, H-1'b), 3.9171 (1H, d, $J = 17.8$ Hz, H-1''a), 3.8415 (1H, d, $J = 18.0$ Hz, H-1''b), 3.5264 (3H, s, MTPA-OMe), 3.2195 (3H, s, OMe), 2.8185 (1H, m, H-3''), 1.1914 (3H, d, $J = 6.9$ Hz, H-4''), 1.1668 (3H, d, $J = 7.1$ Hz, H-5''), 1.1607 (3H, s, H-4'), 1.1361 (3H, s, H-5'); LRESIMS m/z 601.2 [M + Na]⁺ (calcd for C₃₀H₃₃F₃O₈, 578.2).

¹H NMR (Methanol-*d*₄, 600 MHz) data of **3R**: δ_H 7.8988 (1H, d, $J = 9.6$ Hz, H-4), 7.54-7.27 (5H, m, MTPA-Ar), 7.5321 (1H, d, $J = 8.7$ Hz, H-5), 6.9934 (1H, d, $J = 8.7$ Hz, H-6), 6.2601 (1H, d, $J = 9.6$ Hz, H-3), 5.5556 (1H, dd, $J = 1.8, 8.8$ Hz, H-2'), 4.3332 (1H, dd, $J = 1.8, 10.5$ Hz, H-1'a), 4.2047 (1H, dd, $J = 8.8, 10.8$ Hz, H-1'b), 4.004 (1H, d, $J = 17.8$ Hz, H-1''a), 3.9599 (1H, d, $J = 18.0$ Hz, H-1''b), 3.5406 (3H, s, MTPA-OMe), 3.2664 (3H, s, OMe), 2.8688 (1H, m, H-3''), 1.2732 (3H, s, H-4'), 1.2432 (3H, s, H-5'), 1.2220 (3H, d, $J = 6.9$ Hz, H-4''), 1.1829 (3H, d, $J = 7.1$ Hz, H-5''); LRESIMS m/z 601.2 [M + Na]⁺ (calcd for C₃₀H₃₃F₃O₈, 578.2).

¹H NMR (DMSO-*d*₆, 500 MHz) data of **4S**: δ_H 8.0136 (1H, d, *J* = 9.6 Hz, H-4), 7.6402 (1H, d, *J* = 8.6 Hz, H-5), 7.48-7.40 (5H, m, MTPA-Ar), 7.1648 (1H, d, *J* = 8.7 Hz, H-6), 6.3064 (1H, d, *J* = 9.5 Hz, H-3), 5.7644 (1H, dd, *J* = 2.7, 7.5 Hz, H-2'), 5.0210 (1H, s, H-4'a), 4.9536 (1H, s, H-4'b), 4.4049 (1H, dd, *J* = 7.7, 11.1 Hz, H-1'a), 4.3158 (1H, dd, *J* = 2.8, 11.1 Hz, H-1'b), 3.8741 (2H, s, H-1''), 3.4610 (3H, s, MTPA-OMe), 2.7374 (1H, m, H-3''), 1.6729 (3H, s, H-5'), 1.0778 (3H, d, *J* = 6.9 Hz, H-4''), 1.0606 (3H, d, *J* = 6.9 Hz, H-5''); LRESIMS *m/z* 569.2 [M + Na]⁺ (calcd for C₂₉H₂₉F₃O₇, 546.2).

¹H NMR (DMSO-*d*₆, 500 MHz) data of **4R**: δ_H 8.0010 (1H, d, *J* = 9.6 Hz, H-4), 7.6075 (1H, d, *J* = 8.6 Hz, H-5), 7.46-7.39 (5H, m, MTPA-Ar), 7.1111 (1H, d, *J* = 8.7 Hz, H-6), 6.2998 (1H, d, *J* = 9.5 Hz, H-3), 5.7632 (1H, dd, *J* = 2.7, 7.5 Hz, H-2'), 5.1695 (1H, s, H-4'a), 5.1243 (1H, s, H-4'b), 4.3548 (1H, dd, *J* = 7.7, 11.1 Hz, H-1'a), 4.2828 (1H, dd, *J* = 2.8, 11.1 Hz, H-1'b), 3.8213 (2H, s, H-1''), 3.4312 (3H, s, MTPA-OMe), 2.7309 (1H, m, H-3''), 1.8053 (3H, s, H-5'), 1.0829 (3H, d, *J* = 6.9 Hz, H-4''), 1.0516 (3H, d, *J* = 6.9 Hz, H-5''); LRESIMS *m/z* 569.2 [M + Na]⁺ (calcd for C₂₉H₂₉F₃O₇, 546.2).

¹H NMR (Methanol-*d*₄, 500 MHz) data of **5S**: δ_H 7.8802 (1H, d, *J* = 9.6 Hz, H-4), 7.55-7.11 (10H, m, MTPA-Ar), 7.5222 (1H, d, *J* = 8.7 Hz, H-5), 6.9033 (1H, d, *J* = 8.6 Hz, H-6), 6.2437 (1H, d, *J* = 9.5 Hz, H-3), 5.6641 (1H, dd, *J* = 2.5, 11.7 Hz, H-2'), 5.4254 (1H, dd, *J* = 1.9, 8.5 Hz, H-2''),

4.2004 (2H, m, H-1''), 3.5436 (3H, s, MTPA-OMe), 3.3494 (3H, s, MTPA-OMe), 3.2435 (1H, dd, $J = 1.7, 3.3$ Hz, H-1'a), 2.8924 (1H, dd, $J = 2.2, 13.9$ Hz, H-1'b), 1.3513 (3H, s, H-4''), 1.3242 (3H, s, H-5''), 1.1295 (3H, s, H-4'), 1.1244 (3H, s, H-5'); LRESIMS m/z 843.1 $[M - H + HCO_2H]^-$ (calcd for $C_{39}H_{40}F_6O_{11}$, 798.3).

1H NMR (Methanol- d_4 , 500 MHz) data of **5R**: δ_H 7.9166 (1H, d, $J = 9.6$ Hz, H-4), 7.57-7.06 (10H, m, MTPA-Ar), 7.5578 (1H, d, $J = 8.7$ Hz, H-5), 6.9856 (1H, d, $J = 8.6$ Hz, H-6), 6.2909 (1H, d, $J = 9.5$ Hz, H-3), 5.6799 (1H, dd, $J = 2.5, 11.7$ Hz, H-2'), 5.5543 (1H, dd, $J = 1.9, 8.5$ Hz, H-2''), 4.2593 (2H, m, H-1''), 3.5736 (3H, s, MTPA-OMe), 3.4080 (3H, s, MTPA-OMe), 3.4464 (1H, dd, $J = 1.7, 3.3$ Hz, H-1'a), 3.0482 (1H, dd, $J = 2.2, 13.9$ Hz, H-1'b), 1.3299 (3H, s, H-4''), 1.2972 (3H, s, H-5''), 1.2729 (3H, s, H-4'), 1.2729 (3H, s, H-5'); LRESIMS m/z 843.1 $[M - H + HCO_2H]^-$ (calcd for $C_{39}H_{40}F_6O_{11}$, 798.3).

1H NMR (Methanol- d_4 , 600 MHz) data of **6S**: δ_H 7.9102 (1H, d, $J = 9.6$ Hz, H-4), 7.5797 (1H, d, $J = 8.7$ Hz, H-5), 7.56-7.26 (5H, m, MTPA-Ar), 7.0812 (1H, d, $J = 8.7$ Hz, H-6), 6.2606 (1H, d, $J = 9.6$ Hz, H-3), 5.4438 (1H, dd, $J = 1.6, 9.1$ Hz, H-2'), 4.5132 (1H, dd, $J = 1.4, 10.9$ Hz, H-1'a), 4.3540 (1H, dd, $J = 9.1, 10.5$ Hz, H-1'b), 3.9109 (1H, d, $J = 17.8$ Hz, H-1''a), 3.8498 (1H, d, $J = 18.0$ Hz, H-1''b), 3.5338 (3H, s, MTPA-OMe), 2.8174 (1H, m, H-3''), 1.1896 (3H, d, $J = 6.9$ Hz, H-4''), 1.1665 (3H, d, $J = 6.9$ Hz, H-5''), 1.1590 (3H, s, H-4'), 1.1580 (3H, s, H-5'); LRESIMS m/z 565.2 $[M + H]^+$ (calcd for

C₂₉H₃₁F₃O₈, 564.2).

¹H NMR (Methanol-*d*₄, 600 MHz) data of **6R**: δ_H 7.8974 (1H, d, *J* = 9.6 Hz, H-4), 7.5339 (1H, d, *J* = 8.7 Hz, H-5), 7.53-7.26 (5H, m, MTPA-Ar), 6.9936 (1H, d, *J* = 8.7 Hz, H-6), 6.2592 (1H, d, *J* = 9.6 Hz, H-3), 5.4209 (1H, dd, *J* = 1.6, 9.1 Hz, H-2'), 4.3943 (1H, dd, *J* = 1.4, 10.9 Hz, H-1'a), 4.2122 (1H, dd, *J* = 9.1, 10.5 Hz, H-1'b), 3.9874 (1H, d, *J* = 17.8 Hz, H-1''a), 3.9469 (1H, d, *J* = 18.0 Hz, H-1''b), 3.5409 (3H, s, MTPA-OMe), 2.8574 (1H, m, H-3''), 1.2761 (3H, s, H-4'), 1.2612 (3H, s, H-5'), 1.2139 (3H, d, *J* = 6.9 Hz, H-4''), 1.1736 (3H, d, *J* = 6.9 Hz, H-5''); LRESIMS *m/z* 565.2 [M + H]⁺ (calcd for C₂₉H₃₁F₃O₈, 564.2).

Results

1. Compound 1

Compound **1** was obtained as a yellow amorphous powder. The ^1H NMR spectrum indicated a coumarin structure with aromatic signals at δ_{H} 7.60 (1H, d, $J = 9.6$ Hz), 7.30 (1H, d, $J = 8.7$ Hz), 6.85 (1H, d, $J = 8.7$ Hz), and 6.22 (1H, d, $J = 9.2$ Hz), methoxy signal at δ_{H} 3.24 (3H, s).

Absolute configuration of C-2' and C-2'' were determined by an application of the 2-methoxy-2-(trifluoromethyl)phenylacetic acid (MTPA) method (Figure 31).

Based upon the result of combined spectroscopic analyses, the structure of compound **1** was determined to be a new coumarin possessing methoxy group.

2. Compound 2

Compound **2** was obtained as a yellow amorphous powder. The ^1H NMR spectrum indicated a coumarin structure with aromatic signals at δ_{H} 7.88 (1H, d, $J = 9.2$ Hz), 7.48 (1H, d, $J = 8.7$ Hz), 7.05 (1H, d, $J = 8.7$ Hz), and 6.24 (1H, d, $J = 9.6$ Hz), methoxy signal at δ_{H} 3.24 (3H, s).

Absolute configuration of C-2' and C-2'' were determined by an application of the 2-methoxy-2-(trifluoromethyl)phenylacetic acid (MTPA) method

(figure 31).

Based upon the result of combined spectroscopic analyses, the structure of compound **2** was determined to be a new coumarin possessing methoxy group.

3. Compound 3

Compound **3** was obtained as a yellow amorphous powder. The ^1H NMR spectrum indicated a coumarin structure with aromatic signals at δ_{H} 7.86 (1H, d, $J = 9.5$ Hz), 7.51 (1H, d, $J = 8.6$ Hz), 7.04 (1H, d, $J = 8.6$ Hz), and 6.21 (1H, d, $J = 9.5$ Hz), methoxy signal at δ_{H} 3.26 (3H, s). ^{13}C NMR spectrum indicated a ketone signal at δ_{C} 214.0.

Absolute configuration of C-2' was determined by an application of the 2-methoxy-2-(trifluoromethyl)phenylacetic acid (MTPA) method (figure 31).

Based upon the result of combined spectroscopic analyses, the structure of compound **3** was determined to be a new coumarin possessing methoxy group.

4. Compound 4

Compound **4** was obtained as a yellow amorphous powder. The ^1H NMR spectrum indicated a coumarin structure with aromatic signals at δ_{H} 8.00 (1H, d, $J = 9.5$ Hz), 7.61 (1H, d, $J = 8.6$ Hz), 7.10 (1H, d, $J = 8.6$ Hz), and 6.28 (1H, d, $J = 9.6$ Hz), methylene signals at δ_{H} 4.87 (1H, s) and 5.02

(1H, s). ^{13}C NMR spectrum indicated a ketone signal at δ_{C} 210.0.

Absolute configuration of C-2' was determined by an application of the 2-methoxy-2-(trifluoromethyl)phenylacetic acid (MTPA) method (figure 31).

Based upon the result of combined spectroscopic analyses, the structure of compound **4** was determined to be a new coumarin possessing exo-methylene.

5. Compound 5

Compound **5** was obtained as a yellow amorphous powder. The ^1H NMR spectrum indicated a coumarin structure with aromatic signals at δ_{H} 7.90 (1H, d, $J = 9.5$ Hz), 7.49 (1H, d, $J = 8.5$ Hz), 7.05 (1H, d, $J = 8.6$ Hz), and 6.25 (1H, d, $J = 9.4$ Hz).

Absolute configuration of C-2' and C-2'' were determined by an application of the 2-methoxy-2-(trifluoromethyl)phenylacetic acid (MTPA) method (figure 31).

Based upon the result of combined spectroscopic analyses, the structure of compound **5** was identified to be ponciol.

6. Compound 6

Compound **6** was obtained as a yellow amorphous powder. The ^1H NMR spectrum indicated a coumarin structure with aromatic signals at δ_{H} 7.88 (1H, d, $J = 9.4$ Hz), 7.54 (1H, d, $J = 8.7$ Hz), 7.06 (1H, d, $J = 8.6$ Hz),

and 6.23 (1H, d, $J = 9.4$ Hz). ^{13}C NMR spectrum indicated a ketone signal at $\delta_{\text{C}} 214.1$.

Absolute configuration of C-2' was determined by an application of the 2-methoxy-2-(trifluoromethyl)phenylacetic acid (MTPA) method (figure 31).

Based upon the result of combined spectroscopic analyses, the structure of compound **6** was identified to be triphasiol.¹²

7. Compound 7

Compound **7** was obtained as a colorless crystal. The ^1H NMR spectrum indicated a coumarin structure with aromatic signals at δ_{H} 7.85 (1H, d, $J = 9.4$ Hz), 7.45 (1H, d, $J = 8.5$ Hz), 6.79 (1H, dd, $J = 2.3, 8.6$ Hz), and 6.18 (1H, d, $J = 9.4$ Hz).

Based upon the result of combined spectroscopic analyses, the structure of compound **7** was identified to be umbelliferone.^{13,14}

8. Compound 8

Compound **8** was obtained as white needles. The ^1H NMR spectrum indicated a coumarin structure with aromatic signals at δ_{H} 7.59 (1H, d, $J = 16.0$ Hz), 7.45 (1H, d, $J = 8.6$ Hz), 6.80 (1H, d, $J = 8.6$ Hz), and 6.29 (1H, d, $J = 15.9$ Hz).

Based upon the result of combined spectroscopic analyses, the structure of compound **8** was identified to be umbelliferone.¹⁵

9. Compound 9

Compound **9** was obtained as a yellow powder. The ^1H NMR spectrum indicated a flavanone structure with aromatic signals at δ_{H} 7.30 (2H, d, $J = 8.3$ Hz), 6.82 (2H, d, $J = 8.3$ Hz), 6.17 (1H, d, $J = 2.3$ Hz), and 6.15 (1H, d, $J = 2.3$ Hz).

Based upon the result of combined spectroscopic analyses, the structure of compound **9** was identified to be naringin.^{16,17}

10. Compound 10

Compound **10** was obtained as a yellow powder. The ^1H NMR spectrum indicated a flavanone structure with aromatic signals at δ_{H} 7.32 (2H, d, $J = 8.3$ Hz), 6.82 (2H, d, $J = 8.3$ Hz), 6.20 (1H, d, $J = 2.3$ Hz), and 6.17 (1H, d, $J = 2.3$ Hz).

Based upon the result of combined spectroscopic analyses, the structure of compound **10** was identified to be narirutin.^{18,19}

11. Compound 11

Compound **11** was obtained as a white powder. The ^1H NMR spectrum indicated a flavanone structure with aromatic signals at δ_{H} 7.45 (2H, d, $J = 8.7$ Hz), 6.98 (2H, d, $J = 8.7$ Hz), 6.14 (1H, d, $J = 2.1$ Hz), and 6.10 (1H, d, $J = 2.1$ Hz).

Based upon the result of combined spectroscopic analyses, the structure of

compound **11** was identified to be poncirin.^{4,20}

12. Compound 12

Compound **12** was obtained as a white powder. The ¹H NMR spectrum indicated a flavanone structure with aromatic signals at δ_{H} 7.47 (2H, d, $J = 8.7$ Hz), 6.98 (2H, d, $J = 8.7$ Hz), 6.14 (1H, d, $J = 2.3$ Hz), and 6.13 (1H, d, $J = 2.3$ Hz).

Based upon the result of combined spectroscopic analyses, the structure of compound **11** was identified to be neoponcirin.^{21,22,23}





Table 1. ^{13}C and ^1H NMR assignment for compound **1** in Chloroform-*d*

Position	δ_{C}	δ_{H}
2	161.3, C	
3	113.4, CH	6.22, d (9.2)
4	144.0, CH	7.60, d (9.6)
4a	113.3, C	
5	127.0, CH	7.30, d (8.7)
6	108.7, CH	6.85, d (8.7)
7	159.9, C	
8	116.7, C	
8a	153.6, C	
1'	69.6, CH ₂	4.08, dd (7.8, 10.1) 4.35, dd (2.7, 10.2)
2'	74.9, CH	3.81, dd (2.8, 7.4)
3'	76.4, C	
4'	21.7, CH ₃	1.23, s
5'	20.6, CH ₃	1.22, s
1''	25.6, CH ₂	2.99, dd (8.7, 13.8) 3.12, dd (2.4, 13.7)

2''	78.8, CH	3.64, dd (2.3, 10.1)
3''	73.6, C	
4''	25.7, CH ₃	1.34, s
5''	24.7, CH ₃	1.29, s
OMe	49.6, CH ₃	3.24, s




2

Table 2. ^{13}C and ^1H NMR assignment for compound **2** in Methanol- d_4

Position	δ_{C}	δ_{H}
2	161.7, C	
3	113.3, CH	6.24, d (9.6)
4	146.4, CH	7.88, d (9.2)
4a	114.6, C	
5	128.5, CH	7.48, d (8.7)
6	110.2, CH	7.05, d (8.7)
7	161.9, C	
8	118.0, C	
8a	154.9, C	
1'	71.4, CH_2	4.07, dd (7.8, 10.1) 4.42, dd (2.8, 10.1)
2'	77.3, CH	3.76, dd (2.8, 7.8)
3'	72.8, C	
4'	27.2, CH_3	1.28, s
5'	25.1, CH_3	1.25, s
1''	26.2, CH_2	3.09, m
2''	77.4, CH	3.79, dd (3.9, 8.9)

3''	79.3, C	
4''	22.0, CH ₃	1.29, s
5''	20.5, CH ₃	1.29, s
OMe	50.0, CH ₃	3.32, s




3

Table 3. ^{13}C and ^1H NMR assignment for compound **3** in Methanol- d_4

Position	δ_{C}	δ_{H}
2	163.1, C	
3	113.3, CH	6.21, d (9.5)
4	146.2, CH	7.86, d (9.5)
4a	114.4, C	
5	129.3, CH	7.51, d (8.6)
6	110.1, CH	7.04, d (8.6)
7	161.6, C	
8	113.1, C	
8a	154.5, C	
1'	71.9, CH ₂	4.01, dd (8.2, 10.1) 4.33, dd (2.4, 10.2)
2'	76.5, CH	3.78, dd (2.3, 8.1)
3'	77.8, C	
4'	22.4, CH ₃	1.24, s
5'	20.4, CH ₃	1.21, s
1''	36.0, CH ₂	4.09, d (1.4)
2''	214.0, C	
3''	42.0, CH	2.90, m

4''	18.9, CH ₃	1.18, d (1.4)
5''	19.0, CH ₃	1.20, d (1.2)
OMe	49.9, CH ₃	3.26, s



4

Table 4. ^{13}C and ^1H NMR assignment for compound **4** in $\text{DMSO-}d_6$

Position	δ_{C}	δ_{H}
2	160.0, C	
3	111.9, CH	6.28, d (9.6)
4	144.7, CH	8.00, d (9.5)
4a	112.5, C	
5	128.0, CH	7.61, d (8.6)
6	108.9, CH	7.10, d (8.6)
7	159.4, C	
8	112.3, C	
8a	152.8, C	
1'	71.5, CH_2	3.99, dd (2.7, 6.4) 4.04, dd (4.7, 9.8)
2'	72.2, CH	4.23, t (5.4)
3'	145.0, C	
4'	111.5, CH_2	4.87, s 5.02, s
5'	18.0, CH_3	1.70, s
1''	34.4, CH_2	3.95, d (4.7)
2''	210.0, C	

3''	39.6, CH	2.82, m
4''	18.2, CH ₃	1.11, d (2.1)
5''	18.1, CH ₃	1.09, d (2.1)





Table 5. ^{13}C and ^1H NMR assignment for compound **5** in Methanol- d_4

Position	δ_{C}	δ_{H}
2	163.9, C	
3	113.2, CH	6.25, d (9.4)
4	146.5, CH	7.90, d (9.5)
4a	114.6, C	
5	128.5, CH	7.49, d (8.5)
6	110.2, CH	7.05, d (8.6)
7	162.0, C	
8	118.0, C	
8a	154.9, C	
1'	71.5, CH_2	4.07, dd (8.1, 10.1) 4.43, dd (2.8, 10.2)
2'	77.4, CH	3.77, dd (2.8, 8.0)
3'	72.9, C	
4'	27.3, CH_3	1.29, s
5'	25.0, CH_3	1.25, s
1''	26.5, CH_2	3.10, m
2''	79.2, CH	3.69, dd (3.5, 9.2)

3''	74.4, C	
4''	26.3, CH ₃	1.31, s
5''	24.8, CH ₃	1.30, s



6

Table 6. ^{13}C and ^1H NMR assignment for compound **6** in Methanol- d_4

Position	δ_{C}	δ_{H}
2	163.1, C	
3	113.4, CH	6.23, d (9.4)
4	146.2, CH	7.88, d (9.4)
4a	114.5, C	
5	129.4, CH	7.54, d (8.7)
6	110.1, CH	7.06, d (8.6)
7	161.7, C	
8	113.1, C	
8a	154.6, C	
1'	71.9, CH_2	4.03, dd (8.2, 10.2) 4.37, dd (2.4, 10.0)
2'	77.8, CH	3.71, dd (2.5, 8.1)
3'	72.8, C	
4'	27.0, CH_3	1.26, s
5'	25.0, CH_3	1.22, s
1''	36.0, CH_2	4.12, d (6.4)
2''	214.1, C	
3''	42.0, CH	2.91, m

4''	18.9, CH ₃	1.20, d (2.3)
5''	19.0, CH ₃	1.21, d (2.3)




Table 7. ^{13}C and ^1H NMR assignment for compound **7** in Methanol- d_4

Position	δ_{C}	δ_{H}
2	163.8, C	
3	114.6, CH	6.18, d (9.4)
4	146.2, CH	7.85, d (9.4)
4a	113.3, C	
5	130.8, CH	7.45, d (8.5)
6	112.5, CH	6.79, dd (2.3, 8.5)
7	163.3, C	
8	103.5, C	6.71, d (2.3)
8a	157.4, C	





Table 8. ^{13}C and ^1H NMR assignment for compound **8** in Methanol- d_4

Position	δ_{C}	δ_{H}
1	127.5, C	
2	131.2, CH	7.45, d (8.6)
3	116.9, CH	6.80, d (8.6)
4	161.3, C	
5	116.9, CH	6.80, d (8.6)
6	131.2, CH	7.45, d (8.6)
7	146.5, CH	7.59, d (16.0)
8	116.1, CH	6.29, d (15.9)
9	171.4, C	




9

Table 9. ^{13}C and ^1H NMR assignment for the aglycone moiety of compound **9** in Methanol- d_4

Position	δ_{C}	δ_{H}
2	80.8, CH	5.33, dd (2.3, 12.8)
3	44.2, CH ₂	2.73, dd (2.3, 17.4) 3.14, dd (12.8, 17.4)
4	198.6, C	
5	165.1, C	
6	98.0, CH	6.15, d (2.3)
7	166.7, C	
8	96.9, CH	6.17, d (2.3)
9	164.7, C	
10	105.0, C	
1'	130.9, C	
2'	129.2, CH	7.30, d (8.3)
3'	116.5, CH	6.82, d (8.3)
4'	159.2, C	
5'	116.5, CH	6.82, d (8.3)
6'	129.2, CH	7.30, d (8.7)

Table 10. ^{13}C NMR assignment for the sugar moieties ofcompound **9** in Methanol- d_4

Position	δ_{C}
Glc 1''	99.5
2''	79.1
3''	79.0
4''	71.3
5''	78.2
6''	62.4
Rha 1'''	102.6
2'''	72.3
3'''	72.3
4'''	74.0
5'''	70.1
6'''	18.3



10

Table 11. ^{13}C and ^1H NMR assignment for the aglycone moiety of compound **10** in Methanol- d_4


Position	δ_{C}	δ_{H}
2	80.7, CH	5.38, dd (2.3, 12.8)
3	44.2, CH ₂	2.74, dd (2.3, 17.4) 3.14, dd (12.8, 17.4)
4	198.6, C	
5	165.1, C	
6	98.1, CH	6.17, d (2.3)
7	166.9, C	
8	97.3, CH	6.20, d (2.3)
9	159.2, C	
10	105.1, C	
1'	131.0, C	
2'	129.3, CH	7.32, d (8.3)
3'	116.5, CH	6.82, d (8.3)

4'	164.5, C	
5'	116.5, CH	6.82, d (8.3)
6'	129.3, CH	7.32, d (8.3)

Table 12. ^{13}C NMR assignment for the sugar moieties of

compound **10** in Methanol- d_4

Position	δ_{C}
Glc 1''	101.3
2''	74.8
3''	78.0
4''	71.4
5''	77.2
6''	67.5
Rha 1'''	102.2
2'''	72.2
3'''	72.5
4'''	74.2
5'''	69.9
6'''	18.0




11

Table 13. ^{13}C and ^1H NMR assignment for the aglycone moiety of compound **11** in $\text{DMSO-}d_6$

Position	δ_{C}	δ_{H}
2	78.4, CH	5.59, dd (2.7, 12.7)
3	42.1, CH_2	2.78, dd (2.7, 17.2) 3.19, m
4	197.1, C	
5	163.0, C	
6	96.4, CH	6.10, d (2.1)
7	164.9, C	
8	95.2, CH	6.14, d (2.1)
9	162.7, C	
10	103.4, C	
1'	130.4, C	
2'	128.4, CH	7.45, d (8.7)
3'	114.0, CH	6.98, d (8.7)
4'	159.6, C	
5'	114.0, CH	6.98, d (8.7)
6'	128.4, CH	7.45, d (8.7)
OMe	55.2, CH_3	3.77, s

Table 14. ^{13}C NMR assignment for the sugar moieties ofcompound **11** in $\text{DMSO-}d_6$

Position	δ_{C}
Glc 1''	100.4
2''	77.2
3''	76.1
4''	69.6
5''	76.9
6''	60.5
Rha 1'''	97.5
2'''	70.5
3'''	70.4
4'''	71.9
5'''	68.3
6'''	18.1



12

Table 15. ^{13}C and ^1H NMR assignment for the aglycone moiety of compound **12** in $\text{DMSO-}d_6$

Position	δ_{C}	δ_{H}
2	78.3, CH	5.58, dd (2.9, 12.6)
3	41.9, CH_2	2.78, dd (2.8, 17.2) 3.13, m
4	197.0, C	
5	163.0, C	
6	96.4, CH	6.13, d (2.3)
7	165.1, C	
8	95.4, CH	6.14, d (2.3)
9	162.5, C	
10	103.3, C	
1'	103.3, C	
2'	128.4, CH	7.47, d (8.7)
3'	113.9, CH	6.98, d (8.7)

4'	159.5, C	
5'	113.9, CH	6.98, d (8.7)
6'	128.4, CH	7.47, d (8.7)
OMe	55.2, CH ₃	3.77, s

Table 16. ¹³C NMR assignment for the sugar moieties of compound **12** in DMSO-*d*₆

Position	δ _C
Glc 1''	99.4
2''	72.9
3''	76.2
4''	69.6
5''	75.5
6''	66.0
Rha 1'''	100.6
2'''	70.3
3'''	70.7
4'''	72.1
5'''	68.3
6'''	17.9

Discussion

The previous phytochemical investigation of *Poncirus trifoliata* Rafinesque has afforded diverse secondary metabolites such as flavonoids, coumarins, terpenoids. Despite diverse bioactivity of *Poncirus trifoliata* Raf. was reported, however, sortase A which is an enzyme that plays a key role in cell wall protein anchoring and virulence in *Staphylococcus aureus* inhibitory activity was not researched.

In this study, bioactivity-guided chromatographic separation afforded the isolation of in total twelve compounds, eight coumarins and four flavanone glycosides. Among the isolated compounds, the structures of four new coumarins were determined by a combination of spectroscopic analyses.

In addition to determination of the planar structure, the absolute configuration of the new coumarins and known coumarins, previously unassigned, were determined by an application of the 2-methoxy-2-(trifluoromethyl)phenylacetic acid (MTPA) method. Detailed sortase A inhibitory activity of these compounds will be evaluated.

References

1. Antoni P. A. Hendrickx, Jonathan M. Budzik, So-Young Oh and Olaf Schneewind. *Nature Reviews Microbiology*. **2011**, *9*, 166-176.
2. Ikhoon Oh, Woo-Young Yang, Soon-Chun Chung, Tae-Yoon Kim, Ki-Bong Oh, and Jongheon Shin. *Arch. Pharm. Res.* **2011**, *34*, 217-222.
3. R. Tundis, M. Bonesi, V. Sicari, T.M. Pellicano, M.C. Tenuta, M. Leporini, F. Menichini, M.R. Loizzo. *Journal of Functional Foods*. **2016**, *25*, 477-485.
4. Chul Young Kim, Hee Ju Lee, Mi Kyeong Lee, Mi-Jeong Ahn, Jinwoong Kim. *J. Sep. Sci.* **2007**, *30*, 2693-2697.
5. Hwa-Jin Chung, Eun-Jung Park, Yuna Pyee, Guang Hua Xu, Seung-Ho Lee, Young Shik Kim, Sang Kook Lee. *Food and Chemical Toxicology*. **2011**, *49*, 2942-2946.
6. H.M. Kim, H.J. Kim, S.T. Park. *Journal of Ethnopharmacology*. **1999**, *66*, 283-288.
7. Hong Yu Zhou, Eun Myung Shin, Lian Yu Guo, Li Bo Zou, Guang Hua Xu, Seung-Ho Lee, Keum Ryon Ze, Eun-Kyung Kim, Sam Sik Kang, Yeong Shik Kim. *European Journal of Pharmacology*. **2007**, *572*, 239-248.
8. S. Rajkumar, A. Jebanesan. *Parasitol Res.* **2008**, *104*, 19-25.

9. Atiqur Rahman, Minkyun Na, and Sun Chul Kang. *Journal of Food Biochemistry*. **2012**, *36*, 217-223.
10. Tao Feng, Rui-Rui Wang, Xiang-Hai Cai, Yong-Tang Zheng, and Xiao-Dong Luo. *Chem. Pharm. Bull.* **2010**, *58*, 971-975.
11. Jose Manuel Seco, Emilio Quinoa, and Ricardo Riguera. *Chem. Rev.* **2004**, *104*, 17-117.
12. L. B. De Silva, W. H. M. W. Herath, R. C. Jennings, M. Mahendran, and G. P. Wannigama. *phytochemistry*. **1981**, *20*, 2776-2788.
13. Wei Peng, Ting Han, Yang Wang, Wen-Bo Xin, Cheng-Jian Zheng, and Lu-Ping Qin. *Chemistry of Natural Compounds*. **2011**, *46*, 959-960.
14. Jie Li, Qian Shen, Chen-Hao Bao, Li-Ting Chen and Xiang-Rong Li. *Molecules*. **2014**, *19*, 1603-1607.
15. Xiang Yuan, Huaixiu Wen, Yulei Cui, Minxia Fan, Zenggen Liu, Lijuan Mei, Yun Shao, Yiping Wang and Yanduo Tao. *Nat. Prod. Res.* **2017**, *31*, 362-366.
16. Zhi-You Yang, Tomoharu Kuboyama, Kohei Kazuma, Katsuhiro Konno, and Chihiro Tohda. *J. Nat. Prod.* **2015**, *78*, 2297-2300.
17. Mingfeng Xu, Lianqing Shen, Kuiwu Wang. *Fitoterapia*. **2009**, *80*, 461-464.
18. Yoshiharu Matsubara, Hiroyasu Kumamoto, Yoshitomi Iizuka, Tetsuo Murakami, Kozo Okamoto, Hideo Miyake, and Katsumi Yokoi. *Agric. Biol. Chem.* **1985**, *49*, 909-914.

19. Zhonghua Mao, Chunli Gan, Jiuxin Zhu, Nan Ma, Lijun Wu, Libo Wang, Xiaobo Wang. *Bioorganic & Medicinal Chemistry Letters*. **2017**, 27, 2812-2817.
20. Dong-Hyun Kim, Eun-Ah Bae, and Myung Joo Han. *Biol. Pharm. Bull.* **1999**, 22, 422-424.
21. Ming-Liang Li, Ling-Yu Xu, Zhen-Lin Li, Shi-Hui Qian, and Min-Jian Qin. *Chemistry of Natural Compounds*. **2014**, 50, 124-125.
22. Julia Cassani, Anna G. Escalona Araujo, Mariano Martinez-Vazquez, Norberto Manjarrez, Julia Moreno and Rosa Estrada-Reyes. *Molecules*. **2013**, 18, 7584-7599.
23. Jinbin Wei, Quanfang Huang, Facheng Bai, Jun Lin, Jinlan Nie, Shengjuan Lu, Chunyuang Lu, Renbin Huang, Zhongpeng Lu, Xing Lin. *Chemico-Biological Interactions*. **2017**, 261, 118-126.




Figure 1. ^1H and ^{13}C NMR spectrum of compound **1**




Figure 2. COSY spectrum of compound **1**




Figure 3. HMBC spectrum of compound **1**




Figure 4. HSQC spectrum of compound **1**




Figure 5. UV and CD spectrum of compound **1**




Figure 6. ^1H and ^{13}C NMR spectrum of compound **2**




Figure 7. COSY spectrum of compound **2**




Figure 8. HMBC spectrum of compound **2**




Figure 9. HSQC spectrum of compound 2




Figure 10. UV and CD spectrum of compound **2**




Figure 11. ^1H and ^{13}C NMR spectrum of compound **3**




Figure 12. COSY spectrum of compound **3**




Figure 13. HMBC spectrum of compound **3**




Figure 14. HSQC spectrum of compound **3**




Figure 15. UV and CD spectrum of compound **3**




Figure 16. ^1H and ^{13}C NMR spectrum of compound 4




Figure 17. COSY spectrum of compound 4




Figure 18. HMBC spectrum of compound 4




Figure 19. HSQC spectrum of compound 4




Figure 20. UV and CD spectrum of compound **4**




Figure 21. ^1H and ^{13}C NMR spectrum of compound **5**




Figure 22. UV and CD spectrum of compound **5**




Figure 23. ^1H and ^{13}C NMR spectrum of compound **6**




Figure 24. UV and CD spectrum of compound **6**




Figure 25. ^1H and ^{13}C NMR spectrum of compound 7




Figure 26. ^1H and ^{13}C NMR spectrum of compound **8**




Figure 27. ^1H and ^{13}C NMR spectrum of compound **9**




Figure 28. ^1H and ^{13}C NMR spectrum of compound **10**




Figure 29. ^1H and ^{13}C NMR spectrum of compound **11**




Figure 30. ^1H and ^{13}C NMR spectrum of compound **12**




Figure 31. $\Delta\delta$ values ($= \delta_S - \delta_R$; in ppm) obtained for (S)- and (R)-MTPA esters of compounds 1-6

생약 지실의 성분연구

서울대학교 대학원

약학과

천연물과학 전공

이 원 희

Sortase A (SrtA)는 그람양성 세균에 병독성을 갖게 하는 효소로 세균감염을 억제하는 데 있어 중요한 표적효소이다.

전통적으로 다양한 병적 증세의 치료에 사용되는 지실(*Poncirus trifoliata* Rafinesque)은 운향과 (Rutaceae)에 속하는 탕자나무(Trifoliata orange)의 덜 익은 열매로 높은 Srt A 억제를 보였다.

이에 본 연구에서는 Srt A를 억제하는 물질을 지실에서 찾기 위해서 다양한 크로마토그래피 기법을 통해 12종의 물질을 분리하였고, 각각의 분광 자료에 의하여 8종의 coumarins와 4종의 flavanone glycosides 임을 확인하였다. 특히 4종의 coumarins는 자연계에 처음 분리, 보고되는 물질임이 밝혀졌다.

또한 2-methoxy-2-(trifluoromethyl)phenylacetic acid

(MTPA) method를 이용하여 4종의 신규 coumarins와 구조 동정 된 2종의 coumarins의 입체구조를 결정할 수 있었다.

주요어 : herbal medicine, *Poncirus trifoliata* Rafinesque, coumarins, flavanone glycosides, sortase A inhibition.

학번 : 2016-29626