

ORIGINAL ARTICLE

Angiotensin II Type 1–Receptor Activating Antibodies in Renal-Allograft Rejection

Duska Dragun, M.D., Dominik N. Müller, Ph.D., Jan Hinrich Bräsen, M.D., Lutz Fritsche, M.D., Melina Nieminen-Kelhä, B.S., Ralf Dechend, M.D., Ulrich Kintscher, M.D., Birgit Rudolph, M.D., Johan Hoebeke, Ph.D., Diana Eckert, M.D., Istvan Mazak, M.D., Ralph Plehm, Ph.D., Constanze Schönemann, Ph.D., Thomas Unger, M.D., Klemens Budde, M.D., Hans-Hellmut Neumayer, M.D., Friedrich C. Luft, M.D., and Gerd Wallukat, Ph.D.

ABSTRACT

BACKGROUND

Antibodies against HLA antigens cause refractory allograft rejection with vasculopathy in some, but not all, patients.

METHODS

We studied 33 kidney-transplant recipients who had refractory vascular rejection. Thirteen had donor-specific anti-HLA antibodies, whereas 20 did not. Malignant hypertension was present in 16 of the patients without anti-HLA antibodies, 4 of whom had seizures. The remaining 17 patients had no malignant hypertension. We hypothesized that activating antibodies targeting the angiotensin II type 1 (AT₁) receptor might be involved.

RESULTS

Activating IgG antibodies targeting the AT₁ receptor were detected in serum from all 16 patients with malignant hypertension and without anti-HLA antibodies, but in no other patients. These receptor-activating antibodies are subclass IgG1 and IgG3 antibodies that bind to two different epitopes on the second extracellular loop of the AT₁ receptor. Tissue factor expression was increased in renal-biopsy specimens from patients with these antibodies. In vitro stimulation of vascular cells with an AT₁-receptor-activating antibody induced phosphorylation of ERK 1/2 kinase and increased the DNA binding activity of the transcription factors activator protein 1 (AP-1) and nuclear factor- κ B. The AT₁ antagonist losartan blocked agonistic AT₁-receptor antibody-mediated effects, and passive antibody transfer induced vasculopathy and hypertension in a rat kidney-transplantation model.

CONCLUSIONS

A non-HLA, AT₁-receptor-mediated pathway may contribute to refractory vascular rejection, and affected patients might benefit from removal of AT₁-receptor antibodies or from pharmacologic blockade of AT₁ receptors.

From the Departments of Nephrology (D.D., L.F., M.N.-K., D.E., K.B., H.-H.N.) and Pathology (B.R.), and the Center for Cardiovascular Research (D.D., M.N.-K., U.K., T.U.), Charité University Hospital; the Franz Volhard Clinic and HELIOS Klinikum (J.H.B., R.D., I.M., F.C.L.); the Max Delbrück Center for Molecular Medicine (D.N.M., R.P., G.W.); and the HLA Laboratory (C.S.) — all in Berlin; and the Institut de Biologie Moléculaire et Cellulaire, Centre National de la Recherche Scientifique, Strasbourg, France (J.H.). Address reprint requests to Dr. Dragun at the Department of Nephrology and Transplantation, Charité University Hospital, Campus Mitte, Schumannstr. 20–21, 10117 Berlin, Germany, or at duska.dragun@charite.de.

N Engl J Med 2005;352:558–69.

Copyright © 2005 Massachusetts Medical Society.

MANY ASPECTS OF T-CELL-MEDIATED responses in allograft rejection have been elucidated, yet humoral mechanisms are relatively unexplored. Vascular rejection that is refractory to intensified immunosuppression is the most important predictor of early and late graft loss.¹ The association of antidonor humoral reactivity against HLA antigens and vascular rejection has been established.² Other targets of the allograft-directed host response remain elusive. However, alloantibodies against the polymorphic non-HLA system were found in serum obtained before transplantation from patients in whom refractory rejection developed after they received kidney transplants from HLA-identical siblings.³ Allograft endothelium may be a primary target of the cytopathic actions of non-HLA antibodies, triggering endothelial-cell activation or apoptosis.^{4,5} Identification of non-HLA antigens that are relevant to rejection might provide insight into underlying mechanisms, define risk-related phenotypes, and facilitate the development of specific therapies.

We studied kidney-transplant recipients who had severe allograft dysfunction but did not have anti-HLA antibodies; in these patients, rejection was invariably accompanied by accelerated hypertension, and even convulsions, in a manner reminiscent of preeclampsia. In fact, the first patient we examined in this group (the index patient) had had preeclampsia 16 years earlier. Immunologic similarities between graft rejection and fetal survival in pregnancies complicated by preeclampsia are well recognized.⁶ We previously reported that agonistic antibodies that target the angiotensin II type 1 (AT₁) receptor may develop in women with preeclampsia after the 20th week of gestation,⁷ and we observed agonistic AT₁-receptor activity in nine kidney-transplant recipients during an episode of rejection.⁸ We then reasoned that similar mechanisms might be present in refractory allograft rejection and initiated a comprehensive investigation of this issue.

METHODS

STUDY PATIENTS AND BIOPSIES

We studied all patients with steroid-refractory acute renal-allograft rejection seen among those who received kidney transplants at the Charité University Hospital Campus Mitte in Berlin from January 1, 2000, through July 31, 2004. Serum samples were obtained prospectively from all patients during re-

jection episodes and were screened for donor-specific anti-HLA antibodies and also analyzed for agonistic antibodies targeting the AT₁ receptor. Serum samples from patients with steroid-refractory acute vascular rejection from three collaborating centers (the Charité University Hospital Campus Virchow Clinic, and the University Hospital Erlangen-Nürnberg, both in Germany, and the Ospedale Civile in Bergamo, Italy) were also examined. Written informed consent to use serum samples for research purposes was obtained from each patient while he or she was awaiting transplantation. The institutional review board of Charité University Hospital Berlin approved the protocols.

Allograft-biopsy specimens were processed by standard techniques and graded according to the Banff 97 classification, with updated scoring of the degree of humoral rejection.^{9,10} Biopsy specimens were stained with polyclonal antibodies for C4d (Biomedica)^{11,12} and tissue factor (provided by T. Luther, Dresden).

Initial immunosuppressive therapy consisted of a calcineurin inhibitor, mycophenolate mofetil, methylprednisolone, and antibody against interleukin-2 receptor for induction. Patients who were positive for donor-specific anti-HLA antibodies or C4d or who had vascular rejection manifested as endarteritis or necrotizing vasculitis with fibrinoid necrosis were treated with plasmapheresis together with intravenous immune globulin. Patients initially treated with cyclosporine were switched to high-dose tacrolimus when refractory rejection was detected.

BIOASSAYS AND ANALYSIS OF EPITOPES

Patients' serum samples were examined for the presence of AT₁-receptor-activating IgG in a functional bioassay that involves spontaneously beating cultured neonatal-rat cardiomyocytes that express several G-protein-coupled receptors, including the AT₁ receptor. Changes in the activation state of the cells is measured directly by counting the beating rate by means of computerized image analysis. Isolation, cultivation, and detection procedures have been described in detail previously.¹³ The dose-response relationship between activating IgG or angiotensin II concentration and the spontaneous beating rate is linear.

To study the specificity of the AT₁ receptor-mediated response, IgG-stimulated cells were treated with an AT₁-receptor blocker or angiotensin II type 2 (AT₂)-receptor blocker. Short, overlapping

synthetic peptides corresponding to the sequence of the second extracellular loop of the AT₁ receptor were used for identification of epitopes. To identify the IgG subclass, IgG from 10 patients from whom sufficient serum was available were treated with murine monoclonal antihuman IgG1, IgG2, IgG3, and IgG4 antibodies.

SURFACE-PLASMON-RESONANCE ANALYSIS

Binding of AT₁-receptor antibody was verified and quantified by surface-plasmon-resonance analysis (BIAcore). The IgG fraction from affected patients or control human IgG was allowed to pass over biotinylated peptides loaded on a streptavidin-immobilized chip in a biosensor (BIAcore) at different flow rates. After the association phase and the dissociation phase, the relative increase in response of AT₁-receptor peptide and control peptide (β_2 -adrenergic receptor) flow channels was determined. The kinetic variables were calculated with BIA evaluation software (Biosensor, version 3.1).

ASSAYS

Human coronary-artery endothelial cells and vascular smooth-muscle cells were stimulated with AT₁-receptor antibodies in the presence or absence of AT₁-receptor blocker or AT₂-receptor blocker. Protein extraction, sodium dodecyl sulfate–polyacrylamide-gel electrophoresis, and membrane treatment were performed as described by Dechend et al.¹⁴ Primary antibodies against polyclonal extracellular signal-regulated kinase (ERK) 1/2 and phosphorylated ERK 1/2 were used. Triplicate experiments with six patients' IgG preparations were performed. Electromobility-shift assays in vascular smooth-muscle cells stimulated with angiotensin II, AT₁-receptor antibodies, or control IgG were performed as described previously.¹⁴

ANIMAL MODEL OF KIDNEY TRANSPLANTATION

The low-responder Fischer 344-to-Lewis life supporting rat kidney-transplantation model was used to test the physiological effects of various antibodies. AT₁-receptor antibodies (a pool derived from two donors), control IgG, or vehicle was continuously infused into recipient rats by means of intraperitoneal, seven-day-release, osmotic minipumps (Alzet, Charles River Laboratories). After implantation of the minipumps, one kidney was transplanted into each rat (cold-ischemia time was two hours), and kidneys were harvested after seven days (n=4 per treatment group).¹⁵ For continuous blood-

pressure monitoring, telemetric transmitters were placed into the abdominal aortas of six additional rodent recipients, before transplantation. The histologic features of the rat allografts were analyzed after harvesting with standard techniques. Direct immunofluorescence with fluorescein isothiocyanate-labeled antihuman IgG (Dako) was used to determine intragraft binding of AT₁-receptor antibodies; cryosections were counterstained with 4',6-diamidino-2-phenylindole dihydrochloride (Vector).

STATISTICAL ANALYSIS

Continuous data are presented as medians (with ranges). We performed comparisons between groups with use of Fisher's exact test for categorical values and the Mann-Whitney U test for continuous variables.

RESULTS

Thirty-three patients with steroid-refractory rejection were identified (23 at Charité University Hospital Campus Mitte and 10 at collaborating centers). Thirteen had detectable donor-specific anti-HLA antibodies. Sixteen patients who did not have donor-specific anti-HLA antibodies had malignant hypertension, followed by convulsions in four. Four patients had neither donor-specific anti-HLA antibodies nor malignant hypertension. Figure 1 shows representative biopsy specimens from patients with malignant hypertension and refractory rejection who had no donor-specific anti-HLA antibodies; these showed either endarteritis (Fig. 1A) or fibrinoid necrosis (Fig. 1B). Patients whose biopsy specimens had evidence of fibrinoid necrosis underwent magnetic resonance angiography studies that showed multiple perfusion defects consistent with cortical infarctions (Fig. 1C). Subsequently, patients with anti-HLA antibodies had substantially better graft survival than did those who had malignant hypertension but did not have anti-HLA antibodies (Fig. 1D).

We studied the AT₁-receptor agonistic response and IgG subclasses with a bioassay that records the chronotropic response (the increase in the number of beats per minute) of spontaneously beating neonatal-rat cardiomyocytes when exposed to immunoglobulin from patients. Analysis of serum obtained from the 13 anti-HLA-positive patients and from the 16 anti-HLA-negative patients with malignant hypertension before transplantation and at




Figure 1. Features of Refractory Rejection in Patients without Donor-Specific Anti-HLA Antibodies.

Biopsy specimens from representative patients (stained with hematoxylin and eosin) show endarteritis (Panel A) and fibrinoid changes and necrosis in the wall of the interlobular artery, with a mural thrombus consistent with acute vascular rejection (Panel B). A magnetic resonance image of the kidney (Panel C) shows multiple perfusion defects consistent with the occurrence of cortical infarctions. The graph in Panel D demonstrates accelerated allograft loss in patients who had refractory vascular rejection with non-HLA antibodies, compared with patients who had vascular rejection and donor-specific anti-HLA antibodies. AT₁R-AA denotes AT₁-receptor antibodies.


the time of rejection indicated differences between the groups. Figure 2A demonstrates a mean increase of 26 to 30 beats per minute in cells exposed to IgG obtained before transplantation or at the time of rejection from patients without anti-HLA antibodies. Serum from 13 patients with humoral rejection mediated by donor-specific anti-HLA antibodies did not induce such increases. Immunoglobulin from the four remaining patients, who had steroid-refractory rejection without malignant hypertension or HLA-antibodies, did not induce a response (data not shown).

All 16 patients whose serum led to an increased chronotropic response were screened for hereditary and autoimmune causes of thrombophilia as a possible trigger of the observed thrombotic angiopathy; all tests were negative, as were additional

serologic tests for autoimmune disorders. Acute infection with cytomegalovirus was ruled out by pp65 antigen and reverse-transcriptase polymerase-chain-reaction analysis (in three participating German centers) or on clinical grounds (in Bergamo, Italy).

Specific angiotensin II-receptor antagonists were used to test whether the agonistic response to patients' IgG was specific for the AT₁ receptor. Losartan, but not the AT₂-receptor antagonist PD 123319, abolished the functional bioassay response (Fig. 2B). Angiotensin II and IgG from patients achieved results in the bioassay. Thus, IgG fractions of our patients' serum contained AT₁-receptor antibodies.

The demographic data from the 16 patients who were positive for AT₁-receptor antibodies



were compared with those for the 13 patients with donor-specific anti-HLA-antibodies. As shown in Table 1, patients with anti-HLA antibodies were more likely to have staining for C4d in their renal-biopsy specimens and had less rapid allograft loss. Otherwise, there were no significant differences in clinical or demographic characteristics between the two groups.

To estimate the incidence and prevalence of steroid-refractory allograft rejections in patients with AT₁-receptor antibodies, we studied all episodes of rejection that occurred after the 278 consecutive kidney transplantations performed at the Charité University Hospital Campus Mitte in Berlin between January 1, 2000, and July 31, 2004. During this period, 119 biopsy-proved episodes of rejection were

treated; 23 were refractory to steroids (19.3 percent of all rejection episodes). Rejection in patients with donor-specific anti-HLA-antibodies accounted for 9 (39.1 percent) of the 23 steroid-refractory rejection episodes (7.6 percent of all rejection episodes).

In comparison, rejections in association with AT₁-receptor antibodies had a similar incidence (10 episodes, accounting for 43.5 percent of steroid-refractory rejection episodes and 8.4 percent of all rejection episodes). The remaining four steroid-refractory rejection episodes at Charité Campus Mitte (accounting for 17.4 percent of steroid-refractory episodes and 3.4 percent of all rejection episodes) occurred in patients who had neither AT₁-receptor antibodies nor donor-specific anti-HLA antibodies. The crude prevalence of the rejection episodes associated with AT₁-receptor antibodies (10 among 278 kidney transplantations performed) during four years was 3.6 percent.

IgG fractions possessing agonistic activity were used to examine possible epitopes on the second extracellular loop of the AT₁ receptor, to which the antibodies would bind. Two amino acid sequences, AFHYESQ and ENTNIT, inhibited the agonistic activity of IgG obtained from the patients (Fig. 2C), indicating that these sites contain the IgG-binding epitopes for the antibody. Neutralization experiments in which monoclonal antihuman IgG1, IgG2, IgG3, and IgG4 antibodies were added to AT₁-receptor antibodies demonstrated that antibodies of the IgG1 and IgG3 subclass are responsible for the agonist-like effect of these antibodies (Fig. 2D).

The specificity of the bioassay findings was assessed with surface-plasmon-resonance analysis, which confirmed binding of AT₁-receptor antibodies to the peptide of the second extracellular loop of the AT₁ receptor. The BIA evaluation program used to analyze the results calculated the dissociation rate constant at $1.05 \pm 0.29 \times 10^{-3}$ per second and the pseudomolecular association rate constant at $5.02 \pm 1.40 \times 10^{-3}$ per second, revealing a high binding affinity of AT₁-receptor antibodies to the AT₁ receptor.

Thus, IgG fractions from patients contained AT₁-receptor antibodies of the IgG1 and IgG3 subclasses that bound to the second extracellular loop of the AT₁ receptor. Further studies were performed, on the basis of our hypothesis that these antibodies played a role in the pathogenesis of refractory vascular rejection. Vascular lesions in grafts that are rejected because of the presence of anti-


Table 1. Characteristics of Patients with AT₁-Receptor Antibodies and Patients with Donor-Specific Anti-HLA Antibodies but without AT₁-Receptor Antibodies.

Characteristic	AT ₁ -Receptor Antibodies (N=16)	Anti-HLA Antibodies (N=13)	P Value*
Transplant from living donor (no.)	2	4	0.36
Male sex (no.)	10	5	0.12
First kidney allograft (no.)	8	7	1.00
Cold-ischemia time (hr)			0.86
Median	11.2	11.3	
Range	1.6–20.0	1.4–16.2	
No. of HLA mismatches			0.80
Median	3	3	
Range	0–5	2–6	
Peak panel-reactive antibodies (%)			0.92
Median	34	30	
Range	0–100	0–96	
Panel-reactive antibodies at transplantation (%)			0.30
Median	1	0	
Range	0–99	0–84	
Age at transplantation (yr)			0.38
Median	37.6	44.1	
Range	15.1–67.2	20.1–65.2	
Time from transplantation to rejection (days)			0.10
Median	4	9	
Range	2–1217	3–680	
C4d-positive (no.)	5	13	<0.001

* P values were determined with use of Fisher's exact test for categorical variables and the Mann-Whitney U test for continuous variables.

HLA antibodies are generally attributed to complement activation. However, patients with AT₁-receptor antibodies did not have complement-fixing cytotoxic anti-HLA antibodies in their serum, and their renal-biopsy specimens generally did not show staining for C4d. AT₁-receptor antibodies were examined to see whether they could initiate signaling mediated by ERK 1/2 in vitro by means of endothelial cells and vascular smooth-muscle cells (Fig. 3A and 3B). AT₁-receptor antibody induced phosphorylation of ERK 1/2 (Fig. 3A and 3B), which peaked at 10 minutes without a change in total ERK 1/2 in the cells.

We also examined the possibility that the AT₁-receptor antibodies could activate the tran-



scription factor activator protein 1 (AP-1) downstream from ERK 1/2 (Fig. 3C). Both AT₁-receptor antibodies and angiotensin II induced AP-1 activity in vascular smooth-muscle cells, whereas control IgG caused no such induction. DNA-binding activity of nuclear factor-κB (NF-κB), a transcription factor pivotal in initiating immune responses and inflammatory disease,¹⁶ when studied in nuclear extracts from vascular smooth-muscle cells, was activated by AT₁-receptor antibodies, but not by IgG from control patients (Fig. 3D).

Tissue factor, which initiates the extrinsic co-

agulation pathway, is a target gene for AP-1 and NF-κB. Biopsy specimens obtained from patients during episodes of rejection mediated by AT₁-receptor antibodies revealed intense staining of tubular cells, inflammatory infiltrate, and peritubular capillaries (Fig. 4A). In contrast, only weakly positive staining was seen in biopsy specimens obtained after patients had been treated with losartan and plasmapheresis (Fig. 4B). Treatment consisting of plasmapheresis, intravenous immune globulin, and 100 mg of losartan daily in seven patients with AT₁-receptor antibodies resulted in significantly




Figure 4. Influence of Losartan and Plasmapheresis Treatment on Tissue-Factor Expression, Allograft Survival, and AT₁-Receptor-Antibody Activity in AT₁-Receptor-Antibody-Positive Rejection.

A representative biopsy specimen obtained at the time of the diagnosis of refractory rejection shows strong epithelial and endothelial staining for tissue factor (Panel A). Reduction in tissue factor expression after treatment with losartan and plasmapheresis is evident in Panel B. Graft survival in the Kaplan-Meier analysis was significantly increased in the patients with AT₁-receptor antibodies who received losartan and plasmapheresis, as compared with those who received nonspecific treatment (Panel C). Panel D shows decreased activity of AT₁-receptor antibodies six months after rejection in four patients treated with losartan and plasmapheresis. Values before transplantation and at rejection are shown for all AT₁-receptor-antibody-positive patients.

improved allograft survival, as compared with that in patients receiving standard antirejection treatment (Fig. 4C). Serum from four patients with the longest rejection-free follow-up (who are still receiving losartan) became negative for AT₁-receptor antibodies (Fig. 4D).

To demonstrate that the agonistic antibodies could induce vascular rejection, AT₁-receptor antibodies were infused into rats that had received kidney transplants. One week after transplantation, kidneys from all experimental animals showed evidence of endarteritis and intravascular infiltrates (Fig. 5A). Transplants from control animals infused with control IgG showed only endothelial activation (Fig. 5B). Infusion of AT₁-receptor antibodies had no effect on native kidneys in uninephrectomized

animals (data not shown). The act of transplanting non-native kidneys into recipients appeared to make the organs prone to the effects of AT₁-receptor antibodies. AT₁-receptor antibodies containing human AT₁-receptor-activating antibody-positive IgG but not control IgG could be detected within the muscular layer of injured arteries in the rats (Fig. 5C and 5D). The mean arterial pressure, measured by radiotelemetry, was higher in the group of rats with AT₁-receptor antibodies two weeks after transplantation than in controls (Fig. 5E).

DISCUSSION

Patients with refractory vascular allograft rejection and accelerated hypertension but without anti-HLA

antibodies were found to have pathogenic antibodies directed at two epitopes of the second extracellular loop of the AT₁ receptor in their serum. AT₁-receptor antibodies induced phosphorylation of ERK 1/2, AP-1 activation, and NF- κ B activation in vascular cells. An AT₁-receptor antagonist blocked the effects mediated by AT₁-receptor antibodies. Passive infusion of AT₁-receptor antibodies in a rat renal-transplantation model induced vascular changes and increased blood pressure. These data suggest that AT₁-receptor antibodies may be directly involved in the pathogenesis of vascular rejection.

Earlier we had observed AT₁-receptor activity during steroid-refractory, C4d-negative renal-allograft rejection in nine patients with hypertension, but not in eight patients with steroid-sensitive cellular rejection. In the present study, we increased the number of patients studied and clearly defined the pretransplantation activity of IgG1- and IgG3-subclass antibodies that bound to AT₁-receptor epitopes and induced signaling cascades. The clinical presentation of transplant recipients with AT₁-receptor antibodies appears similar to that of transplant recipients with anti-HLA antibodies, but with significantly worse allograft survival.

Similar agonistic antibodies against the AT₁ receptor have been reported in patients with preeclampsia.⁷ Recently, other investigators confirmed the existence of AT₁-receptor antibodies in patients with preeclampsia¹⁷ and also demonstrated that the antibodies induced mobilization of intracellular calcium and activation of a transcription factor, nuclear factor of activated T cells, in AT₁-receptor-transfected Chinese-hamster-ovary cells.¹⁷ The index patient in the present report had had preeclampsia many years earlier. Nevertheless, the antibodies in this patient and the other transplant recipients bound to an epitope that did not entirely coincide with the antibodies described earlier in the patients with preeclampsia.⁷

Antibodies to G-protein receptors have been identified in some patients with idiopathic dilated cardiomyopathy¹⁸ or malignant hypertension.¹⁹ In Graves' disease, autoantibodies activate the thyrotropin G-protein receptor, increasing target-organ activity.²⁰ Anti-endothelial-cell antibodies are non-HLA antibodies that may be related to the pathogenesis of allograft rejection.^{21,22} However, research on anti-endothelial-cell antibodies is hampered by the heterogeneity of endothelial cells in various vascular beds, a fact that makes identifying a stan-

Figure 5 (facing page). Induction of Acute Vascular Rejection in Transplant-Recipient Rats Treated with Human AT₁-Receptor Antibodies and Immunostaining for Human AT₁-Receptor Antibodies in Injured Vascular Media.


A representative section from a rat allograft seven days after transplantation and after seven days of infusion of AT₁-receptor antibodies shows evidence of endarteritis and intravascular inflammatory cells (Panel A, arrow; hematoxylin and eosin). No vascular changes occurred in rats treated with control IgG (Panel B, hematoxylin and eosin). AT₁-receptor antibodies were detected in the vascular layer of the injured vessel (arrowheads); arrows show fragmentation of the elastic membrane (Panel C, direct immunofluorescence). Lack of binding of human control IgG in the control group of rats and an intact elastic membrane are evident in Panel D (direct immunofluorescence). Radiotelemetric intraaortic recordings show an increase in mean arterial pressure in three rats that were infused with AT₁-receptor antibodies (AT₁R-AA), but not in one control animal treated with IgG (Panel E). Numbers of days before (minus sign) or after (plus sign) transplantation are shown in the bar graph. I bars indicate SEs.

dard detection method difficult.²³ Moreover, initially detected target antigens of anti-endothelial-cell antibodies were of unclear relevance.²³

We suggest that AT₁-receptor antibodies have similarities to anti-endothelial-cell antibodies since endothelial cells have one AT₁ receptor,²⁴ and AT₁-receptor antibodies induced phosphorylation of ERK 1/2 in endothelial cells. We further suggest that binding of AT₁-receptor antibodies to the AT₁ receptor is a critical step for activating the downstream signaling cascade, mimicking the action of angiotensin II and inducing damage to the allograft. Emerging information has established that angiotensin II acts as an inflammatory cytokine participating in various vascular disorders.²⁵

AT₁-receptor antibodies may induce inflammatory responses and contribute to allograft rejection by means of activation of NF- κ B target genes. Blockade of NF- κ B activity with decoy oligodeoxynucleotides reduced tubulointerstitial infiltration in rat renal allografts.²⁶ Expression of tissue factor as regulated by NF- κ B and AP-1 may increase procoagulatory activity of the injured vessels.

We previously reported NF- κ B and AP-1 subunit expression in biopsy specimens from several kidney-transplant recipients.⁸ We now report that biopsy specimens of patients with rejection associated with AT₁-receptor antibodies had evidence of increased tissue factor expression and secondary thrombotic occlusions and that tissue factor was



reduced after losartan treatment. It has previously been recognized that the coagulation system has a role in the pathogenesis of rejection.²⁷ An anti-cardiolipin antibody-mediated increase in tissue factor expression contributes to thrombosis in patients with the antiphospholipid syndrome.²⁸ In rat

cardiac allografts, tissue factor plays a critical role in clotting abnormalities and transplant arteriosclerosis.²⁹ Lesions in allografts in rats infused with AT₁-receptor antibodies resemble those observed in kidney-transplant recipients.

It is unknown whether AT₁-receptor antibodies

function only through proinflammatory and procoagulatory activity or also by means of specific immune responses. Experimental studies have implicated the renin-angiotensin system in the regulation of the specific immune response and immune-mediated renal injury.^{30,31} Since AT₁ receptors are present on human mononuclear cells,^{32,33} an effect of AT₁-receptor antibodies on T lymphocytes, monocytes, and dendritic cells appears to be likely. Furthermore, rejection in association with circulating anti-HLA antibodies is commonly characterized by a high incidence of severe vascular lesions with fibrinoid necrosis, whereas endarteritis and mononuclear cell tubulitis predominate in T-cell-mediated rejection.³⁴

We first observed endarteritis and later fibrinoid necrosis, tubulitis, and interstitial-cell infiltration in patients' biopsy specimens. These characteristics may define a distinct type of kidney-transplant rejection mediated by AT₁-receptor antibodies. The incidence of rejection among patients positive for AT₁-receptor antibodies in this study is similar to that of rejection due to donor-specific anti-HLA antibodies. However, it remains unclear why AT₁-receptor antibodies develop in certain patients and why these antibodies target the allograft.

Molecular mimicry may be important; cross-reactivity with microbial antigens has an important role in other pathologic processes associated with antibodies directed against G-protein receptors — for example, myasthenia gravis or the dilated cardiomyopathy seen in Chagas' disease.^{35,36} We have not tested the possibility that the AT₁ receptors in patients in whom AT₁-receptor antibodies develop might harbor specific polymorphisms.³⁷ We spec-

ulate that post-transplantation reperfusion injury may alter the intragraft expression of AT₁ receptor, change its density, or cause conformational changes. A permissive environmental phenomenon might enhance local intragraft immunoreactivity owing to an activated innate immune response.

Transplantation nephrologists have been reluctant to use angiotensin II blockers because of possible decreases in organ perfusion and filtration. However, this concern may be outweighed by the potential advantages of these drugs.³⁸ The seven patients discussed here who were treated with losartan, plasmapheresis, and intravenous immunoglobulin had amelioration of the antibody-mediated rejection process and remain free of rejection while receiving losartan. However, their treatment was neither randomly assigned nor blinded, and the numbers of patients are too small to permit us to draw firm conclusions. We speculate that detection of AT₁-receptor antibodies in patients on a waiting list for a transplant might identify those at risk for refractory rejection.

Dr. Fritsche reports having served as a consultant to Novartis, Hexal, Shire, and GDL, being a stockholder in Novartis, receiving grant support from Novartis and Wyeth, and receiving lecture fees from Hoffmann-La Roche. Dr. Kintscher reports having served on the advisory boards of Novartis and Takeda, receiving grant support from Boehringer Ingelheim, and receiving lecture fees from Takeda. Dr. Unger reports serving as a consultant to Novartis, Abbott, Takeda, and Merck and having received grant support from Takeda, Novartis, Bayer, and Boehringer Ingelheim. Dr. Neumayer reports having received consulting fees from Novartis and Shire and lecture fees from Novartis and Wyeth. Dr. Luft reports having served as a consultant and receiving grant support from Novartis, Boehringer Ingelheim, and CVRx.

We are indebted to Professor U. Frei of Humboldt University in Berlin, Professor G. Remuzzi and Dr. N. Perico of the Mario Negri Institute in Bergamo, Italy, and Professor K. Amann and Dr. C. Hugo of the University of Nürnberg-Erlangen for providing samples.

REFERENCES

- van Saase J, van der Woude FJ, Thorgood J, et al. The relation between acute vascular and interstitial renal allograft rejection and subsequent chronic rejection. *Transplantation* 1995;59:1280-5.
- Jeannet M, Pinn VW, Flax MH, Winn HJ, Russell PS. Humoral antibodies in renal allotransplantation in man. *N Engl J Med* 1970; 282:111-7.
- Ahern AT, Artruch SB, DellaPelle P, et al. Hyperacute rejection of HLA-AB-identical renal allografts associated with B lymphocytes and endothelial reactive antibodies. *Transplantation* 1982;33:103-6.
- Yard B, Spruyt-Gerritse M, Claas F, et al. The clinical significance of allospecific antibodies against endothelial cells detected with an antibody-dependent cellular cytotoxicity assay for vascular rejection and graft loss after renal transplantation. *Transplantation* 1993;55:1287-93.
- Le Bas-Bernardet S, Hourmant M, Coupel S, Bignon JD, Souillou JP, Charreau B. Non-HLA-type endothelial cell reactive alloantibodies in pre-transplant sera of kidney recipients trigger apoptosis. *Am J Transplant* 2003;3:167-77.
- Mellor AL, Munn DH. Immunology at the maternal-fetal interface: lessons for T cell tolerance and suppression. *Annu Rev Immunol* 2000;18:367-91.
- Wallukat G, Homuth V, Fischer T, et al. Patients with preeclampsia develop agonistic antibodies against the angiotensin AT1 receptor. *J Clin Invest* 1999;103:945-52.
- Dragun D, Bräsen JH, Schönemann C, et al. Patients with steroid refractory acute vascular rejection develop agonistic antibodies targeting angiotensin II type 1 receptor. *Transplant Proc* 2003;35:2104-5.
- Racusen LC, Solez K, Colvin RB, et al. The Banff 97 working classification of renal allograft pathology. *Kidney Int* 1999;55: 713-23.
- Racusen LC, Colvin RB, Solez K, et al. Antibody-mediated rejection criteria — an addition to the Banff 97 classification of renal allograft rejection. *Am J Transplant* 2003;3:708-14.
- Böhmig G, Exner M, Habicht A, et al. Capillary C4d deposition in kidney allografts: a specific marker of alloantibody-dependent graft injury. *J Am Soc Nephrol* 2002;13:1091-9.
- Regele H, Exner M, Watschinger B, et al. Endothelial C4d deposition is associated with inferior kidney allograft outcome inde-

- pently of cellular rejection. *Nephrol Dial Transplant* 2001;16:2058-66.
13. Wallukat G, Wollenberger A, Morwinski R, Pitschner HF. Anti- β 1-adrenoreceptor autoantibodies with chronotropic activity from the serum of patients with dilated cardiomyopathy: mapping of epitopes in the first and second extracellular loops. *J Mol Cell Cardiol* 1995;27:397-406. [Erratum, *J Mol Cell Cardiol* 1995;27:2529.]
 14. Dechend R, Homuth V, Wallukat G, et al. AT(1) receptor agonistic antibodies from preeclamptic patients cause vascular cells to express tissue factor. *Circulation* 2000;101:2382-7.
 15. Dragun D, Hoff U, Park JK, et al. Ischemia-reperfusion injury in renal transplantation is independent of the immunologic background. *Kidney Int* 2000;58:2166-77.
 16. Baeuerle P. κ B-NF- κ B structures: at the interface of inflammation control. *Cell* 1998;95:729-31.
 17. Thway TM, Shlykov SG, Day MC, et al. Antibodies from preeclamptic patients stimulate increased intracellular Ca²⁺ mobilization through angiotensin receptor activation. *Circulation* 2004;110:1612-9.
 18. Magnusson Y, Wallukat G, Waagstein F, Hjalmarsen A, Hoebeke J. Autoimmunity in idiopathic dilated cardiomyopathy: characterization of antibodies against the beta1-adrenoreceptor with positive chronotropic effect. *Circulation* 1994;89:2760-7.
 19. Fu ML, Herlitz H, Wallukat G, et al. Functional autoimmune epitope on α 1-adrenergic receptors in patients with malignant hypertension. *Lancet* 1994;344:1660-3.
 20. Rapaport B, McLachlan SM. Thyroid autoimmunity. *J Clin Invest* 2001;108:1253-9.
 21. Paul LC, Baldwin WM III, van Es LA. Vascular endothelial alloantigens in renal transplantation. *Transplantation* 1985;40:117-23.
 22. van der Woude FJ, Deckers JGM, Mallat MJK, et al. Tissue antigens in tubulointerstitial and vascular rejection. *Kidney Int* 1995;52:Suppl:S11-S13.
 23. Praprotnik S, Blank M, Meroni PL, Rozman B, Eldor A, Shoenfeld Y. Classification of anti-endothelial cell antibodies into antibodies against microvascular and macrovascular endothelial cells: the pathogenic and diagnostic implications. *Arthritis Rheum* 2001;44:1484-94.
 24. de Gasparo M, Catt KJ, Inagami T, Wright JW, Unger T. International Union of Pharmacology: XXIII. The angiotensin II receptors. *Pharmacol Rev* 2000;52:415-72.
 25. Ruiz-Ortega M, Lorenzo O, Suzuki Y, Ruperez M, Egido J. Proinflammatory actions of angiotensins. *Curr Opin Nephrol Hypertens* 2001;10:321-9.
 26. Vos IH, Govers S, Gröne HJ, et al. NF- κ B decoy oligodeoxynucleotides reduce monocyte infiltration in renal allografts. *FASEB J* 2000;14:815-22.
 27. Heidenreich S, Dercken C, August C, Koch HG, Nowak-Göttl U. High rate of acute rejections in renal allograft recipients with thrombophilic risk factors. *J Am Soc Nephrol* 1998;9:1309-13.
 28. Dobado-Berrios P, Lopez-Pedraza C, Velasco F, Cuadrado MJ. The role of tissue factor in the antiphospholipid syndrome. *Arthritis Rheum* 2001;44:2467-76.
 29. Hölschermann H, Bohle RM, Zeller H, et al. In situ detection of tissue factor within the coronary intima in rat cardiac graft vasculopathy. *Am J Pathol* 1999;154:211-20.
 30. Hisada Y, Sugaya T, Yamanouchi M, et al. Angiotensin II plays a pathogenic role in immune-mediated renal injury in mice. *J Clin Invest* 1999;103:627-35.
 31. Nataraj C, Oliverio M, Mannon RB, et al. Angiotensin II regulates cellular immune responses through a calcineurin-dependent pathway. *J Clin Invest* 1999;104:1693-701.
 32. Shimada K, Yazaki Y. Binding sites for angiotensin II in human mononuclear leukocytes. *J Biochem (Tokyo)* 1978;84:1013-5.
 33. Nahmod KA, Vermeulen ME, Raiden S, et al. Control of dendritic cell differentiation by angiotensin II. *FASEB J* 2003;17:491-3.
 34. Mauiyyedi S, Crespo M, Collins AB, et al. Acute humoral rejection in kidney transplantation: II. Morphology, immunopathology, and pathologic classification. *J Am Soc Nephrol* 2002;13:779-87.
 35. Schwimmbeck PL, Dyrberg T, Drachman DB, Oldstone MB. Molecular mimicry and myasthenia gravis: an autoantigenic site of the acetylcholine receptor α -subunit that has biologic activity and reacts immunologically with herpes simplex virus. *J Clin Invest* 1989;84:1174-80.
 36. Masuda MO, Levin M, De Oliveira SF, et al. Functionally active cardiac antibodies in chronic Chagas' disease are specifically blocked by *Trypanosoma cruzi* antigens. *FASEB J* 1998;12:1551-8.
 37. Miller JA, Thai K, Scholey JW. Angiotensin II type 1 receptor gene polymorphism predicts response to losartan and angiotensin II. *Kidney Int* 1999;56:2173-80.
 38. Remuzzi G, Perico N. Routine renin-angiotensin system blockade in renal transplantation? *Curr Opin Nephrol Hypertens* 2002;11:1-10.

Copyright © 2005 Massachusetts Medical Society.

APPLY FOR JOBS ELECTRONICALLY AT THE NEJM CAREERCENTER

Physicians registered at the NEJM CareerCenter can apply for jobs electronically using their own cover letters and CVs. You can keep track of your job-application history with a personal account that is created when you register with the CareerCenter and apply for jobs seen online at our Web site. Visit www.nejmjobs.org for more information.