

Document details

[Back to results](#) | 1 of 1

[Export](#) [Download](#) [Print](#) [E-mail](#) [Save to PDF](#) [Add to List](#) [More... >](#)

Metrics [View all metrics >](#)

Journal of Pure and Applied Microbiology
Volume 8, Issue SPEC. ISS. 1, May 2014, Pages 885-890

Effect of biodegradation on the physical properties of palm oil mill effluent (POME) using mixed culture of fungi (Article)

Alade, A.O.^{ab}, Jameel, A.T.^a, Muyibi, S.A.^a, Abdul Karim, M.I.^a, Alam, Z.Md.^a 

^aDepartment of Biotechnology Engineering, Faculty of Engineering, International Islamic University Malaysia, Kuala Lumpur, Malaysia

^bDepartment of Chemical Engineering, Faculty of Engineering, Ladoke Akintola University of Technology, P.M.B. 4000 Ogbomosho, Oyo State, Nigeria

Abstract

This study investigated the effects of biodegradation by the mixed culture of fungi (*Pithomyces sacchari* and *Pestalotiopsis maculans*) on selected physical properties of palm oil mill effluent (POME). Mixed culture inoculum (4% v/v) was added to autoclaved and raw POME samples, which were subjected to biodegradation at 120rpm and 35°C for six days. The pH, electrical conductivity, total dissolve solids and biosolids of the digested samples were quantified at 24 h intervals. These parameters for the autoclaved sample, at the end of the digestion period, were 6.88, 4.38mS/cm, 2.28 g/L and 25.6 g/L, respectively. These values were higher than 6.34, 4.24mS/cm, 2.22 g/L and 22.87g/L for the raw POME sample, respectively. The kinetic studies of the degradation of POME, based on the concentration of the biosolids, were also investigated. The kinetic studies show that the degradation of the raw POME sample best fits the zero order kinetic model ($R^2 = 0.96$), while the degradation of the autoclaved POME sample best fits the first order kinetic model ($R^2 = 0.83$). However, the digested POME may require further treatment in order to meet standard suitable for discharge into the water body.

Author keywords

Biodegradation First order Mixed culture Pestalotiopsis maculans Pithomyces sacchari Zero order

ISSN: 09737510

Source Type: Journal

Original language: English

Document Type: Article

Publisher: Journal of Pure and Applied Microbiology

 Jameel, A.T.; Department of Biotechnology Engineering, Faculty of Engineering, International Islamic University Malaysia, Malaysia; email:abasslad@yahoo.com

© Copyright 2014 Elsevier B.V., All rights reserved.

Metrics [View all metrics >](#)

1 Citation in Scopus

20th Percentile

0.15 Field-Weighted

Citation Impact

Cited by 1 document

Heavy metal ions adsorption from dairy industrial wastewater using activated carbon from milk bush kernel shell

Afolabi, T.J. , Alade, A.O. , Jimoh, M.O.
(2016) Desalination and Water Treatment

[View details of this citation](#)

Inform me when this document is cited in Scopus:

[Set citation alert >](#)

[Set citation feed >](#)

Related documents

Find more related documents in Scopus based on:

[Authors >](#) [Keywords >](#)

[Back to results](#) | 1 of 1

[Top of page](#)

About Scopus

What is Scopus

Content coverage

Scopus blog

Scopus API

Language

日本語に切り替える

切换到简体中文

切换到繁體中文

Русский язык

Customer Service

Help

Contact us

ELSEVIER

[Terms and conditions](#) [Privacy policy](#)

Copyright © 2017 Elsevier B.V. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

Cookies are set by this site. To decline them or learn more, visit our Cookies page.

 RELX Group™