



# Document details

< Back to results | 1 of 1

Export Download Print E-mail Save to PDF Add to List More... >

Full Text

View at Publisher

Marine and Freshwater Behaviour and Physiology  
Volume 49, Issue 3, 3 May 2016, Pages 159-171

## Morphogenesis of free neuromasts in the larvae of brown-marbled grouper *Epinephelus fuscoguttatus* (Article)

Mukai, Y.<sup>a</sup> Lim, L.S.<sup>b</sup>

<sup>a</sup>Department of Marine Science, Kulliyah of Science, International Islamic University Malaysia, Kuantan, Malaysia

<sup>b</sup>Borneo Marine Research Institute, University Malaysia Sabah, Kota Kinabalu, Malaysia

### Abstract

View references (43)

Newly hatched larvae had one pair of free neuromasts behind the eyes. As the larvae grew, free neuromasts increased in number. The apical surface of sensory epithelium widened and subsequently elongated. The number of sensory hair cells increased and the directions of maximum sensitivity became both anteroposterior and dorsoventral on the trunk. Before notochord flexion, only the anteroposterior type was observed. After notochord flexion, two types of neuromasts were observed on the trunk. On the head, the orientation of free neuromasts formed a tangential line to concentric circles around the eyes and nostrils. Free neuromasts on the head could therefore receive stimuli from various angles from predators or zooplanktons. This suggests that these free neuromasts play a role in compensating for a dead angle of vision, and an important role in detecting zooplankton under scotopic vision. Canal organs were observed on the head and operculum in 40-d-old animals. © 2016 Informa UK Limited, trading as Taylor & Francis Group.

### SciVal Topic Prominence ⓘ

Topic: Lateral Line System | Zebrafish | Artificial lateral

Prominence percentile: 84.606 ⓘ

### Author keywords

cupulae fish larvae Free neuromasts lateral line maximum sensitivity neuromast morphology

### Indexed keywords

GEOBASE Subject Index: fish larva morphogenesis morphology physiological response zooplankton

Species Index: Animalia Epinephelus fuscoguttatus

### Metrics ⓘ View all metrics >

3 Citations in Scopus  
25th percentile

0.25 Field-Weighted  
Citation Impact



### PlumX Metrics

Usage, Captures, Mentions,  
Social Media and Citations  
beyond Scopus.

### Cited by 3 documents

Identification of Candidate Growth-Related SNPs and Genes Using GWAS in Brown-Marbled Grouper (*Epinephelus fuscoguttatus*)

Yang, Y. , Wu, L. , Wu, X.  
(2020) *Marine Biotechnology*

Immunity, feed, and husbandry in fish health management of cultured *Epinephelus fuscoguttatus* with reference to *Epinephelus coioides*

Chieng, C.C.Y. , Daud, H.M. , Yusoff, F.M.  
(2018) *Aquaculture and Fisheries*

Feeding and development of warm water marine fish larvae in early life

Hu, J. , Liu, Y. , Ma, Z.  
(2018) *Emerging Issues in Fish Larvae Research*

View all 3 citing documents

Inform me when this document is cited in Scopus:

Set citation alert >

Set citation feed >

References (43)

[View in search results format >](#)

All [Export](#) [Print](#) [E-mail](#) [Save to PDF](#) [Create bibliography](#)

- 1 Blaxter, J.H.S.  
Development of sense organs and behaviour of teleost larvae with special reference to feeding and predator avoidance

(1986) *Transactions of the American Fisheries Society*, 115 (1), pp. 98-114. Cited 497 times.  
doi: 10.1577/1548-8659(1986)115[98:NLFCD0]2.0.CO;2

[View at Publisher](#)

- 2 Blaxter, J.H.S.  
Structure and development of the lateral line  
(1987) *Biol Rev*, 62, pp. 471-514. Cited 92 times.

- 3 Blaxter, J.H.S., Fuiman, L.A.  
Function of the free neuromasts of marine teleost larvae  
(1989) *The mechanosensory lateral line*, pp. 481-499. Cited 92 times.  
Coombs S., Görner P., Münz H., (eds), New York(NY): Springer-Verlag

- 4 Blaxter, J.H.S., Gray, J.A.B., Best, A.C.G.  
Structure and development of the free neuromasts and lateral line system of the herring  
(1983) *Journal of the Marine Biological Association of the United Kingdom*, 63 (2), pp. 247-260. Cited 61 times.  
doi: 10.1017/S0025315400070648

[View at Publisher](#)

- 5 Diaz, J.P., Prié-Granié, M., Kentouri, M., Varsamos, S., Connes, R.  
Development of the lateral line system in the sea bass  
(2003) *Journal of Fish Biology*, 62 (1), pp. 24-40. Cited 23 times.  
doi: 10.1046/j.1095-8649.2003.00004.x

[View at Publisher](#)

- 6 Disler, N.N.  
(1971) *Lateral line sense organs and their importance in fish behavior*. Cited 27 times.  
Mills H., Yariv M., (eds), Jerusalem: Israel Program for Science Translation

- 7 Flock, Å.  
Electron microscope and electrophysiological studies on the lateral line canal organ  
(1965) *Acta Otolaryngol Suppl*, 199, pp. 1-90. Cited 211 times.

Structure and development of free neuromasts in barramundi, *Lates calcarifer* (block)

Mukai, Y. , Chai, L.L. , Shaleh, S.R.M.  
(2007) *Zoological Science*

Development of sensory organs in larvae of African catfish *Clarias gariepinus*

Mukai, Y. , Tuzan, A.D. , Lim, L.S.  
(2008) *Journal of Fish Biology*

Neuromast formation in the prehatching embryos of the cod fish, *Gadus macrocephalus Tilesius*

Otsuka, M. , Nagai, S.-I.  
(1997) *Zoological Science*

[View all related documents based on references](#)

[Find more related documents in Scopus based on:](#)

[Authors >](#) [Keywords >](#)