

[< Back to results](#) | 1 of 1[Export](#) [Download](#) [Print](#) [E-mail](#) [Save to PDF](#) [Add to List](#) [More... >](#)[Full Text](#) [View at Publisher](#)Jurnal Teknologi  
Volume 79, Issue 5-3, 2017, Pages 81-87

## A model for arabidopsis thaliana cell suspension growth and sugar uptake kinetics (Article)

Puad, N.I.M.<sup>a</sup> [✉](#), Abd-Karim, K.<sup>b</sup>, Mavituna, F.<sup>c</sup> [👤](#)<sup>a</sup>Bioprocess and Molecular Engineering Research Unit (BPMERU), Department of Biotechnology Engineering, Kulliyah of Engineering, International Islamic University Malaysia, P.O. Box 10, Kuala Lumpur, Malaysia<sup>b</sup>School of Chemical Engineering, Universiti Sains Malaysia (USM) Engineering Campus, Seri Ampangan, Pulau, Nibong Tebal, Pinang, Malaysia<sup>c</sup>School of Chemical Engineering and Analytical Science, The University of Manchester, Oxford Rd, Manchester, United Kingdom

### Abstract

Arabidopsis thaliana (*A. thaliana*) is a small weed which is related to the cabbage and mustard family. This flowering plant has been used widely as a model plant in genetics and molecular biology research since it is the first plant the full sequenced genome. Thus, *A. thaliana* literature provides plentiful information from genomics and transcriptomics point of view. However, there is still a lack of physiological information regarding its cell suspension cultures which can be powerful research tools in Plant Biotechnology and especially in Plant Systems and Synthetic Biology. In this study, cell growth and sugar uptake of *A. thaliana* Col ecotype grown in the continuous dark condition were modelled using the modified Monod and Michelis-menten equations. The model included sucrose hydrolysis by the cell-wall invertase enzyme into hexoses (glucose and fructose) and consumption of these hexoses at different rates to support cell growth. All kinetic model parameters were obtained from a control experiment where Col cells were grown on 30 gL<sup>-1</sup> sucrose as well as other independent experiments where Col cells were supplied with different concentrations and combinations of sugars. The model adequately described and predicted the growth and sugars profile of *A. thaliana* cells. This model can also be applied for larger scale of growth with extended expressions for oxygen uptake rate, carbon dioxide production rate etc. © 2017 Penerbit UTM Press. All rights reserved.

### Author keywords

Arabidopsis thaliana Cell growth Columbia (Col) ecotype Kinetic mode Sugar uptake

### Funding details

| Funding number | Funding sponsor          | Acronym |
|----------------|--------------------------|---------|
|                | University of Manchester |         |

### Funding text

The authors would like to thank Professor Simon Turner's research group (Faculty of Life Sciences, University of Manchester) for kindly providing Arabidopsis plant seeds. They would also like to acknowledge Malaysian of Higher Education for financial support.

ISSN: 01279696

Source Type: Journal

Original language: English

DOI: 10.11113/jt.v79.11331

Document Type: Article

Publisher: Penerbit UTM Press

### Metrics [🔗](#)

0 Citations in Scopus

0 Field-Weighted

Citation Impact

### Cited by 0 documents

Inform me when this document is cited in Scopus:

[Set citation alert >](#)[Set citation feed >](#)

### Related documents

Find more related documents in Scopus based on:

[Authors >](#) [Keywords >](#)

## About Scopus

What is Scopus  
Content coverage  
Scopus blog  
Scopus API  
Privacy matters

## Language

日本語に切り替える  
切换到简体中文  
切换到繁體中文  
Русский язык

## Customer Service

Help  
Contact us

**ELSEVIER**

[Terms and conditions](#) [Privacy policy](#)

Copyright © 2017 Elsevier B.V. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

Cookies are set by this site. To decline them or learn more, visit our [Cookies page](#).

 RELX Group™