

Scopus

[Search](#) [Sources](#) [Alerts](#) [Lists](#) [Help](#) [SciVal](#) [Register](#) [Login](#)

Document details

[Back to results](#) | 1 of 1
[Export](#) [Download](#) [Print](#) [E-mail](#) [Save to PDF](#) [Add to List](#) [More...](#)
[Full Text](#) [View at Publisher](#)

 Tropical Journal of Pharmaceutical Research
 Volume 16, Issue 10, October 2017, Pages 2315-2324
[Open Access](#)
Comparative assessment of poly (D,L-lactide-co-glycolide) nanoparticles modified by either cetyltrimethylammonium bromide or chitosan for plasmid DNA adsorption (Article)

 Doolaanea, A.A.^{2b}, Mansor, N.I.², Nor, N.H.M.², Shafri, M.A.B.M.², Sukmasari, S.², Mohamed, F.^{2bc}
^aDepartment of Pharmaceutical Technology, International Islamic University Malaysia (IIUM), Kuantan, Malaysia

^bIKOP Sdn. Bhd, Pilot Plant Pharmaceutical Manufacturing, Faculty of Pharmacy, International Islamic University Malaysia (IIUM), Kuantan, Malaysia

^cKulliyah of Allied Health Sciences, International Islamic University Malaysia (IIUM), Kuantan, Malaysia
View additional affiliations [v](#)

Abstract

[View references \(25\)](#)

Purpose: To evaluate poly (D,L-lactide-co-glycolide) PLGA nanoparticles modified by cetyltrimethyl ammonium bromide (CTAB) or chitosan for plasmid DNA adsorption. Methods: PLGA nanoparticles were prepared by solvent diffusion method and modified by including CTAB in the aqueous (F1) or oil phase (F2), or by including low (F3) or medium (F4) molecular weight chitosan. The nanoparticles were characterised by differential scanning calorimetry (DSC) and Fourier transform infrared spectroscopy (FTIR), as well as for cell toxicity, cell uptake and transfection. Results: CTAB failed to confer positive charge on the nanoparticles. CTAB desorbed easily from F1 surface. This resulted in negative zeta potential, increased cytotoxicity as well as decreased cell uptake and transfection. In F2, CTAB was located mainly in PLGA matrix, resulting in negative charge with decreased cytotoxicity, and increased cell uptake and transfection compared to F1. On the other hand, chitosan-modified nanoparticles (F3 and F4) showed stronger interaction between chitosan and PLGA, leading to positively-charged particles, decreased cytotoxicity, as well as increased cell uptake and transfection. Amongst the four formulations, F4 exhibited the highest transfection. Conclusion: These results should aid in understanding how PLGA nanoparticles are modified by CTAB and chitosan. Modification with chitosan yields PLGA nanoparticles with higher DNA adsorption and transfection with lower cytotoxicity. © Pharmacotherapy Group.

Author keywords

[Cetyltrimethyl ammonium bromide \(CTAB\)](#) [Chitosan](#) [Gene therapy](#) [Nanoparticle](#) [Plasmid DNA adsorption](#) [Poly \(D,L-lactide-co-glycolide\) PLGA](#)

Indexed keywords

EMTREE drug terms:

[cetrimide](#) [chitosan](#) [nanoparticle](#) [plasmid DNA](#) [polyglactin](#)

EMTREE medical terms:

[adsorption](#) [Article](#) [biological activity](#) [comparative study](#) [cytotoxicity](#) [differential scanning calorimetry](#) [drug screening](#)
[fluorescence microscopy](#) [genetic transfection](#) [infrared spectroscopy](#) [nerve cell membrane conductance](#) [nonhuman](#)
[physical chemistry](#) [zeta potential](#)

Chemicals and CAS Registry Numbers:

cetrimide, 57-09-0, 6899-10-1, 8044-71-1; chitosan, 9012-76-4; polyglactin, 26780-50-7, 34346-01-5

ISSN: 15965996

Source Type: Journal

Original language: English

DOI: 10.4314/tjpr.v16i10.2

Document Type: Article

Publisher: University of Benin

References (25)

[View in search results format](#)
 All [Export](#) [Print](#) [E-mail](#) [Save to PDF](#) [Create bibliography](#)

- 1 Bala, I., Hariharan, S., Kumar, M.N.V.R.
 PLGA nanoparticles in drug delivery: The state of the art
 (2004) *Critical Reviews in Therapeutic Drug Carrier Systems*, 21 (5), pp. 387-422. Cited 437 times.
[doi: 10.1615/CritRevTherDrugCarrierSyst.v21.i5.20](https://doi.org/10.1615/CritRevTherDrugCarrierSyst.v21.i5.20)
[View at Publisher](#)
- 2 Doolaanea, A.A., Ismail, A.F.H., Mansor, N.I., Mohd Nor, N.H., Mohamed, F.
 Effect of surfactants on plasmid DNA stability and release from poly (D,L-lactide-co-glycolide) microspheres
 (2015) *Tropical Journal of Pharmaceutical Research*, 14 (10), pp. 1769-1778. Cited 2 times.
http://www.tjpr.org/admin/12389900798187/2015_14_10_6.pdf
[doi: 10.4314/tjpr.v14i10.6](https://doi.org/10.4314/tjpr.v14i10.6)
[View at Publisher](#)

Metrics

0 Citations in Scopus

0 Field-Weighted Citation Impact

 PlumX Metrics
 Usage, Captures, Mentions,
 Social Media and Citations
 beyond Scopus

Cited by 0 documents

Inform me when this document is cited in Scopus:

[Set citation alert >](#)[Set citation feed >](#)

Related documents

Find more related documents in Scopus based on:

[Authors >](#) [Keywords >](#)

cells

 Wu, G., He, X.-L., Zhang, H.-L.
 (2008) *Acta Academiae Medicinae Sinicae*

 Design of biodegradable nanoparticles: A novel approach to encapsulating poorly soluble phytochemical ellagic acid
 Bala, I., Bhardwaj, V., Hariharan, S.
 (2005) *Nanotechnology*

View all related documents based on references

Find more related documents in Scopus based on:

[Authors >](#) [Keywords >](#)