



Document details

< Back to results | 1 of 1

[Download](#)
[Print](#)
[Save to PDF](#)
[☆ Add to List](#)
[More... >](#)
[Full Text](#)
[View at Publisher](#)

 ScienceAsia [Open Access](#)
 Volume 44, Issue 2, June 2018, Pages 179-186

Determination of antioxidant activity of gum Arabic : An exudation from two different locations (Article)

 Mirghani, M.E.S.^{a,b} , Elnour, A.A.M.^{a,c}, Kabbashi, N.A.^a, Alam, M.Z.^a, Musa, K.H.^d, Abdullah, A.^d 
^aBioenvironmental Engineering Research Centre, Biotechnology Engineering, Faculty of Engineering, International Islamic University, Malaysia, PO Box 10, Gombak, Kuala Lumpur, 50728, Malaysia

^bInternational Institute for Halal Research and Training, International Islamic University Malaysia, PO Box 10, Gombak, Kuala Lumpur, 50728, Malaysia

^cDepartment of Biochemistry and Gum Processing, Gum Arabic Research Centre, University of Kordofan, Box 160, Elobied, Sudan

[View additional affiliations](#) 


Abstract

[View references \(37\)](#)

Gum Arabic (GA) is the main product of acacia trees. As a raw and commercial samples, GA was extracted with methanol and analysed to measure the antioxidant activity using five methods: 2,2-diphenyl-1-picrylhydrazyl (DPPH), Folin-Ciocalteu indexes (FCI), which indicate total phenolic compounds (TPC), oxygen radical absorbance capacity (ORAC), ferric reducing antioxidant power (FRAP), and cupric reducing antioxidant capacity (CUPRAC). This study used antioxidant assays to detect TPC and selected appropriate and inexpensive methods to determine the antioxidant capacity of GA samples. The results reveal that the FCI, ORAC, and CUPRAC are correlated highly with FRAP. Person's correlation coefficient (r) values are 0.98, 0.93, and 0.99, respectively, based on the sample size of ($n = 8$). This means that the TPC of GA is highly correlated with their antioxidant activities that are measured by these three methods. Hence the FCI, ORAC, and the CUPRAC methods are more effective and simpler. They had similar predictive power to the FRAP of GA antioxidant activity. Consequently, GA is generally recognized as being slightly acidic which May have been obtained from appropriate methods of the antioxidant capacity detection. This acidity is due to the electronic transfer mechanism based on the selection of the working pH. © 2018 Science Society of Thailand under Royal Patronage. All right reserved.

SciVal Topic Prominence

Topic: Voltammetry | Antioxidants | pulse voltammetry

 Prominence percentile: 94.800 

Reaxys Database Information


[View Compounds](#)

Author keywords

[Antioxidants](#)
[CUPRAC](#)
[DPPH](#)
[Extraction](#)
[FRAP](#)
[ORAC](#)
[TPC](#)

Funding details

Funding sponsor

NEW! SciVal Topic Prominence is now available in Scopus. 

 Which Topic is this article related to? [View the Topic.](#)

Metrics

0 Citations in Scopus

 0 Field-Weighted
Citation Impact


PlumX Metrics

 Usage, Captures, Mentions,
Social Media and Citations
beyond Scopus.

Cited by 0 documents

 Inform me when this document
is cited in Scopus:

[Set citation feed >](#)

Related documents

 Measurement of oxygen radical
absorbance capacity in biological
samples

 Cao, G. , Prior, R.L.
(1998) *Methods in Enzymology*

 Screening Methods to Measure
Antioxidant Activity of Sorghum
(Sorghum bicolor) and Sorghum
Products

 Awika, J.M. , Rooney, L.W. , Wu,
X.
(2003) *Journal of Agricultural and
Food Chemistry*

 Comparative study of different
methods to measure antioxidant
activity of resveratrol in the
presence of cyclodextrins

 Lucas-Abellán, C. , Mercader-
Ros, M.T. , Zafrilla, M.P.
(2011) *Food and Chemical
Toxicology*

 View all related documents based
on references

 Find more related documents in
Scopus based on:

International Islamic University Malaysia

IIUM

Universiti Kebangsaan Malaysia

UKM

Funding text

Acknowledgements: The authors would like to express their gratitude to the University of Kordofan Sudan, Gum Arabic Research Centre, Bioenvironmental Engineering Research Centre, Department of Biotechnology Engineering at the Faculty (Kulliyyah) of Engineering, International Islamic University Malaysia for providing necessary lab facilities and UKM. For this opportunity, authors also extend the appreciation to Dr Elbasheir Sallam for his significant, continuous, and unlimited financial support.

ISSN: 15131874

DOI: 10.2306/scienceasia1513-1874.2018.44.179

Source Type: Journal

Document Type: Article

Original language: English

Publisher: Science Society of Thailand under Royal Patronage

References (37)

View in search results format >

 All Export Print E-mail Save to PDF Create bibliography
 1 Chawla, R., Patil, G.R.

Soluble dietary fiber (Open Access)

(2010) *Comprehensive Reviews in Food Science and Food Safety*, 9 (2), pp. 178-196. Cited 104 times.<http://onlinelibrary.wiley.com/doi/10.1111/j.1541-4337.2009.00099.x/pdf>

doi: 10.1111/j.1541-4337.2009.00099.x

View at Publisher

 2 Kong, H., Yang, J., Zhang, Y., Fang, Y., Nishinari, K., Phillips, G.O.

Synthesis and antioxidant properties of gum arabic-stabilized selenium nanoparticles

(2014) *International Journal of Biological Macromolecules*, 65, pp. 155-162. Cited 58 times.

doi: 10.1016/j.ijbiomac.2014.01.011

View at Publisher

 3 Ahmed, A.A., Fedail, J.S., Musa, H.H., Kamboh, A.A., Sifaldin, A.Z., Musa, T.H.

Gum Arabic extracts protect against hepatic oxidative stress in alloxan induced diabetes in rats

(2015) *Pathophysiology*, 22 (4), pp. 189-194. Cited 8 times.www.elsevier.com/locate/pathophys

doi: 10.1016/j.pathophys.2015.08.002

View at Publisher

 4 Musa, K.H., Abdullah, A., Kuswandi, B., Hidayat, M.A.

A novel high throughput method based on the DPPH dry reagent array for determination of antioxidant activity

(2013) *Food Chemistry*, 141 (4), pp. 4102-4106. Cited 40 times.www.elsevier.com/locate/foodchem

doi: 10.1016/j.foodchem.2013.06.112

View at Publisher

NEW! SciVal Topic Prominence is now available in Scopus.



Which Topic is this article related to? View the Topic.

- 5 Arnao, M.B., Cano, A., Acosta, M.
Methods to measure the antioxidant activity in plant material. A comparative discussion
(1999) *Free Radical Research*, 31 (SUPPL.), pp. S89-S96. Cited 104 times.
-
- 6 Lemańska, K., Szymusiak, H., Tyrakowska, B., Zieliński, R., Soffers, A.E.M.F, Rietjens, I.M.C.M
The influence of pH on antioxidant properties and the mechanism of antioxidant action of hydroxyflavones
(2001) *Free Radical Biology and Medicine*, 31 (7), pp. 869-881. Cited 265 times.
doi: 10.1016/S0891-5849(01)00638-4
[View at Publisher](#)
-
- 7 Arabaci, B., Gulcin, I., Alwasel, S.
Capsaicin: A potent inhibitor of carbonic anhydrase isoenzymes ([Open Access](#))
(2014) *Molecules*, 19 (7), pp. 10103-10114. Cited 78 times.
<http://www.mdpi.com/1420-3049/19/7/10103/pdf>
doi: 10.3390/molecules190710103
[View at Publisher](#)
-
- 8 Liang, N., Kitts, D.D.
Antioxidant property of coffee components: Assessment of methods that define mechanism of action ([Open Access](#))
(2014) *Molecules*, 19 (11), pp. 19180-19208. Cited 69 times.
<http://www.mdpi.com/journal/molecules>
doi: 10.3390/molecules191119180
[View at Publisher](#)
-
- 9 Bondet, V., Brand-Williams, W., Berset, C.
Kinetics and mechanisms of antioxidant activity using the DPPH* free radical method
(1997) *LWT - Food Science and Technology*, 30 (6), pp. 609-615. Cited 695 times.
<http://www.elsevier.com/inca/publications/store/6/2/2/9/1/0/index.htm>
doi: 10.1006/food.1997.0240
[View at Publisher](#)
-
- 10 Karabegović, I.T., Stojičević, S.S., Veličković, D.T., Todorović, Z.B., Nikolić, N.T., Lazić, M.L.
The effect of different extraction techniques on the composition and antioxidant activity of cherry laurel (*Prunus laurocerasus*) leaf and fruit extracts
(2014) *Industrial Crops and Products*, 54, pp. 142-148. Cited 36 times.
doi: 10.1016/j.indcrop.2013.12.047
[View at Publisher](#)
-
- 11 Cao, G., Alessio, H.M., Cutler, R.G.
Oxygen-radical absorbance capacity assay for antioxidants
(1993) *Free Radical Biology and Medicine*, 14 (3), pp. 303-311. Cited 1105 times.
doi: 10.1016/0891-5849(93)90027-R
[View at Publisher](#)

- 12 Sueishi, Y., Ishikawa, M., Yoshioka, D., Endoh, N., Oowada, S., Shimmei, M., Fujii, H., (...), Kotake, Y.
Oxygen radical absorbance capacity (ORAC) of cyclodextrin-solubilized flavonoids, resveratrol and astaxanthin as measured with the ORAC-EPR method (Open Access)

(2012) *Journal of Clinical Biochemistry and Nutrition*, 50 (2), pp. 127-132. Cited 30 times.
http://www.jstage.jst.go.jp/article/jcfn/50/2/127/_pdf
doi: 10.3164/jcfn.11-21

[View at Publisher](#)

- 13 Cao, G., Sofic, E., Prior, R.L.
Antioxidant Capacity of Tea and Common Vegetables

(1996) *Journal of Agricultural and Food Chemistry*, 44 (11), pp. 3426-3431. Cited 938 times.
<http://pubs.acs.org/journal/jafcau>
doi: 10.1021/jf9602535

[View at Publisher](#)

- 14 Soto-Vaca, A., Gutierrez, A., Losso, J.N., Xu, Z., Finley, J.W.
Evolution of phenolic compounds from color and flavor problems to health benefits

(2012) *Journal of Agricultural and Food Chemistry*, 60 (27), pp. 6658-6677. Cited 72 times.
doi: 10.1021/jf300861c

[View at Publisher](#)

- 15 Niki, E.
Assessment of antioxidant capacity in vitro and in vivo

(2010) *Free Radical Biology and Medicine*, 49 (4), pp. 503-515. Cited 443 times.
doi: 10.1016/j.freeradbiomed.2010.04.016

[View at Publisher](#)

- 16 Smina, T.P., Mathew, J., Janardhanan, K.K., Devasagayam, T.P.A.
Antioxidant activity and toxicity profile of total triterpenes isolated from *Ganoderma lucidum* (Fr.) P. Karst occurring in South India

(2011) *Environmental Toxicology and Pharmacology*, 32 (3), pp. 438-446. Cited 49 times.
doi: 10.1016/j.etap.2011.08.011

[View at Publisher](#)

- 17 Prior, R.L., Wu, X., Schaich, K.
Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements

(2005) *Journal of Agricultural and Food Chemistry*, 53 (10), pp. 4290-4302. Cited 2566 times.
doi: 10.1021/jf0502698

[View at Publisher](#)

- 18 Apak, R., Güçlü, K., Özyürek, M., Karademir, S.E.
Novel total antioxidant capacity index for dietary polyphenols and vitamins C and E, using their cupric ion reducing capability in the presence of neocuproine: CUPRAC method

(2004) *Journal of Agricultural and Food Chemistry*, 52 (26), pp. 7970-7981. Cited 812 times.
<http://pubs.acs.org/journal/jafcau>
doi: 10.1021/jf048741x

[View at Publisher](#)

NEW! SciVal Topic Prominence is now available in Scopus.

Which Topic is this article related to? [View the Topic.](#)



- 19 Benzie, I.F.F., Strain, J.J.
The ferric reducing ability of plasma (FRAP) as a measure of 'antioxidant power': The FRAP assay
(1996) *Analytical Biochemistry*, 239 (1), pp. 70-76. Cited 8890 times.
<http://www.elsevier.com/inca/publications/store/6/2/2/7/8/1/index.htm>
doi: 10.1006/abio.1996.0292
[View at Publisher](#)
-
- 20 Wayner, D.D.M., Burton, G.W., Ingold, K.U., Locke, S.
Quantitative measurement of the total, peroxy radical-trapping antioxidant capability of human blood plasma by controlled peroxidation. The important contribution made by plasma proteins ([Open Access](#))
(1985) *FEBS Letters*, 187 (1), pp. 33-37. Cited 520 times.
doi: 10.1016/0014-5793(85)81208-4
[View at Publisher](#)
-
- 21 Caillet, S., Yu, H., Lessard, S., Lamoureux, G., Ajdukovic, D., Lacroix, M.
Fenton reaction applied for screening natural antioxidants
(2007) *Food Chemistry*, 100 (2), pp. 542-552. Cited 75 times.
doi: 10.1016/j.foodchem.2005.10.009
[View at Publisher](#)
-
- 22 Brand-Williams, W., Cuvelier, M.E., Berset, C.
Use of a free radical method to evaluate antioxidant activity
(1995) *LWT - Food Science and Technology*, 28 (1), pp. 25-30. Cited 9185 times.
doi: 10.1016/S0023-6438(95)80008-5
[View at Publisher](#)
-
- 23 Gheldof, N., Engeseth, N.J.
Antioxidant capacity of honeys from various floral sources based on the determination of oxygen radical absorbance capacity and inhibition of in vitro lipoprotein oxidation in human serum samples
(2002) *Journal of Agricultural and Food Chemistry*, 50 (10), pp. 3050-3055. Cited 337 times.
doi: 10.1021/jf0114637
[View at Publisher](#)
-
- 24 Chumark, P., Khunawat, P., Sanvarinda, Y., Phornchirasilp, S., Morales, N.P., Phivthong-ngam, L., Ratanachamnong, P., (...), Pongrapeeporn, K.-u.S.
The in vitro and ex vivo antioxidant properties, hypolipidaemic and antiatherosclerotic activities of water extract of *Moringa oleifera* Lam. leaves
(2008) *Journal of Ethnopharmacology*, 116 (3), pp. 439-446. Cited 132 times.
doi: 10.1016/j.jep.2007.12.010
[View at Publisher](#)
-

- 25 Apak, R., Özyürek, M., Güçlü, K., Çapanoğlu, E.
Antioxidant activity/capacity measurement. 1. Classification, physicochemical principles, mechanisms, and electron transfer (ET)-based assays (Open Access)
(2016) *Journal of Agricultural and Food Chemistry*, 64 (5), pp. 997-1027. Cited 85 times.
<http://pubs.acs.org/journal/jafcau>
doi: 10.1021/acs.jafc.5b04739
View at Publisher
-
- 26 Soares, A.A., de Souza, C.G.M., Daniel, F.M., Ferrari, G.P., da Costa, S.M.G., Peralta, R.M.
Antioxidant activity and total phenolic content of *Agaricus brasiliensis* (*Agaricus blazei* Murri) in two stages of maturity
(2009) *Food Chemistry*, 112 (4), pp. 775-781. Cited 109 times.
doi: 10.1016/j.foodchem.2008.05.117
View at Publisher
-
- 27 Apak, R., Güçlü, K., Özyürek, M., Çelik, S.E.
Mechanism of antioxidant capacity assays and the CUPRAC (cupric ion reducing antioxidant capacity) assay
(2008) *Microchimica Acta*, 160 (4), pp. 413-419. Cited 184 times.
doi: 10.1007/s00604-007-0777-0
View at Publisher
-
- 28 Musa, K.H., Abdullah, A., Jusoh, K., Subramaniam, V.
Antioxidant Activity of Pink-Flesh Guava (*Psidium guajava* L.): Effect of Extraction Techniques and Solvents
(2011) *Food Analytical Methods*, 4 (1), pp. 100-107. Cited 71 times.
doi: 10.1007/s12161-010-9139-3
View at Publisher
-
- 29 Shahidi, F., Zhong, Y.
Measurement of antioxidant activity
(2015) *Journal of Functional Foods*, Part B 18, pp. 757-781. Cited 147 times.
http://www.elsevier.com/wps/find/journaldescription.cws_home/717426/description#description
doi: 10.1016/j.jff.2015.01.047
View at Publisher
-
- 30 Thaipong, K., Boonprakob, U., Crosby, K., Cisneros-Zevallos, L., Hawkins Byrne, D.
Comparison of ABTS, DPPH, FRAP, and ORAC assays for estimating antioxidant activity from guava fruit extracts
(2006) *Journal of Food Composition and Analysis*, 19 (6-7), pp. 669-675. Cited 1124 times.
doi: 10.1016/j.jfca.2006.01.003
View at Publisher
-
- 31 Gul, H.I., Kucukoglu, K., Yamali, C., Bilginer, S., Yuca, H., Ozturk, I., Taslimi, P., (...), Supuran, C.T.
Synthesis of 4-(2-substituted hydrazinyl)benzenesulfonamides and their carbonic anhydrase inhibitory effects (Open Access)
(2016) *Journal of Enzyme Inhibition and Medicinal Chemistry*, 31 (4), pp. 568-573. Cited 45 times.
doi: 10.3109/14756366.2015.1047359

NEW! SciVal Topic Prominence is now available in Scopus.

View at Publisher

Which Topic is this article related to? View the Topic.



32 Wang, H., Cao, G., Prior, R.L.

Total antioxidant capacity of fruits

(1996) *Journal of Agricultural and Food Chemistry*, 44 (3), pp. 701-705. Cited 1286 times.

<http://pubs.acs.org/journal/jafcau>

doi: 10.1021/jf950579y

[View at Publisher](#)

33 Giovanelli, G., Buratti, S.

Comparison of polyphenolic composition and antioxidant activity of wild Italian blueberries and some cultivated varieties

(2009) *Food Chemistry*, 112 (4), pp. 903-908. Cited 164 times.

doi: 10.1016/j.foodchem.2008.06.066

[View at Publisher](#)

34 Moylan, S., Berk, M., Dean, O.M., Samuni, Y., Williams, L.J., O'Neil, A., Hayley, A.C., (...), Maes, M.

Oxidative & nitrosative stress in depression: Why so much stress?

(2014) *Neuroscience and Biobehavioral Reviews*, 45, pp. 46-62. Cited 156 times.

www.elsevier.com/locate/neubiorev

doi: 10.1016/j.neubiorev.2014.05.007

[View at Publisher](#)

35 Edet, E., Ofem, J., Igile, G., Ofem, O., Zainab, D., Ak-Waowo, G.

Antioxidant capacity of different African seeds and vegetables and correlation with the contents of ascorbic acid, phenolics and flavonoids

(2015) *J Med Plant Res*, 9, pp. 454-461.

36 Sahu, N., Saxena, J.

Different methods for determining antioxidant activity: A review

(2013) *Indo Am J Pharmaceut Res*, 3, pp. 7025-7028.

37 Tabart, J., Kevers, C., Pincemail, J., Defraigne, J.-O., Dommes, J.

Evaluation of spectrophotometric methods for antioxidant compound measurement in relation to total antioxidant capacity in beverages

(2010) *Food Chemistry*, 120 (2), pp. 607-614. Cited 34 times.

doi: 10.1016/j.foodchem.2009.10.031

[View at Publisher](#)

🔍 Mirghani, M.E.S.; Bioenvironmental Engineering Research Centre, Biotechnology Engineering, Faculty of Engineering, International Islamic University, Malaysia, PO Box 10, Gombak, Kuala Lumpur, Malaysia; email:elwathig@iiium.edu.my

© Copyright 2018 Elsevier B.V., All rights reserved.

ELSEVIER

[Terms and conditions ↗](#) [Privacy policy ↗](#)

Copyright © 2018 Elsevier B.V. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.
We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies.

 RELX Group™