

Close

Web of Science
Page 1 (Records 1 -- 1)

Print

**Record 1 of 1****Title:** Assessment of Peripartum Psychological Disturbances in Relation to Modes of Delivery**Author(s):** Radeef, AS (Radeef, Ali S.); Al-Kubaisi, MK (Al-Kubaisi, Muna Kh); John, AB (John, Akbar B.); Khairulnisa, AIN (Khairulnisa, Nur A., I)**Source:** MAKARA JOURNAL OF HEALTH RESEARCH **Volume:** 23 **Issue:** 1 **Pages:** 1-6 **DOI:** 10.7454/msk.v23i1.9773 **Published:** APR 2019**Times Cited in Web of Science Core Collection:** 0**Total Times Cited:** 0**Usage Count (Last 180 days):** 1**Usage Count (Since 2013):** 1**Cited Reference Count:** 29

Abstract: Background: Anxiety disorders are common during peripartum period. This study aimed to identify which mode of delivery was more prone to peripartum psychological disturbances. Methods: This prospective cohort study conducted among 541 pregnant women presented for spontaneous labour, induction or elective caesarean section in a Malaysian hospital. The severity of depression, anxiety and stress symptoms are assessed by using self-rated depression anxiety and stress scales in four different occasions; pre-labour/operation, 24 hours postnatal, 2 weeks, and at 6 weeks. Results: In the prenatal period, there was no statistically significant difference in the mean score of depression between women coming for labour (4.17 +/- 5.28) and elective CS (4.21 +/- 5.87). During puerperium; at 24 hours postnatal, 2 weeks and 6 weeks assessments there was no significant difference in depression score among patients with different modes of delivery. The overall scores were (4.23 +/- 5.46, 3.26 +/- 4.85, 0.78 +/- 2.37, 0.18 +/- 0.93) for patients going through vaginal delivery, instrumentation, emergency or elective CS. The same trend was found when anxiety and stress were analysed. Conclusions: The most prominent peripartum psychological disturbances symptom is anxiety. Minority of patients will experience persistence anxiety by the end of 6 weeks postpartum. The caesarean section and induction of labour does not increase the psychological impact.

Accession Number: WOS:000467042700001**Language:** English**Document Type:** Article**Author Keywords:** anxiety; caesarean section; depression; induction of labour; stress**KeyWords Plus:** POSTNATAL DEPRESSION SCALE; RISK-FACTORS; CESAREAN-SECTION; ANXIETY; STRESS; PREVALENCE; PREGNANCY; MALAYSIA; MOTHERS; HEALTH**Addresses:** [Radeef, Ali S.] Int Islamic Univ Malaysia, Kulliyah Med, Dept Psychiat, Kuantan 25200, Pahang, Malaysia.

[Al-Kubaisi, Muna Kh; Khairulnisa, Nur A., I] Int Islamic Univ Malaysia, Kulliyah Med, Dept Obstet & Gynaecol, Kuantan 25200, Pahang, Malaysia.

[John, Akbar B.] Int Islamic Univ Malaysia, Kulliyah Sci, Kuantan 25200, Pahang, Malaysia.

Reprint Address: Al-Kubaisi, MK (reprint author), Int Islamic Univ Malaysia, Kulliyah Med, Dept Obstet & Gynaecol, Kuantan 25200, Pahang, Malaysia.**E-mail Addresses:** mkubaysi@gmail.com**Publisher:** UNIV INDONESIA**Publisher Address:** DIRECTORATE RESEARCH & PUBLIC SERV, UI CAMPUS, KAMOUS UNIV INDONESIA, DEPOK, 16424, INDONESIA**Web of Science Categories:** Medicine, Research & Experimental**Research Areas:** Research & Experimental Medicine**IDS Number:** HW9WZ**ISSN:** 2356-3664**eISSN:** 2356-3656**29-char Source Abbrev.:** MAKARA J HEALTH RES**ISO Source Abbrev.:** MAKARA J. HEALTH RES.**Source Item Page Count:** 6**Funding:**

Funding Agency	Grant Number
International Islamic University Malaysia (IIUM Research Initiative Grant scheme)	RIGS 15-080-0080

International Islamic University Malaysia (IIUM Research Initiative Grant scheme, RIGS 15-080-0080).

Open Access: DOAJ Gold**Output Date:** 2019-07-31

Close

Web of Science
Page 1 (Records 1 -- 1)

Print

**Clarivate**

Accelerating innovation

© 2019 Clarivate

[Copyright notice](#)[Terms of use](#)[Privacy statement](#)[Cookie policy](#)[Sign up for the Web of Science newsletter](#)[Follow us](#)