



Scopus

Search Sources Lists SciVal ↗



Create account

Sign in

Document details

[Back to results](#) | [Previous](#) 2 of 345 [Next](#)
[Export](#) [Download](#) [Print](#) [E-mail](#) [Save to PDF](#) [Add to List](#) [More...](#)
[Full Text](#) [View at Publisher](#)

Saudi Journal of Biological Sciences [Open Access](#)
 Volume 26, Issue 7, November 2019, Pages 1519-1524

Lifestyle interventions for non-alcoholic fatty liver disease (Review) [\(Open Access\)](#)

Ahmed, I.A.^a Mikail, M.A.^b, Mustafa, M.R.^a, Ibrahim, M.^c, Othman, R.^a

^aThe Center for Natural Products Research and Drug Discovery (CENAR), Wellness Research Cluster, University of Malaya, Kuala Lumpur, 50603, Malaysia

^bDepartment of Biomedical Science, Faculty of Science, Lincoln University College, Malaysia

^cInternational Islamic University Malaysia, Kuantan Campus, Malaysia

Abstract

[View references \(54\)](#)

Non-alcoholic fatty liver disease (NAFLD) is a multi-factorial disease and the most common of chronic liver diseases worldwide. The four clinical-pathological entities which are usually followed by NAFLD course include non-alcoholic steatosis, non-alcoholic steatohepatitis, advanced fibrosis/cirrhosis, and hepatocellular carcinoma. The cornerstones of NAFLD management and treatment, however, are healthy lifestyles such as dietary modifications, regular physical activity, and gradual weight loss. At present, no drugs or pharmacological agents have been approved for long-term treatment of NAFLD. Therefore, lifestyle modification is considered the main clinical recommendation and an initial step for the management of NAFLD. © 2019 King Saud University

SciVal Topic Prominence


Topic: Fatty Liver | Liver Diseases | Non-alcoholic steatohepatitis

Prominence percentile: 99.849



Chemistry database information

Substances



Author keywords

Diet Exercise Lifestyle Liver disease Weight loss

Metrics [View all metrics](#)



PlumX Metrics

Usage, Captures, Mentions, Social Media and Citations beyond Scopus.

Cited by 0 documents

Inform me when this document is cited in Scopus:

[Set citation alert >](#)
[Set citation feed >](#)

Related documents

Phenolic-rich baccarea angulata modulates inflammatory biomarkers of atherosclerosis

Ibrahim, M., Mikail, M.A., Ahmed, I.A. (2018) *Journal of Nutrition and Metabolism*

Improvements in clinical characteristics of patients with non-alcoholic fatty liver disease, after an intervention based on the Mediterranean lifestyle: A randomised controlled clinical trial

Katsagoni, C.N., Papatheodoridis, G.V., Ioannidou, P. (2018) *British Journal of Nutrition*

Treatment of Non-Alcoholic Steatohepatitis : from dietary means to bariatric surgery | Traitement de la stéatohépatite non-alcoolique: Des mesures hygiéno-diététiques à la chirurgie bariatrique

Lassailly, G., Gnemmi, V., Baud, G.

References (54)

[View in search results format >](#)

- 1 Ahmed, I.A., Mikail, M.A.
Paradigm shift: focusing on plant-based natural antimicrobials
(2017) *J. Microbiol. Experiment.*, 5 (2), pp. 1-2.
- 2 Ahmed, I.A., Mikail, M.A., Ibrahim, M.
Baccaurea angulata fruit juice ameliorates altered hematological and biochemical biomarkers in diet-induced hypercholesterolemic rabbits
(2017) *Nutrition Research*, 42, pp. 31-42. Cited 5 times.
www.elsevier.com/locate/nutres
doi: 10.1016/j.nutres.2017.04.012
[View at Publisher](#)
- 3 Ahmed, I.A., Mikail, M.A., Ibrahim, M.
Baccaurea angulata fruit juice protects rabbit's liver from hypercholesterolemia-induced injury
(2017) *Malaysian J. Med. Res.*, 1 (4), pp. 18-21. Cited 2 times.
- 4 Ahmed, I.A., Mikail, M.A., Bin Ibrahim, M., Bin Hazali, N., Rasad, M.S.B.A., Ghani, R.A., Wahab, R.A., (...), Yahya, M.N.A.
Antioxidant activity and phenolic profile of various morphological parts of underutilised Baccaurea angulata fruit
(2015) *Food Chemistry*, 172, pp. 778-787. Cited 26 times.
www.elsevier.com/locate/foodchem
doi: 10.1016/j.foodchem.2014.09.122
[View at Publisher](#)
- 5 Ahmed, M.
Non-alcoholic fatty liver disease in 2015 [\(Open Access\)](#)
(2015) *World Journal of Hepatology*, 7 (11), pp. 1450-1459. Cited 88 times.
<http://www.wjgnet.com/1948-5182/pdf/v7/i11/1450.pdf>
doi: 10.4254/wjh.v7.i11.1450
[View at Publisher](#)
- 6 Bambha, K., Wilson, L.A., Unalp, A., Loomba, R., Neuschwander-Tetri, B.A., Brunt, E.M., Bass, N.M.
Coffee consumption in NAFLD patients with lower insulin resistance is associated with lower risk of severe fibrosis
(2014) *Liver International*, 34 (8), pp. 1250-1258. Cited 38 times.
<http://www.wiley.com/bw/journal.asp?ref=1478-3223>
doi: 10.1111/liv.12379
[View at Publisher](#)

[View all related documents based on references](#)

Find more related documents in Scopus based on:

[Authors >](#) [Keywords >](#)